# FINAL REPORT

Investigating Potential Risk to Threatened and Endangered Species from Per- and Polyfluoroalkyl Substances (PFAS) on Department of Defense (DoD) Sites

SERDP Project ER18-1626

**JUNE 2020** 

Jamie Suski

EA Engineering, Science, and Technology, Inc., PBC

Distribution Statement A
This document has been cleared for public release





This report was prepared under contract to the Department of Defense Strategic Environmental Research and Development Program (SERDP). The publication of this report does not indicate endorsement by the Department of Defense, nor should the contents be construed as reflecting the official policy or position of the Department of Defense. Reference herein to any specific commercial product, process, or service by trade name, trademark, manufacturer, or otherwise, does not necessarily constitute or imply its endorsement, recommendation, or favoring by the Department of Defense.



#### REPORT DOCUMENTATION PAGE

Form Approved OMB No. 0704-0188

The public reporting burden for this collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing the burden, to Department of Defense, Washington Headquarters Services, Directorate for Information Operations and Reports (0704-0188), 1215 Jefferson Davis Highway, Suite 1204, Arlington, VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to any penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.

PLEASE DO NOT RETURN YOUR FORM TO THE ABOVE ADDRESS.

1. REPORT DATE (DD-MM-YYYY)	2. REPORT TYPE		3. DATES COVERED (From - To)	
30/06/2020	SERDP Final Report		6/25/2018 - 6/24/2021	
TITLE AND SUBTITLE  Investigating Potential Risk to Threatened and Endangered Species from Per- and			5a. CONTRACT NUMBER 18-C-0080	
Polyfluoroalkyl Substances (PFAS) on		5b. Gl	5b. GRANT NUMBER	
			ROGRAM ELEMENT NUMBER	
6. AUTHOR(S) Jamie Suski		ER18-1		
			ASK NUMBER	
		5f. Wo	ORK UNIT NUMBER	
7. PERFORMING ORGANIZATION N EA Engineering, Science, and Techno 225 Schilling Circle, Suite 400 Hunt Valley, Maryland 21031			8. PERFORMING ORGANIZATION REPORT NUMBER 6333101	
<ol> <li>SPONSORING/MONITORING AGE Strategic Environmental Research and 4800 Mark Center Drive, Suite 16F16 Alexandria, VA 22350-3605</li> </ol>	( )		10. SPONSOR/MONITOR'S ACRONYM(S) SERDP	
Alexandra, VA EEGGG GGGG			11. SPONSOR/MONITOR'S REPORT NUMBER(S) ER18-1626	
12. DISTRIBUTION/AVAILABILITY S	STATEMENT			
DISTRIBUTION STATEMENT A. Appr	oved for public release: distribution unlimited.			
13. SUPPLEMENTARY NOTES				

#### 14. ABSTRACT

Federal and State threatened and endangered fauna species occupy many Department of Defense (DoD) sites. The diversity of the species is great ranging from invertebrates to higher trophiclevel predators. Importantly, many DoD sites are also impacted by historical activities that result in chemical contamination. One of those activities was widespread use of Aqueous Film Forming Foams (AFFF) to suppress fires and in fire training exercises that has resulted in Per- and Polyfluoroalkyl Substances (PFAS) distributed in many environmental compartments.

A species-specific exposure potential to PFAS was developed to provide an initial screen of impacted habitat using spatial analysis. Protected species (or taxa) can then be ranked by those that occupy the greatest proportion of PFAS impacted habitat. Those species may be prioritized for hazard and/or risk assessment. Here, we carry two species (northern pine snake and bald eagle) through to develop a probabilistic risk framework on site specific PFOS concentrations.

#### 15. SUBJECT TERMS

Potential Risk, Threatened and Endangered Species, Per- and Polyfluoroalkyl Substances, PFAS, Department of Defense, DoD

16. SECURITY CLASSIFICATION OF:		17. LIMITATION OF 18. NUMBER		19a. NAME OF RESPONSIBLE PERSON Jamie Suski	
a. REPORT	b. ABSTRACT	c. THIS PAGE	ABSTRACT	OF PAGES	Janne Suski
UNCLASS	UNCLASS	UNCLASS	UNCLASS	142	<b>19b. TELEPHONE NUMBER</b> (Include area code) 410-584-7000

This page intentionally left blank

June 2020

# **TABLE OF CONTENTS**

				<u>Pa</u>	age
	OF ACE			ABBREVIATIONS	
EXEC 1.					
2. 3.				ETHODS	
	3.1	PROT	ECTED S	SPECIES ON DOD SITES	7
		3.1.1	Species	List Development	7
			3.1.1.1 3.1.1.2 3.1.1.3	Joint Base McGuire-Dix-Lakehurst  Naval Air Station Patuxent River  Barksdale Air Force Base	7
		3.1.2	Species	Habitat Designation	8
	3.2	DESK	TOP GIS	ANALYSIS	8
		3.2.1	Existing	Data	8
			3.2.1.1 3.2.1.2 3.2.1.3	Joint Base McGuire-Dix-Lakehurst Naval Air Station Patuxent River Barksdale Air Force Base	9
		3.2.2 3.2.3	Data Re	s Methodology for Spatial Exposure Potentialliability and Limitations for the Spatial Exposure Potential	
			3.2.3.1 3.2.3.2 3.2.3.3	Joint Base McGuire-Dix-Lakehurst  Naval Air Station Patuxent River  Barksdale Air Force Base	11
		3.2.4	Threater	ned and Endangered Species GIS Data Layers	.12
4.	RESU	LTS A	ND DISC	USSION	13
	4.1	SPAT	IAL ANA	LYSIS OF PFAS EXPOSURE POTENTIAL	13
		4.1.1 4.1.2 4.1.3	Naval A	se McGuire-Dix-Lakehurstir Station Patuxent River	.19

EA Project No.: 6333101 SERDP Project No. ER.18-1626 Version: FINAL June 2020

# EA Engineering, Science, and Technology, Inc., PBC

	4.2	SPEC	IES-SPECIFIC PROBABILISTIC EXPOSURE MODELS22	,
			Probabilistic exposure and risk framework for <i>Pituophis melanoleucus</i> 23 Probabilistic exposure and risk framework for <i>Haliaeetus leucocephalus</i> 30	
5. 6.			ONS AND IMPLICATIONS FOR FUTURE RESEARCH	

June 2020

# LIST OF FIGURES

<u>Number</u>	<u>Title</u>
Figure 1	Percent of AFFF impacted habitat by protected taxa on JBMDL
Figure 2	Percent of protected taxa-specific landcover estimated to be impacted by AFFF by base
Figure 3	Percent of estimated AFFF impacted landscape of protected avian species by dietary requirements
Figure 4	Percent impacted habitat by taxa on NAS Patuxent River before and after GIS updates
Figure 5	Daily energy consumed for an individual snake. Consumption is not constant because snakes consume whole prey which can exceed their daily energy requirements
Figure 6	Estimated PFOS exposure for individual snakes with a probability of consuming PFOS-contaminated prey of 0.5 (top) and 0.1365 (bottom)
Figure 7	Log10 estimated PFOS exposure for 1000 snakes exposed for 90 days
Figure 8	Post-hatch growth of eaglets until fledging ~62-days.
Figure 9	Realization of eaglet ingestion rate through the 60-day developmental period.
Figure 10	Contours are 0.5km and extend East from Cooper Bayou.
Figure 11	Representative PFOS dietary exposure to post-hatch eaglet.
Figure 12	Estimate of PFOS risk to eaglets at Barksdale Air Force Base.
	Appendix C
Figure C-1	Habitat Types – Lakehurst (JBMDL)
Figure C-2	Habitat Types – McGuire-Dix (JBMDL)
Figure C-3	Habitat Types – Pre-site Visit (NAS Patuxent River)
Figure C-4	Habitat Types – Post Site Visit (NAS Patuxent River)
Figure C-5	Habitat Types – Original (Barksdale AFB)
Figure C-6	Habitat Types (Original) – Area of AFFF Release (Barksdale AFB)
Figure C-7	Habitat Types - Updated (Barksdale AFB)
Figure C-8	Habitat Types (Updated) – Area of AFFF Release (Barksdale AFB)
Figure C-9	Habitat Impacted for <i>Pituophis melanoleucus</i> – McGuire (JBMDL)
Figure C-10 Figure C-11	Habitat Impacted for <i>Haliaeetus leucocephalus</i> – McGuire (JBMDL) Habitats Impacted for <i>Haliaeetus leucocephalus</i> – Revised layer
riguie C-11	(Barksdale AFB)

# LIST OF TABLES

<u>Number</u>	<u>Title</u>		
Table 1	Estimated landcover impacted by AFFF migration from release locations		
Table 2	Landcover type by base that has $\geq 10\%$ of available habitat impacted by AFFF migration		
Table 3	Protected Species Exposure Potential to PFAS by base on JBMDL		
Table 4	Protected Species Exposure Potential to PFAS on NAS Patuxent River		
Table 5	Exposure model parameters for <i>P. m. melanoleucus</i>		
	Appendix A		
Table A-1	Estimated exposure potential of landscapes at Joint Base McGuire Dix Lakehurst		
Table A-2	Estimated exposure potential of landscapes at NAS Patuxent River		
Table A-3	<u> </u>		
	Appendix B		
Table B-1	Spatial exposure potential of protected species at McGuire of Joint Base McGuire-Dix-Lakehurst		
Table B-2	Spatial exposure potential of protected species at Dix of Joint Base McGuire-Dix- Lakehurst		
Table B-3	Spatial exposure potential of protected species at Lakehurst of Joint Base McGuire-Dix-Lakehurst		
Table B-4	Spatial exposure potential of protected species at Naval Air Station Patuxent River		

SERDP Project No. ER.18-1626 Version: FINAL June 2020

#### LIST OF ACRONYMS AND ABBREVIATIONS

ac acre

ADI Average Daily Intake
AF Assessment Factors
AFB Air Force Base

AFFF Aqueous Film Forming Foams

AUF Area Use Factor

BAFB Barksdale Air Force Base

BSAF Biota-to-Soil-Accumulation Factors

C-F bond Carbon-Fluorine Bond DoD Department of Defense DOE Department of Energy

Dom Dominated

EPA Environmental Protection Agency

ERSON Environmental Restoration Statement of Need

ESA Endangered Species Act

ESTCP Environmental Security Technology Certification Program

FFTS Former Fire Training Area FMR Field Metabolic Rate

GIS Geographic Information System

INRMP Integrated Natural Resource Management Plans
IPaC Information for Planning and Consultation

JBMDL Joint Base McGuire Dix Lakehurst

LC50 Lethal Concentration for 50 Percent of Organisms

LiDAR Light Detection and Ranging

LOAEL Lowest Observable Adverse Effect Level MD DNR Maryland Department of Natural Resources

MDP Maryland Department of Planning

NAS Naval Air Station

NJDEP New Jersey Department of Environmental Protection

NLCD National Land Cover Dataset

NRCS National Resources Conservation Service

NWI National Wetland Inventory PA Preliminary Assessment

PFAS Per- and Polyfluoroalkyl substances

PFHxS Perfluorohexanoic Acid PFOS Perfluorooctane sulfonate

BSAF Biota-to-Soil Accumulation Factor

sd Standard Deviation

SEEM Spatially-Explicit Exposure Model

SERDP Strategic Environmental Research and Development Program

SIR Site Investigation Report T&E Threatened and Endangered

EA Project No.: 6333101 SERDP Project No. ER.18-1626

Version: FINAL June 2020

# EA Engineering, Science, and Technology, Inc., PBC

TRV **Toxicity Reference Values** United States Department of Agriculture USDA United States Environmental Protection Agency **USEPA** United States Fish and Wildlife Service

**USFWS** 

United States Geological Survey **USGS** 

EA Project No.: 6333101 SERDP Project No. ER.18-1626 Version: FINAL

June 2020

EA Engineering, Science, and Technology, Inc., PBC

This page intentionally left blank

EA Project No.: 6333101 SERDP Project No. ER.18-1626 Version: FINAL

EA Engineering, Science, and Technology, Inc., PBC

June 2020

#### **ABSTRACT**

#### Introduction:

Federal and State threatened and endangered fauna species occupy many Department of Defense (DoD) sites. The diversity of the species is great ranging from invertebrates to higher trophic-level predators. Importantly, many DoD sites are also impacted by historical activities that result in chemical contamination. One of those activities was widespread use of Aqueous Film Forming Foams (AFFF) to suppress fires and in fire training exercises that has resulted in Per- and Polyfluoroalkyl Substances (PFAS) distributed in many environmental compartments. Objectives and Technical Approach:

A species-specific exposure potential to PFAS was developed to provide an initial screen of impacted habitat using spatial analysis. Protected species (or taxa) can then be ranked by those that occupy the greatest proportion of PFAS impacted habitat. Those species may be prioritized for hazard and/or risk assessment. Here, we carry two species (northern pine snake and bald eagle) through to develop a probabilistic risk framework on site specific PFOS concentrations. Results:

Investigating three DoD installations generally resulted in high exposure potential for those species that have smaller ranges (invertivores) compared to species that move great distances over the landscape (carnivorous mammals). Furthermore, exposure potential models provide a map that illustrates the overlap of critical habitat with areas of suspected PFAS contamination which is useful for natural resource managers in prioritizing environmental remediation. Results from the spatial exposure potential model and the statement of need (SON) requesting information on species that occupy multiple DoD sites, two species were selected for developing the risk framework; those were the state listed (New Jersey/Tennessee) northern pine snake and the federally protected bald eagle. These species were selected based on high exposure potential, occupancy on multiple installations and no available data on risk of PFAS to these receptors. Currently, there are no published data of PFAS risk to reptiles and importantly, there are many reptilian species that occupy DoD sites. Probabilistic risk frameworks were developed for the northern pine snake and the bald eagle that incorporates species-specific physiology and behavioral traits. Results of the risk model were compared to working Toxicity Reference Values (TRV) for each species. The risk frameworks focused on PFOS alone, as this is currently, the most frequently encountered and believed to be the most hazardous of the PFAS to wildlife. The probabilistic model indicates the northern pine snake is not likely to be affected even with 100% of dietary items captured from PFOS-contaminated habitats (~16 µg PFOS/kg soil) on Joint Base McGuire Dix Lakehurst (JBMDL). Conversely, the bald eagle model that focuses on the eaglet life-stage shows there may be risk resulting from PFOS exposures. Eaglets have small body-size with high energetic demands resulting in high energetic demands that drive exposure. Nests in closer proximity to PFAS-impacted water bodies are assumed to receive a greater proportion of PFOS-contaminated prey items. With 10% of prey items from an impacted water body, the eaglet model shows the TRV is exceeded regularly over the course of development to fledging (60-days). Although the models are comprehensive, further information on exposure estimates from field research would be useful in refining the probabilistic risk estimates. In addition, given the high exposure potential to invertebrates and invertivores, toxicity data and exposure/risk modeling to these receptors is needed. Benefits:

EA Project No.: 6333101 SERDP Project No. ER.18-1626

Version: FINAL

EA Engineering, Science, and Technology, Inc., PBC

June 2020

This research benefits Department of Defense installations by applying a rapid screening-level approach to determine which wildlife receptors may be at greatest risk of exposure. Given the current limitations on toxicity (or hazard) information addressing exposure may be a more rapid route to take in determining if wildlife receptors are expected to come into contact with PFAS impacted habitat. Furthermore, the probabilistic risk framework developed for reptiles and avian predators illustrate an initial path forward in conducting full ecological risk assessments on DoD sites.

EA Project No.: 6333101 SERDP Project No. ER.18-1626 Version: FINAL

June 2020

EA Engineering, Science, and Technology, Inc., PBC

This page intentionally left blank

EA Engineering, Science, and Technology, Inc., PBC

#### **EXECUTIVE SUMMARY**

#### 1. Introduction:

Per- and Polyfluoroalkyl Substances (PFAS) are emerging contaminants of concern and are found on many military installations that have airborne missions. These chemicals are extremely persistent and somewhat mobile in environmental media and have been measured in nearly all environmental compartments and biota (e.g. Houtz et al. 2013, Long and Porter, 2015, Geisy and Kannnan 2001, Custer et al. 2014). As such, there is growing interest regarding the potential effects of PFAS to public and environmental health, as well as concern from the Department of Defense (DoD) regarding liability under federal statutes such as the Endangered Species Act (ESA). The predominant source of PFAS on DoD sites originates from historical fuel-based fire suppression activities using Aqueous Film Forming Foam (AFFF) which contained a mixture of largely perfluorooctane sulfonate (PFOS) in combination with other PFAS (Houtz et al. 2013). The physicochemical properties that render PFAS compounds effective in fire suppression also translate to resistance to biodegradation, photooxidation, direct photolysis and hydrolysis.

Studies conducted by DoD and researchers at U.S. Air Force installations confirm PFAS detections in environmental compartments and transportation of PFAS from sites below AFFF training areas to surface water, sediment and groundwater (Lanza et al. 2017, Moody et al. 2002, Gewurtz et al. 2014, Karrman et al. 2011, Oakes et al. 2010). Biota may be exposed via direct and indirect pathways; such as, direct contact with soil or drinking water or bioaccumulation. Identifying exposure pathways to ecological receptors can be complex; however, determining risk may prove more challenging based on overall lack of toxicity data on many PFAS.


Given the lack of information concerning the potential risk of PFAS to federal and state listed Threatened &Endangered (T&E) species on DoD lands, we developed a framework to determine T&E species occurrence and potential overlap with PFAS contamination. This project provides immediate and useful insights into the potential for PFAS exposure to T&E species and points the way toward methods for location and species-specific assessments for PFAS risk.

## 2. Objectives:

The overall objective of this research was to develop a framework for natural resource managers to quickly assess on-site threatened and endangered species likelihood of PFAS exposure and risk. To address this objective, we developed (1) methodology for determining spatial overlap of T&E species on DoD sites with areas of AFFF release, (2) prioritization for T&E species with greatest exposure potential and (3) conducted a species-specific probabilistic risk assessment for T&E species to determine risk of PFOS exposure.

### 3. Technical Approach:

The technical approach followed the schematic below in two main phases with two decision points (as indicated by stars). Phase 1 (silver box) was used to develop the T&E (this point forward referred to as 'protected') species exposure potential on installations and Phase 2 (blue box) is the determination and/or development of the individual based probabilistic models. The decision points were used at particular junctures to (1) determine if there are enough data to move forward with generating the spatial exposure potential and (2) determine if there are any protected species that warrant a probabilistic exposure assessment to determine a risk.



Phase 1: Exposure Potential Modeling

To develop the spatial exposure potential modeling, installations with known AFFF release locations and environmental PFAS concentrations in the form of a Site Investigation Report (SIR) or published literature were identified. Using Geographic Information System (GIS), landscape layers were obtained from the installation or other publicly available sources, AFFF release locations from repeated fire training exercises or single use were digitized from the SIR creating a base map of land cover layers (habitat) and AFFF release sites. Transport of PFAS from AFFF release sites is expected, and therefore, using landscape topography, known conveyances and surface water flow direction areas impacted by AFFF and PFAS have been extrapolated out from the release locations terminating at the nearest surface water body. Those extrapolations cross landscape layers that allow for estimation of habitat specific PFAS impacts from AFFF release sites.

Site specific protected species were identified using the Integrated Natural Resource Management Plans (INRMP) and Federal/State/County T&E species lists. Habitat requirements

EA Engineering, Science, and Technology, Inc., PBC

of each species were identified through literature searches and were matched with the available landscape layers on the base map in GIS to determine where those species may be located on the installation. Furthermore, species-specific dietary requirements were also incorporated into the spatial model to capture potential foraging areas.

#### Phase 2: Risk Assessment

Risk Assessment of two protected species were conducted. The bald eagle (*Haliaeetus leucocephalus*) and the northern pine snake (*Pituophis melanoleucus*) were selected based on the spatial exposure potential of the three installations evaluated and a broad application to multiple installations. Although the bald eagle is delisted from the Endangered Species Act (ESA) it continues to be a protected species under the Bald and Golden Eagle Protection Act (1940), the Migratory Bird Treaty Act (1918) and the Lacy Act (1900), given the number of laws in place to protect this species and the broad distribution of bald eagles on coastal installations it was prioritized for the risk assessment. The second species, the northern pine snake was selected because the exposure potential was relatively high compared to other protected taxa on Joint Base McGuire-Dix-Lakehurst (JBMDL), many DoD installations have one or more protected reptilian species and there is a significant gap in reptilian risk assessment (Weir et al. 2010) especially pertaining to PFAS (Conder et al. 2019). The northern pine snake is also a federally protected species at Arnold Air Force Base in Tennessee (Petersen et al. 2017).

A probabilistic exposure and risk model were used to estimate PFAS exposure to the bald eagle and the northern pine snake. The models were based on allometric equations to predict field metabolic rate (energy needs) in avian and reptilian receptors. For the bald eagle, an exposure model was developed, assuming a realistic diet of mammal and fish with a probability of PFOS contamination using published PFOS concentrations (Chang et al. 2012, Lanza et al., 2017). The study focused on the eaglet (day 0 -60, fledgling) as this may be the most vulnerable life stage because of the disproportionally high ingestion rates compared to body weights. Model simulations were comprised of 1000 replicate eaglets and the mean exposure over the developmental period was the average daily intake (ADI). The 1000 replicate estimates of ADI were averaged and compared to the Toxicity Reference Value (TRV) for raptors of 0.021 mg/kg/d (Newsted et al. 2005).

The probabilistic exposure model for the northern pine snake followed a similar construct to the bald eagle model. Where a probability of capturing PFOS-contaminated prey items, feeding frequency and PFOS concentration was incorporated into the exposure model. Model results were also comprised of 1000 simulation runs of individual snakes foraging for 90-days; the resulting mean PFOS exposure was the ADI. The overall mean ADI for snakes with varied exposure potential to PFOS was compared to the working PFOS TRV for reptiles of 2 mg/kg/d (Salice et al. *In prep*).

#### 4. Results and Discussion:

Results indicate that a screening-level spatial exposure potential assessment will help installations identify protected species that are likely to be exposed to PFAS through historical

EA Engineering, Science, and Technology, Inc., PBC

AFFF release. Moreover, this analysis provides a ranking of impacted landscapes and thus those protected species dependent on landscapes ranked higher may have comparably greater exposure potential to those on landscapes ranked lower. Broadly, this tool will aid in identifying habitat that supports non-protected ecological receptors as there is likely overlap in habitat selection by taxa. Using the exposure mapping also provides a spatial component for natural resource managers to visualize areas of potential concern based on habitat type.

Protected species at Joint Base McGuire-Dix-Lakehurst, Naval Air Station Patuxent River and Barksdale Air Force Base were prioritized by impacted habitat exposure potential by taxa, taxa and diet and individual species. Results indicate taxa with small home ranges have the greatest exposure potential as those that travel further are able to move across the landscape increasing their habitat range. Whereas the inverse is true for taxa with small home (i.e. invertebrates) ranges should their habitat be limited to an impacted area. In addition to providing a framework for ranking species-specific exposure potential our aim was to conduct full risk estimates on two species that were identified as having high exposure potential.

The risk estimates for the eaglet life-stage of the Bald Eagle indicate there may be significant risk from dietary PFOS exposures. The probabilistic model developed indicates that the life stage from hatching to fledgling (60-days) may be the most sensitive compared to adult exposures due to the high metabolic rate and smaller body size. As such, eaglets can receive a high PFOS exposure potential in dietary items from prey captured daily by parent eagles. Using available PFOS concentrations in fish at Barksdale AFB (Lanza et al. 2017), eaglets may receive PFOS exposures greater than the currently available TRV of 0.021 mg/kg/d on a daily basis. In comparison, the preliminary modeled exposure for the northern pine snake suggests PFOS exposures do not exceed the working TRV of 2 mg/kg/d for reptiles.

The spatial exposure potential assessment is dependent on a few key factors. First, accurate and available landscape layers in GIS is critical at identifying appropriate habitat suitable for protected species on the installation. Secondly, the spatial exposure analysis is limited to reported AFFF release locations. To account for environmental movement of PFAS we extrapolated the area of impact out from the known release site following topography, groundwater and surface water conveyances. Therefore, without environmental sampling of PFAS the true migration is unknown and thus we provide best estimation of migration. Furthermore, there may be other sites of AFFF release and measurable PFAS concentrations that are unknown or not available for this effort.

#### 5. Implications for Future Research and Benefits:

There are some excellent opportunities for future research that have become apparent during this effort. To address uncertainties in extrapolated PFAS migration within the landscape environmental sampling along a gradient or at the expected surface water termination would be beneficial for proof-of-concept. Protected species are ranked based on the exposure potential which is a critical first step in understanding which species and/or taxa may be receiving the greatest exposure to PFAS. We acknowledge there are limitations to this based on historical

EA Engineering, Science, and Technology, Inc., PBC

sampling of known AFFF release locations, however, using those limited data, environmental sampling could fill in the gaps of expected PFAS transport.

In addition to environmental sampling, <u>a critical need</u> in understanding true ecological receptor PFAS exposure scenarios is sampling of biota in concert with environmental compartment sampling. This is especially true for the reptilian model as there are very few (if any) data available for PFAS tissue concentrations. Our model suggests the northern pine snake may occupy a PFOS impacted site; however, exposures may not elicit an apical response. Sampling blood or tissue of a surrogate reptilian species would be notably informative to this model. Importantly, these data should be collected in conjunction with animal tagging so true PFAS exposure can be ascertained based on a temporal and spatial scale. Other avenues of research to determine exposure would be sampling of shedded reptilian skin as depuration of contaminants have been previously noted via this route of elimination. These data would then be incorporated into the snake model to refine expected reptile PFAS exposure scenarios.

Finally, and perhaps most important would be developing a field component to determining PFOS exposures to bald eagle eaglets. As the current probabilistic model suggests there may be risk to eaglets through high dietary exposures. We assumed proximity of nest location to PFAS impacted sites plays a role in the eaglet exposure profile and would suggest nest sampling at near and far distances including nests that are used in consecutive years. One non-lethal approach to determining PFOS exposure to eaglets would be to sample feathers that have been shed, this approach also results in minimum disturbance to the animals. In addition to feather sampling, droppings may also be sampled for PFOS, the concern with this approach alone is that fecal material may be from parent or the eaglet. Whereas feathers can be more easily identified as adult or juvenile.

The abovementioned additional effort could serve as a 'ground truth' to our models developed under this statement of need. Moreover, the risk assessment community frequently requests more information on observed field effects to those modeled and/or inferred by environmental sampling alone.

### 1. OBJECTIVES

The technical objectives of this research were developed in response to Strategic Environmental Research and Development Program's (SERDP), Environmental Restoration Statement Of Need (ER SON)-18-L2 that requested an approach for assessing risk of PFAS to federal and state threatened and endangered (T&E) species. The SON highlighted the need to assess PFAS risk of shared T&E species among installations and development of PFAS exposure factors. Our approach involved a thorough investigation of the available data on T&E species occurrence, life-histories, and habitat requirements to identify the likelihood and routes of potential PFAS exposure to prioritize T&E species for further risk assessment. Upon identification of those species with the greatest exposure potential probabilistic exposure models were developed on two candidate species. The objects and associated tasks are broken down as follows:

Objective 1: Develop Species-specific spatial PFAS exposure estimates on impacted Department of Defense (DoD) sites

- Task 1: Identification of DoD installations that are or may be contaminated with PFAS resulting from the historical use of Aqueous Film Forming Foam (AFFF) in fire suppression activities
- Task 2: Determine Locations, Occurrences, and Exposure of Rare, Threatened or Endangered Species on DoD Installations identified in Task 1 as PFAS-contaminated (or likely contaminated). Geographic Information System (GIS) geodatabase data layers and taxa specific exposure potentials will be used to rank T&E species for further investigation

Objective 2: Develop tiered assessment approach for T&E species risk assessments

- Task 3: Synthesize outcomes from Objective 1 into an overall conceptual framework for assessing risk to protected species that inhabit PFAS-contaminated installations
- Task 4: Develop two probabilistic exposure models for identified protected species from the tasks above

EA Project No. 6333101 SERDP Project No. ER.18-1626 Version: FINAL

June 2020

EA Engineering, Science, and Technology, Inc., PBC

This page intentionally left blank

#### 2. BACKGROUND

Per- and Polyfluoroalkyl Substances (PFAS) are emerging contaminants of concern and are found on many military installations that have airborne missions. These chemicals are extremely persistent and somewhat mobile in environmental media and have been measured in drinking water (Hu et al. 2016), groundwater (Long and Porter, 2015, Houtz et al. 2013), surface water (Long and Porter, 2015, Awad et al. 2011), soils and sediment (Long and Porter, 2015; Higgins and Luthy 2007) and a wide variety of biota (Fish, Geisy and Kannnan 2001, Hoff et al. 2005; Birds, Kannan et al. 2001, Custer et al. 2014; Mammals, Falandysz et al. 2007). As such, there is growing interest regarding the potential effects of PFAS to public and environmental health, as well as concern from the Department of Defense (DOD) regarding liability under federal statutes such as the Endangered Species Act (ESA). The predominant source of PFAS on DOD sites originates from historical fuel-based fire suppression activities using Aqueous Film Forming Foam (AFFF) which contained a mixture of largely perfluorooctane sulfonate (PFOS) in combination with other PFAS (Houtz et al. 2013). The physicochemical properties (i.e. smothering surfactant layer) that render PFAS compounds effective in fire suppression also translate to resistance to biodegradation, photooxidation, direct photolysis and hydrolysis. Specifically, while carbon chain length and terminal group vary among PFAS, all share a carbon backbone that is either fully saturated with Fluorine (perfluoroalkyl substance) or partially saturated with Florine (polyfluoroalkyl substance) (Buck et al. 2011). The C-F bond is especially strong and imparts the salient properties for PFAS chemicals. AFFF has been used by the military since the 1960s, though the exact PFAS mixture that comprised the AFFF used over the years changed with respect to supplier. It has been estimated, however, that the 3M mixture was used predominantly since the 1980s and in 2004, 75% of the military stockpile was produced by 3M (Darwin 2004). Given the persistent nature of PFAS, continuous exposure occurs even though AFFF use in fire suppression and training exercises has been reduced. For example, PFAS detections in sediment and surface water near an accidental release of AFFF from a malfunctional fire alarm at L.B. Pearson Toronto International Airport in 2000 were detected nine years later (Awad et al. 2011). In addition to PFAS detections in multiple environmental compartments, PFAS are found within blood and tissue of humans and wildlife (Apelberg et al. 2007, Geisy and Kannan 2001, Kannan et al. 2004, Kannan et al. 2006). Accordingly, concentrations of PFAS in environmental compartments and in biota near the Toronto Airport release decreased with distance from the epicenter of release (Awad et al. 2011). Importantly, this highlights the ubiquitous nature of this class of chemicals and the need to further understand how they may be impacting wildlife, especially imperiled species. Studies conducted by DOD and researchers at U.S. Air Force installations confirm PFAS detections in environmental compartments and transportation of PFAS from sites below AFFF training areas to surface water, sediment and groundwater (Lanza et al. 2017, Moody et al. 2002, Gewurtz et al. 2014, Karrman et al. 2011, Oakes et al. 2010). Biota may be exposed via direct and indirect pathways; such as, direct contact with soil or drinking water or bioaccumulation. Identifying exposure pathways to ecological receptors can be complex; however, determining risk may prove more challenging based on overall lack of toxicity data on many PFAS.

Most studies investigating ecotoxicology of PFAS have focused on PFOS (Giesy et al. 2010, Ding et al. 2013) and PFOA (Hekster et al. 2003) within the aquatic environment.

EA Engineering, Science, and Technology, Inc., PBC

Unfortunately, ecotoxicity data are significantly lacking with regard to terrestrial organisms and on the toxicity of other PFAS in general, such as Perfluorohexane sulfonic acid (PFHxS) and Perfluorohexanoic acid (PFHxA), which are routinely found in surface water, sediment, and soils together on DOD installations with AFFF contamination (Long and Porter 2015). Should taxa specific ecotoxicity data be available it does not necessarily reflect the classic taxa-exposure-response relationship, for example, Custer et al. 2014 found effects in hatching success of tree swallows correlated to PFOS concentrations (283 ng/g ww) lower than proposed Avian Toxicity Reference Values (TRV; 600 ng/g ww [Northern Bobwhite Quail and Mallard], Newsted et al. 2005). This highlights the importance of considering species-specific differences and life-history traits (e.g. foraging range) when quantifying exposure and risk scenarios to Threatened & Endangered (T&E) species.

Section 7(a)(2) of the ESA states "each federal agency shall, in consultation with the Secretary, insure that any action authorized, funded, or carried out by such agency is not likely to jeopardize the continued existence of threatened and endangered species or destroy or result in the destruction or adverse modification of habitat of such species which is determined by the Secretary...". Therefore, putting both proactive and responsive actions on federal agencies with respect to protection of T&E species. The United States Environmental Protection Agency (USEPA) conducts risk assessments on T&E species for unintentional pesticide exposures; the T&E risk assessments are species-specific and the majority are on Rana draytonii (California Red-legged frog; 39% of total assessments). The US Environmental Protection Agency (US EPA) frequently estimates exposures using allometric equations for food and water consumption from the Wildlife Exposures Factors Handbook (USEPA 1993); although dated at this point, the wildlife exposures factors handbook is still comprehensive in providing estimates of food and water ingestion rates for terrestrial and aquatic receptors through drinking water or diet. Commonly incorporated into protected species risk assessments are assessment factors (AF), which reflect confidence in the data being used, i.e. soundness of the study and relevance of the data to the assessment (USEPA 2003). For acute data, the AF is applied by multiplying the lethal concentration by 0.5; this is very close to the estimate of 0.63 that was derived from acute toxicity tests using standard models and endangered species (Dwyer et al. 2005). Dwyer et al. (2005) found that there were different sensitivities among species and there was not a dominantly sensitive species of the 20 tested. Importantly, PFAS exposures will not follow an acute duration and therefore, chronic exposures are most relevant to this research. As Part II of the Dwyer et al. (2005) research, Besser et al. (2005) conducted 30-day exposures on two commonly used test species and two protected species expanding the investigation on comparing sensitivities of common test species to those protected using two toxicants. The endangered fountain darter (Etheostoma fonticola) was the most sensitive to both toxicants compared to the two common and one endangered species (Besser et al. 2005). As expected, Besser et al. (2005) found that non-lethal endpoints were more sensitive than survival. Given the increased sensitivity of an endangered species to toxicant exposure compared to common laboratory models the inclusion of assessment factors for chronic studies is appropriate and likely the commonly used approach to protecting species on high concern.

Given the lack of information concerning the potential risk of PFAS to T&E species on DoD lands, we proposed to compile and synthesize data on species occurrence and potential overlap

Version: FINAL

EA Engineering, Science, and Technology, Inc., PBC

June 2020

with PFAS contamination and to outline a tiered approach to assess risk to T&E species with specific suggestions for data needs, as well as uncertainties related to each tier of the approach. This project provides immediate, useful insights into the potential for PFAS exposure to T&E species and points the way toward methods for location and species-specific assessments for PFAS risk.

EA Project No. 6333101 SERDP Project No. ER.18-1626 Version: FINAL

June 2020

EA Engineering, Science, and Technology, Inc., PBC

This page intentionally left blank

#### 3. MATERIALS AND METHODS

#### 3.1 PROTECTED SPECIES ON DOD SITES

Pursuant to the provisions of the Sikes Act (16 United States Code §670a et seq.), DoD is required to conserve and protect natural resources present on DoD lands. The Sikes Act requires that the DoD develop and implement Integrated Natural Resources Management Plans (INRMPs) for installations with natural resources conservation programs. These INRMPs serve as planning documents for an installation's management of natural resources in a manner that supports the military mission while promoting stewardship of natural resources and compliance with legal requirements. INRMPs also allow for stakeholder coordination and are developed and implemented in collaboration with the US Fish and Wildlife Service (USFWS) and State fish and wildlife agencies.

INRMPs provide detailed information and management strategies of federally- and state-protected species that are known to occur or have the potential to occur on an installation. The Sikes Act directs that INRMPs be reviewed and revised every 5 years. Installations with a recently updated INRMP provided an up-to-date species list for this research.

### 3.1.1 Species List Development

Protected species with potential and/or known occurrence at the study sites were developed to provide a geographically defined species list. Species-specific habitat type and ranges were determined through federal, state and/or published literature. These data provide the foundation for determining expected species locations on the study sites.

#### 3.1.1.1 Joint Base McGuire-Dix-Lakehurst

To establish expected protected species on JBMDL, the INRMP was compared against the New Jersey Pinelands Commission Threatened & Endangered Animals of the New Jersey Pinelands (Revised June 2015) as well as the USFWS Information for Planning and Consultation (IPaC) system to identify federal T&E species potentially at JBMDL.

#### 3.1.1.2 Naval Air Station Patuxent River

An updated INRMP was completed at Naval Air Station (NAS) Patuxent River in 2017 and includes a list of federal and state rare, threatened or endangered species with known or expected occurrence at the installation. Because this INRMP was developed in coordination with USFWS and provides site-specific species data, it was used to develop the species list for those expected to be found at NAS Patuxent River. In addition, the USFWS IPaC system was used to identify any federal species with the potential to occur on the installation.

#### 3.1.1.3 Barksdale Air Force Base

Like NAS Patuxent River, the INRMP was completed in 2017. The only protected species known to be present is the Bald Eagle (*Haliaeetus leucocephalus*). However, there is potential for the federally-listed northern long-eared bat as this species has been found during mist net surveys in Winn and Grant Parishes, which are southeast of the Air Force Base.

#### 3.1.2 Species Habitat Designation

Following species list development, species preferred habitat and home range information were collected from the New Jersey Department of Environmental Protection (NJDEP) Endangered and Non-Game Species Program website, Maryland Department of Natural Resources (MD DNR) website, USFWS IPaC system, NatureServe, and from the primary literature. Home range is defined here as the distance an organism may travel beyond the identified habitat preference.

#### 3.2 DESKTOP GIS ANALYSIS

A geospatial desktop analysis was conducted using a Geographic Information System (GIS) (ArcGIS version 10.4.1 or newer) to identify AFFF release sites, fate and transport pathways, and protected species receptor habitats. This approach provided a quasi-quantitative PFAS exposure potential to T&E species which can be used to rank species for further risk analysis.

#### 3.2.1 Existing Data

Data were obtained from the respective military installations and/or best available public data sources. Data gaps were identified if no publicly available data exists and/or the installation does not have sufficient data layers.

#### 3.2.1.1 Joint Base McGuire-Dix-Lakehurst

The following GIS data layers were obtained from JBMDL in 2018:

- Installation boundary, military ranges, training areas, buildings, impervious surfaces, and emergency response sites
- U.S. Geological Survey (USGS) streams and waterbodies
- U.S. Fish and Wildlife Service (USFWS) National Wetland Inventory (NWI) wetlands

#### Publicly Available Sources:

- New Jersey Department of Environmental Protection (NJDEP) (2012) land use/land cover
- U.S. Department of Agriculture (USDA) Natural Resources Conservation Service (NRCS) soils
- USGS and/or best available topography/Light Detection and Ranging (LiDAR) data

#### 3.2.1.2 Naval Air Station Patuxent River

The following GIS data layers were obtained from NAS Patuxent River in 2018:

- Installation boundary, military ranges, training areas, buildings, impervious surfaces, environmental remediation sites, monitoring well locations, and previous sampling locations
- Streams and waterbodies
- Wetlands

#### **Publicly Available Sources:**

- Maryland Department of Planning (MDP) for St. Mary's County (2010) land use/land cover
- USDA NRCS soils
- USGS and/or best available topography/LiDAR data

#### 3.2.1.3 Barksdale Air Force Base

The following GIS data layers were obtained from Barksdale AFB in 2019:

- Installation boundary, military ranges, road lines, restricted land use areas, environmental remediation sites, and previous sampling locations
- Prescribed burn and wildfire areas, natural resource survey areas, and bat survey areas
- Bald eagle habitat areas and nest locations
- Groundwater contours

#### Publicly Available Sources:

- USDA NRCS soils
- USGS streams and waterbodies
- USFWS NWI wetlands
- USGS National Land Cover Dataset (NLCD) (2011) land cover (2017)

#### 3.2.2 Analysis Methodology for Spatial Exposure Potential

Data were imported into a GIS framework from sources previously identified or were obtained and revised from field site visits (NAS Patuxent River). AFFF release areas were obtained from the installation if available or digitized from the Site Investigation Report (Aerostar SES LLC 2018 [JBMDL], Amec Foster Wheeler 2019 [Barksdale]) and Preliminary Assessment for NAS Patuxent River (CH2M 2018). Those sites included historical fire extinguishing events, fire stations, fire training areas and emergency response sites. Additionally, suspected AFFF release sites were also identified based on the SIR, for example, landfill locations. If exact location of release was known a point was used to identify that AFFF site; however, if a general area was reported a polygon was assigned.

AFFF drainage areas were created to map the horizontal extent of aboveground PFAS migration. Drainage areas were created by referencing the surface water pathways identified in the SIRs and Preliminary Assessment (PA) Reports and hydrologic interpretation of surface and groundwater

June 2020

EA Engineering, Science, and Technology, Inc., PBC

flow direction (HydroGeoLogic, Inc. 2015a, b [JBMDL, Barksdale, respectively], CH2M 2018 [NAS Patuxent R.]). Extents were developed to include AFFF release areas and downgradient areas to the connection of the closest surface water features (i.e. streams and ditches).

AFFF drainage areas were overlain with T&E species habitat to calculate PFAS exposure potential impacts to T&E species as a percentage of overall habitat available at each site. Habitat by species, taxa, diet, and taxa-diet were calculated using this approach.

#### 3.2.3 Data Reliability and Limitations for the Spatial Exposure Potential Estimate

The best available data for each site was variable based on scale, date of creation/collection, and methodology. To improve reliability, data layers were interpolated where data gaps were identified and/or collected from a field visit. Importantly, the spatial exposure potential estimates are not quantitative with respect to PFAS exposure concentrations within a given media (i.e. soils or surface water). The goal of this foundational step is to determine if exposure may occur within a species preferred habitat and if so, what is the percentage of habitat that may be impacted by PFAS.

#### 3.2.3.1 Joint Base McGuire-Dix-Lakehurst

The land use/land cover data layer was obtained from NJDEP for the entire State (2012). Aerial photointerpretation and a review of the data layer at the site demonstrated that the habitat data layer boundaries and land use/land cover types were at an acceptable resolution and detail for use in creating the T&E species habitat layer. These data appear to be at a fine scale with distinctions between many habitat features, such as differences in water body types and forest cover based on crown closure (Figure C-1 and C-2). Importantly, the ability to visually differentiate between the landscapes is not critical, figures C-1 and C-2 are included to provide an overview of the level of detail and complexity within the landscape mosaic.

Groundtruthing was conducted at Lakehurst (only, Figure C-1) to determine the accuracy of the obtained data layer. The data layers were refined with the results of the groundtruthing and further desktop analysis. Habitat types and boundaries were also revised using aerial photointerpretation. As a result, several habitat types were removed from one or more of the bases that combine to make JBMDL. These included: agricultural wetlands (modified), bridge over water, industrial, orchards/vineyards/nurseries/horticultural areas, other agriculture, residential rural single unit, and severe burned upland vegetation. "Phragmites dominant interior" habitat type was added to increase detail.

Due to the large size of McGuire and Dix (Figure C-2), habitat types were not groundtruthed; however, data layers were refined using aerial photointerpretation. As a result, several habitat types were removed from one or more bases. These included: altered lands, bridge over water, extractive mining, former agricultural wetland, industrial, major roadway, mixed residential, orchards/vineyards/nurseries/horticultural areas, wetland rights-of-way.

The data layers were further refined by integrating developed areas (i.e. buildings, airfields, roads, and sidewalks) and hydrolines (i.e. streams and canals) provided by the military installation.

#### 3.2.3.2 Naval Air Station Patuxent River

The land use/land cover data layer was obtained from MDP for St. Mary's County (2010). Aerial photointerpretation and a review of the data layer at the site revealed that the data layer boundaries and land use/land cover types were at an *unacceptable* resolution and detail for use in creating the T&E species habitat layer (Figure C-3). Comparatively, the habitat data layers are much coarser than those for JBMDL. The data layer boundaries were created at too low of a resolution, especially within the developed military areas, and a shoreline habitat type of a T&E shorebird species was absent.

Groundtruthing was conducted at NAS Patuxent River and those modifications were made to the data layers (Figure C-4). Also, habitat types and boundaries were revised using aerial photointerpretation. As a result, bare exposed rock, extractive, and industrial habitat types were removed and the varying qualifiers for residential density were consolidated into one habitat type called residential. Water was differentiated between nontidal and tidal water as species likely occupy one or the other unless euryhaline. Wetlands were differentiated between emergent and forested as canopy cover can influence species composition (Quesnelle et al. 2013). Sandy shoreline and old field habitat types were also added.

The data layer was further refined by integrating developed areas (i.e. buildings, airfields, and roads) and hydrology (i.e. streams and canals, waterbodies and wetlands) obtained from the military installation and publicly available data sources.

#### 3.2.3.3 Barksdale Air Force Base

The land use/land cover data layer was obtained from Barksdale AFB (2017); it represented a slightly revised version of the USGS National Land Cover Dataset (NLCD) (2011). Aerial photointerpretation and a review of the data layer at the site revealed that the data layer boundaries and land use/land cover types were at too course of resolution and detail for use in creating the T&E species habitat layer (Figures C-5&C-6). Comparatively, the habitat data layers are much coarser than those for JBMDL which was determined to be at an acceptable resolution for this analysis. Habitat types and boundaries were revised using aerial photointerpretation. As a result, the "barren land" habitat type was removed and open water was refined as lakes/ponds or streams/canals.

The data layer was further refined by integrating developed areas (i.e. buildings, airfields, and roads) and hydrology (i.e. streams and canals, waterbodies, and wetlands) through digitizing data and publicly available data sources. USFWS NWI wetlands were grouped into USGS NLCD wetland types for consistency (Figures C-7&C-8).

EA Engineering, Science, and Technology, Inc., PBC

June 2020

# 3.2.4 Threatened and Endangered Species GIS Data Layers

T&E species habitat layers were created by using the best available and corrected land use/land cover data layer and performing a tabular data join with the list of species that may inhabit each land use/land cover type.

Version: DRAFT
June 2020

#### 4. RESULTS AND DISCUSSION

Results of the spatial analyses are presented as (1) percent of impacted habitat and (2) percent of impacted habitat by taxa, dietary preference and individual T&E species. By incorporating the geographically influenced fate and transport of PFAS into the GIS analyses the likely pathways of chemical movement into defined landscapes become more apparent. Given this approach is limited to sampling locations from the Site Investigation Report (SIR) we believe it may be underestimating the total percent of impacted habitat. Furthermore, this approach was limited to the installation boundary, yet we know chemicals and wildlife may go beyond such boundaries.

#### 4.1 SPATIAL ANALYSIS OF PFAS EXPOSURE POTENTIAL

#### 4.1.1 Joint Base McGuire-Dix-Lakehurst

Using installation and habitat boundaries within GIS with the AFFF release and estimated PFAS migration pathway landscape layers we estimated the percent of impacted landscape on each base of JBMDL. The spatial analysis estimated 1,005 of the 42,151 available landcover acres (approximately 2.4%) are likely impacted by PFAS. The area of Dix has a disproportionate effect on this statistic given the large size (31,167 acres). Taken individually, results of expected PFAS contamination on each base are approximately 9.7, 1.4 and 3% of available landcover on McGuire, Dix and Lakehurst, respectfully (Table 1). Comparatively, we have estimated McGuire has the greatest percent of AFFF impacted landcover and Lakehurst has the greatest total estimated acres of impact (433 ac).

Table 1. Estimated landcover impacted by AFFF migration from release locations

Base	se Total Acres Impacted Acres		Percent Impacted (%)
McGuire	3,560	345	9.69
Dix	7,424	227	3.06
Lakehurst	31,167	433	1.39

For JBMDL there were 46 landcover types in GIS (Figure C-1&C-2). Twelve landcover types were identified as having a portion of impacted habitat across two or more bases (Appendix A). Additionally, landcover types that are expected to be impacted with approximately 10% or more of the available habitat on a base was identified and reported in Table 2 with the other bases of the installation for comparison.

June 2020

Table 2. Landcover type by base that has  $\geq 10\%$  of available habitat impacted by AFFF migration

Landcover Type	McGuire	Dix	Lakehurst
Military Installations	13.5	<10	8.0*
Stormwater Basin	10.1	<10	<10
Deciduous Forest (10-50% Crown Closure)	11.6	<10	<10
Old Field (< 25% Brush Covered)	18.7	<10	<10
Deciduous Brush/Shrubland	10.6	<10	<10
Artificial Lakes	19.5	<10	<10
Coniferous Wooded Wetlands	14.8	<10	<10
Herbaceous Wetlands	14.5	<10	<10
Disturbed Wetlands (Modified)	10.6	<10	<10
Transitional Areas	21.4	16.5	<10
Recreational Land	<10	26.7	<10
Agricultural Wetlands (Modified)	<10	98.8	n/a

<sup>\*</sup> greatest percent impacted landcover type on Lakehurst, n/a – removed from Lakehurst after identified as incorrect landscape cover for this base in GIS


Military Installation landcover type is a 'catchall' for landcover that does not fall into one of the listed categories and will include military fence lines and auxiliary land use. All other habitat layers are self-evident based on the name (i.e. Artificial Lake is a manmade lake).

Following the migration of AFFF through the landcover type, percent of impacted habitat specific to wildlife was evaluated (Table A-1). Results on protected taxa that occupy JBMDL indicate that insects appear to have the greatest percent (15.7%) impacted habitat, followed by herpetofauna (14%) (Fig. 1). These data suggest there are taxa specific constraints in movement that result in the likelihood of exposure. The only protected mammal on JBMDL is the bobcat (*Lynx rufus*) which can have a home range of 23 – 60 km² (Lovallo and Anderson 1996) compared to protected insects (Lepidoptera, approximately 0.75 km over lifetime, Schultz 1998). As such, taxa that have limited home ranges are likely going to have greater exposure potential if restricted to a location near AFFF release.

Version: DRAFT

June 2020

Figure 1. Percent of AFFF impacted habitat by protected taxa on JBMDL



McGuire has the greatest percent of estimated AFFF impacted habitat used by protected species that may occur on the installation, with exception to the state endangered bobcat (Figure 2, Appendix B). A dominant habitat (landcover layer) that the bobcat occupies is the coniferous forest (> 50% crown closure); importantly, it is not the case that this habitat is only available on Lakehurst but that the estimated AFFF impact to this habitat is greatest on Lakehurst (3.3%) compared to near zero percent impacted on McGuire and Dix. In total there are 10, 6,216 and 1,38 acres of coniferous forest (>50% crown closure) on McGuire, Dix and Lakehurst; respectively. Likewise, for the protected insects that have habitat preference for old fields and herbaceous wetlands, these habitat types are distributed across the entire installation; however, we estimate that these habitats are most impacted by historical AFFF activities on McGuire and therefore exposure potential is greatest for protected Lepidoptera on McGuire compared to Dix and Lakehurst bases. Given the comparatively small size of McGuire and yet similar number of known AFFF release locations on Dix and Lakehurst the PFAS exposure potential is disproportionately greater on this base (figure 2).




Figure 2. Percent of protected taxa-specific landcover estimated to be impacted by AFFF by base

Further investigation shows that there are differences in exposure potential to protected taxa by dietary requirements. For example, protected insectivorous avian taxa appear to have greater exposure potential based on preferred habitat type (Figure 3). Likewise, is true for protected reptilian taxa; where the estimated AFFF impacted habitat by dietary requirements are greatest for Omnivores (Squamates) > Carnivores (Serpentines) > Herbivores (Testudines). Again, the greatest estimated AFFF impacted habitat across reptile diet type is within McGuire compared to Dix and Lakehurst.




Figure 3. Percent of estimated AFFF impacted landscape of protected avian species by dietary requirements

#### Protected Species Ranking by Exposure Potential

Species exposure potential to PFAS is location specific, meaning species may be at greater risk of exposure on one site verses another. This is exactly the case at JBMDL where the species identified as having the greatest exposure potential to PFAS on McGuire are not the same as those identified on Dix nor Lakehurst (Table 3). This is due to a number of factors (1) habitat availability on the particular base (2) proximity of AFFF release to a given habitat and/or (3) best judgment in matching habitat preference with GIS landscape layer. For example, osprey (*Pandion haliaetus*) is identified as having greater exposure potential compared to the bald eagle (*Heliaeetus leucocephalus*) on McGuire base. Although there is habitat overlap among these species, osprey are almost exclusive piscivores and therefore habitats such as "wooded wetlands" which are assumed to have less fish than "lakes/ponds" were not designated as osprey habitat. Bald eagles may inhabit old growth forests and wooded wetlands and therefore the proportion of available habitat increases for Bald eagles but not for osprey (nearly obligate piscivores). Based on this type of spatial analyses it is not appropriate to draw commonalities in species ranking across the bases as habitat availability and AFFF release locations are variable. This approach is useful to identify species with greatest exposure potentials at a given location, the fine-scale GIS

EA Project No.: 6333101 SERDP Project No. ER.18-1626

Version: DRAFT

EA Engineering, Science, and Technology, Inc., PBC.

June 2020

landcover layer and ability to compartmentalize the large size of JBMDL allows for multiple lists of species ranking.

Version: DRAFT

June 2020

EA Engineering, Science, and Technology, Inc., PBC.

Table 3. Protected Species Exposure Potential to PFAS by base on JBMDL

Latin Name	Common Name	McGuire	Latin Name	Common Name	Dix	Latin Name	Common Name	Lakehurst
P. haliaetus	Osprey	19.47%	F. sparverius	American Kestrel	12.48%	A. flammeus	Short-eared Owl	3.65%
D. oryzivorus	Bobolink	18.72%	D. oryzivorus	Bobolink	9.97%	C. cyaneus	Northern Harrier	3.26%
P. gramineus	Vesper Sparrow	18.72%	P. sandwichensis	Savannah Sparrow	8.60%	P. rubriventris (Adult)	Red-bellied Turtle - Adult	2.82%
A. savannarum	Grasshopper Sparrow	17.00%	C. cyaneus	Northern Harrier	2.96%	S. varia	Barred Owl	2.17%
A. arogos arogos	Arogos Skipper	17.00%	P. haliaetus	Osprey	1.79%	L. rufus	Bobcat	2.16%
C. irus	Frosted Elfin	17.00%	P. rubriventris (Adult)	Red-bellied Turtle - Adult	1.63%	E. guttata	Corn Snake	2.16%
B. longicauda	Upland Sandpiper	16.99%	P. rubriventris (Juvenile)	Red-bellied Turtle-juvenile	1.50%	C. platensis	Sedge Wren	2.04%
A. henslowii	Henslow's sparrow	14.69%	H. chrysocelis	Southern Gray Tree Frog	1.47%	A. gentilis	Northern Goshawk	2.01%
P. montanus	Eastern Mud Salamander	14.37%	P. gramineus	Vesper Sparrow	1.02%	B. lineatus	Red-shouldered Hawk	2.01%
B. selene myrina	Silver-bodered Fritillary	14.35%	B. longicauda	Upland Sandpiper	0.95%	M. erythrocephalus	Red-headed woodpecker	2.01%
C.muhlenbergii	Bog Turtle	14.35%	S. varia	Barred Owl	0.89%	L. ludovicianus	Loggerhead Shrike	2.00%
C. platensis	Sedge Wren	14.32%	A. flammeus	Short-eared Owl	0.85%	A. otus	Long eared owl	1.97%
B. lentiginosos	American bittern	14.22%	A. gentilis	Norther Goshawk	0.72%	C. horridus	Timber Rattlesnake	1.92%
H. andersonii	Pine Barrens Tree Frog	14.15%	B. lineatus	Red-shouldered Hawk	0.72%	C. insculpta	Wood Turtle	1.79%
P. m melanoleucus	Northern Pine Snake	13.65%	M. erythrocephalus	Red-headed woodpecker	0.72%	N. violaceus	Yellow-crowned night heron	1.61%
P. podiceps	Pied-billed Grebe	12.52%	A. otus	Long eared owl	0.62%	C. muhlenbergii	Bog Turtle	1.52%
P. rubriventris (Juvenile	Red-bellied Turtle-juvenile	12.19%	P. montanus	Eastern Mud Salamander	0.50%	H. leucocephalus	Bald Eagle	1.45%
H. chrysocelis	Southern Gray Tree Frog	12.13%	N. violaceus	Yellow-crowned night heron	0.44%	N. nycticorax	Black-crowned night heron	1.45%
C. guttata	Spotted Turtle	11.95%	C. guttata	Spotted Turtle	0.43%	B. lentiginosos	American bittern	1.44%
C. insculpta	Wood Turtle	11.92%	H. leucocephalus	Bald Eagle	0.43%	B. selene myrina	Silver-bodered Fritillary	1.43%
H. leucocephalus	Bald Eagle	11.91%	N. nycticorax	Black-crowned night heron	0.43%	C. guttata	Spotted Turtle	1.42%
N. violaceus	Yellow-crowned night heron	11.91%	L. rufus	Bobcat	0.41%	P. podiceps	Pied-billed Grebe	1.15%
N. nycticorax	Black-crowned night heron	11.91%	E. guttata	Corn Snake	0.38%	H. chrysocelis	Southern Gray Tree Frog	1.14%
L. ludovicianus	Loggerhead Shrike	9.73%	C. insculpta	Wood Turtle	0.38%	P. rubriventris (Juvenile)	Red-bellied Turtle-juvenile	1.07%
C. cyaneus	Northern Harrier	9.08%	C. horridus	Timber Rattlesnake	0.34%	H. andersonii	Pine Barrens Tree Frog	0.93%
P. rubriventris (Adult)	Red-bellied Turtle - Adult	8.10%	P. podiceps	Pied-billed Grebe	0.27%	A. arogos arogos	Arogos Skipper	0.24%
A. flammeus	Short-eared Owl	7.53%	H. andersonii	Pine Barrens Tree Frog	0.24%	C. irus	Frosted Elfin	0.24%
A. otus	Long eared owl	4.33%	L. ludovicianus	Loggerhead Shrike	0.18%	P. montanus	Eastern Mud Salamander	0.23%
F. sparverius	American Kestrel	4.16%	A. arogos arogos	Arogos Skipper	0.13%	D. oryzivorus	Bobolink	0.19%
P. sandwichensis	Savannah Sparrow	4.15%	C. irus	Frosted Elfin	0.13%	A. savannarum	Grasshopper Sparrow	0.19%
E. guttata	Corn Snake	3.10%	A. savannarum	Grasshopper Sparrow	0.13%	A. henslowii	Henslow's sparrow	0.16%
C. horridus	Timber Rattlesnake	2.15%	A. henslowii	Henslow's sparrow	0.11%	P. sandwichensis	Savannah Sparrow	0.12%
A. gentilis	Norther Goshawk	1.31%	P. m melanoleucus	Northern Pine Snake	0.07%	P. gramineus	Vesper Sparrow	0.11%
B. lineatus	Red-shouldered Hawk	1.31%	C. platensis	Sedge Wren	0.05%	B. longicauda	Upland Sandpiper	0.11%
M. erythrocephalus	Red-headed woodpecker	1.31%	B. selene myrina	Silver-bodered Fritillary	0.03%	F. sparverius	American Kestrel	0.09%
L. rufus	Bobcat	0.16%	C. muhlenbergii	Bog Turtle	0.03%	P. m melanoleucus	Northern Pine Snake	0.08%
S. varia	Barred Owl	0.00%	B. lentiginosos	American bittern	0.03%	P. haliaetus	Osprey	0.07%

#### 4.1.2 Naval Air Station Patuxent River

Analysis on estimating impacts of AFFF to NAS Pax River habitat resulted in approximately 2% of the installation identified as PFAS impacted (Table A-2). Greatest estimated impacts (8%) were to evergreen forests, all other habitats had 4% or less of the available landscape impacted. As noted above in the "data reliability" section, the spatial scale of the available GIS layers were not well defined. Habitat (landscape layer) was updated following the in-field habitat assessment and GIS was updated with new landscapes and boundaries. Although, this update did not change the total estimated impacted landscape on NAS Patuxent River, it does influence the estimates of impacted habitat per species and per taxa. The exposure potential for most taxa using the original GIS was lower than the respective exposure potentials using the updated GIS landscapes (Figure 4).




Figure 4. Percent impacted habitat by taxa on NAS Patuxent River before and after GIS updates

The greatest difference in exposure potential can be seen in fish that jumps from <1% of the habitat exposed to 2.29%. Importantly, this estimate of impacted freshwater habitat is likely conservative as the migration of PFAS from the AFFF release site terminates at the surface water body and therefore the model does not include downstream portions of that landcover layer. There is a loss of habitat exposure potential in reptiles and marine mammals as the original GIS landscape has only one "water" landcover layer that is broken down into "tidal water", "non-tidal water", "emergent wetland" and "forested wetland" in the updated GIS. There were no AFFF

June 2020

EA Engineering, Science, and Technology, Inc., PBC

releases that resulted in PFAS impacts to the tidal water habitat in the updated GIS and therefore, no exposure potential to marine mammals and reptiles, which included only protected sea turtles. In comparison with the estimated impacted habitat of JBMDL, NAS Patuxent River has lower percentage of protected species exposure potential to PFAS. Interestingly, the insect taxa appear to have a high exposure potential at both installations indicating that further hazard/risk assessment may be warranted to those terrestrial receptors using non-traditional species (i.e. crickets). Furthermore, insectivorous species such as passerine birds may also experience greater exposure potential as resources may occupy impacted habitat. Further supporting the need to investigate, not only direct toxicity to terrestrial invertebrates but also PFAS accumulation potential. Non-traditional invertebrates are of interest as insectivores prey upon other invertebrate orders (i.e. Coleoptera, Diptera, etc., Nyffeler et al. 2018) more frequently than Lumbriculida (earthworms) a standard ecotoxicity test organism.

# Protected Species Ranking by Exposure Potential

Protected species on NAS Patuxent River were ranked by potential exposure to PFAS using the GIS spatial analysis. Ranking is provided as pre- and post-site visit where GIS landscape layers were updated to reflect the change in exposure potential by species (Table 4). The olive sided fly catcher was identified as having the greatest PFAS exposure potential following the site-visit; where pre-site visit it was the frosted elfin (non-migratory butterfly), which was still ranked comparatively high (2.1%) in the post-site visit. Freshwater fish, Ironcolor shiner and Flier were ranked as having greater exposure potential in the post-site visit compared to the pre-site visit where the "water" GIS landcover layer was redefined as "tidal" and "nontidal" water landcover. This update resulted in greater exposure potential to "nontidal" habitats and no exposure potential to "tidal" habitat thus, reduced exposure potentials to protected marine species (e.g. sea turtles and marine mammals). Furthermore, exposure potentials for the Common Gulled-billed Tern, Royal Tern and Black Skimmer were also reduced as these species occupy tidal coastal regions.

Importantly, there are a number of species (i.e. Henslow's Sparrow, Short-eared Owl) where exposure potential is relatively similar in the pre- and post-site visit GIS analysis. This is due to little or no change in those GIS landcover layers following updates. Although, updating the GIS landscape layers did not alter exposure potentials of some protected species, it would be imprudent to assume this is always the case. The spatial exposure potential analyses is very specific to geographic location and therefore any generic assumptions such as "wooded wetlands are always well defined" should not be made.

Table 4. Protected Species Exposure Potential to PFAS on NAS Patuxent River

Latin Name	Common Name	Patuxent River (pre-site visit)	Patuxent River (post-site visit)
C. cooperi	Olive sided Fly catcher	1.62%	2.44%
C. macropterus	Flier	0.99%	2.29%
N. chalybaeus	Ironcolor Shiner	0.99%	2.29%
C. platensis	Sedge Wren	1.34%	2.23%
C. d. dorsalis	tiger beetle	0.00%	2.19%
C. irus	Frosted Elfin	1.97%	2.12%
A. henslowii	Henslow's Sparrow	1.91%	2.02%
B. longicauda	Upland Sandpiper	1.97%	1.99%
L. ludovicianus	Loggerhead shrike	1.78%	1.90%
A. flammeus	short eared owl	1.89%	1.89%
G. philadelphia	Mourning warbler	1.44%	1.84%
S. fusca	Blackburnian warbler	1.45%	1.69%
H. leucocephalus	Bald Eagle	1.43%	1.61%
G. carolinensis	Narrow-mouthed toad	1.43%	1.59%
A. gentilis	Northern Goshawk	1.45%	1.58%
S. antillarum	Least Tern	0.83%	0.90%
C. melodus	Piping Plover	1.31%	0.45%
A. oxyrhynchus	Atlantic Sturgeon	0.99%	0.00%
E. glacialis	North Atlantic Right Whale	0.99%	0.00%
M. novaeangliae	Humpback Whale	0.99%	0.00%
G. nilotica	Common gulled-billed tern	0.83%	0.00%
R. niger	Black Skimmer	0.83%	0.00%
T. maximus	Royal Tern	0.83%	0.00%
L. kempii	Kemp's Ridley Sea Turtle	0.83%	0.00%
D. coriacea	Leatherback Sea Turtle	0.80%	0.00%
C. caretta	Loggerhead Sea Turtle	0.80%	0.00%

### 4.1.3 Barksdale Air Force Base

Spatial exposure estimates of AFFF on Barksdale AFB have resulted in approximately 110 of the 22,000 acres identified as impacted. All exposure scenarios are confined to the western portion of the base near the runway and other developed land (Figure C-8). The greatest exposure potential per landscape type was streams and canals at approximately 8% of those estimated to be impacted by PFAS (Table A-3). Currently, the only protected species found on this installation is the Bald Eagle. Estimated bald eagle impacted habitat on Barksdale AFB is less than 1% of total available. Given the large range of a bald eagle, nest locations relative to impacted sites is paramount as prey capture frequency from nearby water/field is likely greater than obtaining prey from locations further from the nest. Thus, PFAS exposure potential is expected to be correlated to nest locations near impacted sites, as seen in bald eagles near Lake Superior in Wisconsin where nestlings closer to the area of impact had higher contaminate loads compared to those at further distances (Kozie and Anderson 1991).

#### 4.2 SPECIES-SPECIFIC PROBABILISTIC EXPOSURE MODELS

The objective for this phase of the project was to develop probabilistic exposure models and preliminary risk estimates for two ecological receptors that are federally or state protected and whose habitat overlaps with PFAS contamination at our case study installations. The two species chosen were the Northern Pine Snake (*Pituophis melanoleucus*) and the Bald Eagle (*Heliaeetus leucocephalus*). These species represent two very different receptors and each present certain challenges with regard to estimating PFAS exposure and therefore illustrate important species-specific traits that might need to be considered when conducting PFAS exposure estimation.

The overlap of species habitat and estimated PFAS contamination suggests a potential for exposure. As a preliminary risk screen, those species whose habitat most overlaps with potential PFAS contamination are also most likely to be exposed to PFAS. At McGuire (of JBMDL) the Northern Pine Snake had a percent habitat impacted by PFAS of 13.6% which was among the highest for the reptilian taxa. Other herpetofauna had similarly high percentage of habitat overlap with PFAS contamination and importantly, herpetofauna occupy many DoD sites in general (Petersen et al. 2017). The pine snake was chosen as the receptor of interest because it represents a high-end PFAS exposure and, hence, if no exposure or risk was identified, it seems likely that risk of PFAS exposure to other terrestrial herpetofauna may be similarly low. The bald eagle was chosen as the other receptor of interest because it is federally protected and is widespread (common at the three installations studied here) and, hence, likely to be an important receptor at many DoD installations in North America. As well, percent impacted habitat was relatively high for avian raptors and these species have a relatively low TRV (Newsted et al., 2005) which is used to determine if there is concern of eliciting an affect at a particular exposure concentration. The sections below describe the methods and results for each of these focal receptors.

Arguably, the most commonly used approach for risk estimation is a deterministic approach in which a measure of effect is divided by a measure of exposure (or vice versa) and the resulting quotient compared to an established threshold (U.S EPA 1997). If the quotient exceeds the threshold, then the assessor concludes that there is a potential for adverse effects or risk (e.g., U.S. EPA 1997). While this type of deterministic approach is fairly easy to implement is it not an actual estimate of risk and suffers from some critical assumptions that impact both the result and inferences of the analysis. As an example, the measure of exposure in a quotient-based assessment is a point estimate which might be the mean or a higher confidence limit. Several significant research efforts have pointed to the problems and uncertainties associated with the deterministic approach and have suggested probabilistic methods that more explicitly incorporate spatial, temporal, and organismal variability which therefore generate actual risk estimates where risk is the probability of an adverse outcome.

We use a probabilistic exposure framework here along the lines of what has been conducted for song birds (Johnson et al. 2007) and small mammals (Williams et al. 2014) exposed to lead from small arms firing ranges. The premise of these and similar studies on probabilistic risk assessment is that the landscape is heterogenous with respect to contaminant distribution and in how organisms use the habitat. By more explicitly accounting for these heterogeneities, more accurate exposure estimates can be generated.

June 2020

EA Engineering, Science, and Technology, Inc., PBC

Implementation of probabilistic exposure estimation involves the use of Monte-Carlo simulations whereby receptor exposure is simulated through time and, in some cases, space (e.g., Hope 2000, Johnson et al. 2007, Williams et al. 2014). When many virtual receptors are simulated, the resulting distribution of exposure can be used to generate the probability that effects thresholds are exceeded (e.g., Hope 2000). This approach also lends itself well to explicit analysis of uncertainties whereby the effects of uncertain parameters can be explored via simulation. Probabilistic exposure estimation can therefore be helpful when exploring the potential risk of PFAS to threatened and endangered species.

Probabilistic exposure estimation approaches are most useful in cases where the ecological receptor of interest has a relatively large foraging range and/or where contaminant distribution is heterogenous with respect to the foraging range (Williams et al. 2014). The receptors chosen for this this case-study exploration of PFAS risk to threatened and endangered species, the bald eagle and northern pine snake, both can forage over a relatively large area and are not restricted to very specific habitat features at a small spatial scale. These species also present unique challenges with regard to exposure estimation and methods presented here may help for other, similar species whose critical habitat may overlap PFAS (or other)-contaminated areas.

# 4.2.1 Probabilistic exposure and risk framework for Pituophis melanoleucus

#### Species Background

The Northern pine snake, is a large-bodied colubrid. There are three subspecies of pine snakes with all three limited to the eastern United States and all considered rare. The Northern pine snake is the only pine snake found in New Jersey where JBMDL is also located. The range of this species in New Jersey is limited to a narrow fringe of pinelands. Hence, any reference to "pine snake" in the current document is in reference to the Northern pine snake. Pine snakes are large, nonvenomous snakes that are excellent at digging hibernacula. Adults can be as long as 2m and have a white-gray base color with brown or black blotching. In New Jersey, pine snakes are generally active mid-April to mid-October spending the remainder of the year in hibernacula. Females are oviparous and produce clutches that are typically 4-16 eggs that are laid in underground hibernaculum (Tennant and Barlett 2000). Pine snakes have been documented to live for up to 23 years in the wild (Golden et al. 2009). Nests are found almost exclusively in open areas with loose sandy soils which are easier to excavate compared to clay-soil compositions. The burrowing abilities of pine snakes clearly aids in nesting but also in the pursuit of subterranean prey such as moles, voles, and shrews. As a general rule, these snakes prefer well-drained, sandy, upland pine and pine-oak forests (Burger and Zappalorti 1986, Zappalorti et al. 2009, Golden et al. 2009). Mating occurs in mid-May and female snakes excavate nests and lay eggs from mid-June to early July. Pine snakes frequently reuse nesting sites which suggests that suitable nesting sites may be limited (Burger and Zappalorti 1992). Pine snakes overwinter in underground hibernacula they share with other pine snakes and sometimes other species of snakes. Pine snakes consume small

June 2020

EA Engineering, Science, and Technology, Inc., PBC

mammals, birds, and bird eggs. Juvenile or smaller pine snakes also consume invertebrates.

As mentioned, the pine snake preferred habitat characteristics include soil and forest types common only to the pine barren habitats in New Jersey. The population that occupies the pine barrens is totally isolated from other pine snake populations throughout the country (Golden et al. 2009). This isolation adds to the importance and vulnerability of this population as no natural recolonization or immigration is possible.

# PFAS Exposure Conceptual Framework for the Northern Pine Snake

There are no existing published studies that we are aware of in which any species of terrestrial snake has been collected and analyzed for PFAS. Similarly, there are limited data available for other reptiles. Several studies; however, have been published in which PFAS tissue concentrations were reported for several testudine species (turtles, e.g. Bangma et al. 2019, De Silva et al. 2011) but these animals have vastly different life histories, physiology and habitat requirements compared to the pine snake. Nevertheless, we can infer that the most likely PFAS exposure routes for snakes would be dietary, dermal, and possibly drinking water. Of these, dietary and dermal are likely to be most significant for pine snakes given the close contact with soils and PFAS contaminated food sources.

There is considerable uncertainty regarding PFAS uptake from direct dermal contact. At the time of this report, there are insufficient data to effectively estimate or model dermal exposure of PFAS to snakes. Given that pine snakes construct and use burrows dermal exposure likely warrants additional research attention.

For the purposes of this case-study risk assessment, dietary exposure is assumed to be the most likely and most important route of PFAS exposure. Pine snakes consume prey whole and are also, given their prey preference, likely to consume some soil (Beyer et al. 1994). Although there are no published estimates for incidental soil ingestion rates for snakes, estimates for other wildlife species range from <2% to approximately 10% (or more). Consumption of actual prey items (e.g., rodents) can be determined by allometric equations in which food ingestion rate or field metabolic rate are estimated from receptor body weight (US EPA 1993). Because snakes consume prey items whole, it is very possible that an individual snake could obtain several days worth of energetic content from a single prey item. Using the allometric equation for field metabolic rate (FMR) of snakes seems most appropriate and would allow modeling of intermittent feeding. Field metabolic rate estimates are obtained from organisms in the field and as such include energetic demands associated with activities such as foraging, predator avoidance, and mating in addition to basal and resting metabolic rates.

The equation for field metabolic rates in snakes was obtained from the Wildlife Exposure Factors Handbook (U.S. EPA 1993):

 $FMR (kcal/day) = 0.0535 * (body mass (g))^0.799$ 

Table 5. Exposure model parameters for *P. m. melanoleucus* 

Exposure model parameter	Value (variability), distribution	Reference		
Snake body mass	2000 g (sd = 500), normal	Golden et al. 2009		
Prey mass (rodent)	30  g (sd = 5), normal	Assumed (reasonable wild rodent)		
Gross energy prey	$1.7 \text{ kcal/g (sd} = 0.28), normal}$	US EPA 1993		
Assimilation efficiency	0.8	US EPA 1993		
Soil-to-Biota Accumulation	19.7	D'Hollander et al. 2014,		
Factor (SBAF)		Bogdanska et al. 2011, Brown et		
		al. 1997		
Exposure Duration	90 days	Period of activity/exposure		
		duration of PFOS studies		
Incidental Soil Ingestion	0.07  (min) - 0.13  (max),	Beyer et al. 1994, US EPA 1993		
	uniform			
PFOS soil concentration	16.86 (sd=75.69), lognormal	JBMDL Site Investigation Report		
(mean±sd surface soil at		(Aerostar SES LLC 2018)		
McGuire)				

Concentrations of PFAS in prey items can be based on actual measured concentrations if available for site-specific assessments or can be estimated from PFAS soil concentrations using biota-to-soil-accumulation factors (BSAF). There are actually few BSAF estimates for rodents and small birds for PFAS including PFOS (Conder et al. 2019). D'Hollander et al. (2014) conducted a field study on terrestrial receptors occupying habitats near a Fluorochemical plant in Belgium. They focused on PFOS and reported soil concentrations of 69±9 ng/g and liver concentrations in co-located rodents (wood mice) of 5,810±1,300 ng/g. Colubrids consume prev in their entirety (Figure 5), therefore, liver PFOS concentrations should be converted to wholebody PFOS. There are few data, however, where tissue and whole-body concentrations of PFAS have been reported in rodents. Bodanska et al. (2011) reported tissue distributions of PFOS in laboratory mice and showed that 45% of the PFOS was recovered in the liver followed by ~15% in bone, 10% in blood, 10% in skin and 10% in muscle. Brown et al. (1997) provided body weight and organ weights for rodents including mice. The liver, bone, blood, skin and muscle comprised 0.055, 0.11, 0.08, 0.16, and 0.38 of the total bodyweight of mice. The data from D'Hollander et al. (2014), Bodanska et al. (2011), and Brown et al. (1997) were combined to generate a Biota-to-Soil Accumulation Factor (BSAF) of 19.7.




Figure 5. Daily energy consumed for an individual snake. Consumption is not constant because snakes consume whole prey which can exceed their daily energy requirements.

Pine snake exposure on PFAS-contaminated sites is assumed to occur primarily via the diet since data on dermal exposure are limited although studies in the Salice lab are ongoing. Further, the prevailing hypothesis is that dermal exposure may be unlikely to contribute strongly to total PFAS exposure because the chemical structures of many PFAS are not conducive to significant uptake across lipid barriers (Conder et al. 2007). For diet items, the BSAF is used to estimate a whole body PFAS contaminant load in rodent prey items from soil PFAS data which are commonly available for sites in which PFAS monitoring has occurred. For most DoD installations where AFFF has been used, the dominant soil PFAS is PFOS (East et al., in preparation). Ongoing monitoring data continues to support this statement as do data from JBMDL. Also, because PFOS is among the most toxic PFAS studied thus far, this assessment focuses on estimating exposure and risk to PFOS only; this focus has been applied to recently published ecological risk assessments on DoD installations (e.g., Salice et al. 2018, Larson et al. 2018).

The pine snake occurs at JBMDL and has been observed occupying a range of habitats on the base (personal communication). Importantly, overlying PFAS contamination layers with pine snake habitat and range layers, shows a potential for overlap of 13.65% for McGuire, specifically (Figure C-9). Because pine snakes can move and forage over a relatively large area a significant challenge for exposure modeling lies in the fact that there will be considerable variability as to whether snakes consume PFOScontaminated prey. To account for this uncertainty, a random value from a binomial distribution was chosen to determine on each day that a snake would feed, whether that individual snake would consume a PFOScontaminated prey. In the absence of detailed, individual-based data we cannot know exactly whether and how often pine snakes consume PFOS-contaminated prev. The binomial distribution was populated with values ranging from 0.05 to 1.0 with 0.05 representing a very low likelihood of consuming PFOScontaminated prey and 1.0 representing 100% probability of consuming contaminated prey. Because the overlap of pine snake habitat and PFOS-contaminated soil was estimated at 13.65%, it seems reasonable to conclude that some snakes are likely to have relatively high exposure but not likely 100% and some are likely to have much




Figure 6. Estimated PFOS exposure for individual snakes with a probability of consuming PFOScontaminated prey of 0.5 (top) and 0.1365 (bottom).

lower exposures. Figure 6 provides output for the PFOS exposure model for simulated individual snakes with different probabilities of encountering and consuming PFOS-contaminated prey.

As expected, as the proportion of snake habitat that overlaps PFOS-contaminated areas on McGuire increases so too does PFOS exposure (Figure 7). Preliminary estimates of risks of PFOS to pine snakes can be obtained by comparing estimated PFOS exposure to available Toxicity Reference Values (TRVs). At the time of this report, there are no available TRVs for reptilian receptors. However, Salice et al. are currently in the process of submitting a manuscript for review on the toxicity of PFOS to juvenile and adult male brown anoles, *Anolis sagrei*. This represents the first available toxicity data for reptiles. The predominant effect of PFOS on lizards appears to be on growth or size. A PFOS exposure of 0.25 mg/kg/d is associated with a 10% decrease in growth of juvenile brown anoles. Although toxicity studies are ongoing and further consideration is needed to develop final reptilian TRVs, we used an exposure level of 0.25 mg/kg/d as an available threshold to obtain a sense of whether and under what circumstances pine snakes at McGuire might be exposed to PFOS at levels that would exceed current effect

concentrations. The red horizontal line in Figure 7 corresponds to the 0.25 mg/kg/d effects level. The Avian Level IV Predator TRV (Newsted et al. 2005) was also included on the figure as avian toxicity data can sometimes be used for reptilian risk assessments. Only for snakes in which a high proportion of their habitat was contaminated with PFOS did some exceedances of the avian TRV occur. There were no exceedances of the current toxicity threshold for reptilian species even with 100% overlap between habitat/forage area and PFOS contamination.




Figure 7. Log10 estimated PFOS exposure for 1000 snakes exposed for 90 days. Red horizontal line corresponds to 0.25 mg/kg/d PFOS, the existing toxicity threshold for reptiles; purple line corresponds to 0.021 mg/kg/d which is the avian predator TRV provided for comparison purposes

To our knowledge, this represents a first attempt to estimate risk to a reptilian receptor exposed to PFOS. Further, we did not locate any published studies in which risks of any chemical stressor to snakes had been assessed. Our results suggest that PFOS exposure to pine snakes is likely to be below current toxicity thresholds, even under circumstances when a high proportion of prey items are from PFOS-contaminated habitats. However, it is critical to consider that there are many uncertainties that can reduce confidence in the risk estimates.

A critical uncertainty lies in the lack of data that allows us to generate reliable estimates of PFOS concentrations in prey items from PFOS concentrations in soil. We located only a single study (D'Hollander et al. 2014) in which co-located soil and biota samples were analyzed for PFOS. The lack of this type of data has also been echoed by Conder et al. (2019). Further complicating

June 2020

EA Engineering, Science, and Technology, Inc., PBC

the analysis for pine snakes, is that these animals consume their prey whole and D'Hollander et al. (2014) reported only liver concentration in wood mice. A series of extrapolations and analysis using PFOS tissue distribution data (Bodanska et al. 2011) combined with relative organ weights in mice (Brown et al. 1997) were used to estimate a BSAF. Given the limited data and the need to use several sources, the current BSAF should be considered preliminary and highly uncertain. More data on PFAS concentrations in co-located soil and terrestrial prey species is needed to increase confidence in the current estimates.

In addition to the SBAF, snake movement patterns, foraging and burrowing are likely critical for accurate exposure estimation. Here, only foraging was explicitly modeled. The assumption was that as the overlap of pine snake habitat and PFOS-contamination increased, so too would exposure to PFOS-contaminated prey. This is often modeled as an area use factor (AUF) adjustment whereby an exposure estimate is simply multiplied by the proportion of the habitat that is contaminated. Instead, in the current effort, individual snakes were modeled for 90 days with the probability of encountering and consuming PFOS-contaminated prey dictated by the overlap of habitat. While not spatially explicit, per-se, this construct provides the framework for including greater spatial detail as well as providing a better representation of how exposure occurs. Moreover, the current analysis generates a probabilistic output that translates to an actual estimate of risk (probability of adverse outcome). This effort differed compared to Johnson et al. (2007) and efforts under ESTCP project ER-0197 (Williams et al. 2014) in that habitat suitability was not explicitly incorporated. In this case, available information and personal communications suggest that Northern pine snakes use and forage over a wide range of habitats and can be found on roads and in fields. Also, we do not have PFAS biota data matched to soil concentrations. The lack of strong habitat preferences and robust spatial PFAS data would obviate the utility of a true spatially-explicit modeling effort. Instead, we captured uncertainties that relate to habitat use and PFOS in prey by simulating exposure over a range of different probabilities of consuming PFOScontaminated prey. Because we generated only a few estimates where exposure exceeded the existing toxicity threshold, it seems unlikely that better spatial data would alter apparent risk conclusions.

As mentioned, pine snakes are excellent excavators and can consume subterranean prey (e.g., moles). If snakes happened to burrow in PFAS-contaminated soils, it seems this could result in a significant exposure. No data are available concerning dermal exposure in reptiles although the Salice Lab has conducted a preliminary study in which brown anoles were maintained on PFOS-amended sand. The resulting data showed an apparent uptake of PFOS from dermal exposure and/or incidental soil ingestion although the data were not robust enough to confirm that dermal exposure occurs and is an important exposure route. Nonetheless, the data do suggest that dermal exposure is possible and that additional studies may be warranted. Incidental soil ingestion may also be higher than was used here (~10%) given pine snake behaviors and preferred prey. Although increasing the incidental soil ingestion to 25% had little impact on risk estimates (data not shown).

Another important uncertainly is that there are few relevant toxicity data for reptiles exposed to PFOS or any other PFAS. As mentioned, the Salice lab has conducted a series of experiments in which brown anoles were exposed to PFOS. The resulting toxicity data suggests that adverse

June 2020

EA Engineering, Science, and Technology, Inc., PBC

effects on growth or size occur in the sub mg/kg/d range (~0.25 or perhaps lower). It is possible that the toxicity data being developed using a lizard model is not suitable for snakes or turtles although no robust comparisons for any chemical class had been conducted to determine how responses among reptiles differ. This remains an ongoing and important uncertainty.

This assessment focused on PFOS because the existing data from JBMDL has indicated that PFOS has been detected most frequently and at the highest concentrations. Similarly, an analysis of a large dataset of PFAS soil concentrations on US Air Force Bases shows that PFOS is the dominant PFAS. Nonetheless, it is important to consider and acknowledge that other PFAS may be present but not included as analytes and this could be an important uncertainty. Available toxicity data, however, also supports a focus on PFOS as it tends to be more toxic than other commonly studied and/or commonly occurring PFAS.

# 4.2.2 Probabilistic exposure and risk framework for Haliaeetus leucocephalus

# Species background

The bald eagle, is a large raptor with a range that extends over much of North America. Adult birds range from about 3000 to 6300g (Buehler 2000). Bald Eagles nest in forested areas adjacent to large bodies of water with nest usually placed in trees. The majority of bald eagle diet consists of fish that may be captured live, stolen from other animals, or as carrion. Bald eagles are opportunists, however, an also consume other animals and carrion. Although the species was once declining due to pesticides such as DDT and poaching, with protection the species has made an impressive comeback but is still protected in North America by the Bald Eagle Protection Act (USFWS), the Migratory Bird Species Act, and the Lacey Act.

Given the extensive range of Bald Eagles, it is likely that habitat layers that meet the requirements for Bald Eagles overlap with many DoD installations. For example, Bald Eagles appeared on the protected species lists for the three installations used for the case studies described above. It was, in fact, the only species of conservation interest for Barksdale Air Force Base (BAFB). Bald Eagles represent a potentially important and common predatory receptor for exposure to contaminants on DoD installations.

# <u>PFAS Exposure Conceptual Framework for the Bald Eagle</u>

The dietary exposure route for any contaminant, including PFAS, is generally considered the most relevant for bald eagles. Indeed, several published risk assessments for bald eagles have focused entirely on the dietary route (Rumbold 2005). Eagles spend a considerable portion of time in trees or airborne and, hence, dermal exposure is likely very low. A key driver of dietary exposure is the daily energetic requirement which determines the quantity of food needed and, hence, the delivery of contaminants. Specifically, field metabolic rate (FMR) is a reasonable determinant of energetic requirements of free-living animals compared to basal or resting metabolic rate. In an analysis of FMR for mammal and birds, Nagy (1987) showed that FMR was specific to body mass but generally similar among taxa although endothermic animals had considerably higher FMR than ectothermic animals, as expected. For avian species, developing

nestlings, in particular, have very high energetic demands because of the need to meet basal metabolic requirements as well as growth. For bald eagles, eaglets were selected as the most vulnerable life stage because of their high energetic needs and relatively long exposures. Eaglets take approximately 60 days to develop into a 4 kg subadult (Bortolotti 1984). Several studies on growth and metabolic demands of eaglets are available (e.g. Bortolotti 1984) and have been used to generate eaglet exposure models for ecological risk assessment (Rumbold 2005).

Unlike adult birds, growing eaglets show drastic changes in metabolic demands and mass through time. Rumbold (2005) provides a framework (used here) for estimating exposure to eaglets to methylmercury via consumption of contaminated fish from a constructed wetland in Southern Florida. The values and equations used to estimate exposure to eaglets and these were incorporated into the current case-study assessment (Table 2 *in* Rumbold 2005).

First, a generic growth model based on empirical data for eaglets is used to provide a daily mass estimate. The growth equation is:

$$dW/dt = -kaW (log_eW)$$

Where W is mass in g, k is a growth coefficient, and a is the asymptotic weight of an eaglet.

Figure 8 is a representation of eaglet growth from the equation described above. Body weight for any given day from this growth model eventually is used to calculate a daily intake of PFAS (mg PFAS/kg body mass/day).




Figure 8. Post-hatch growth of eaglets until fledging ~62-days.

As mentioned, FMR of eaglets is the driving element to determine food consumption and PFAS exposure. For eaglets, available studies on eaglet metabolism were focused either on earlier

development (days 1-38) or later development (days 39-60). Energetic needs, in kilojoules are then converted to ingestion of prey using energetic content of prey items combined with assimilation efficiency for avian species (US EPA, 1993). Figure 9 shows a realization of eaglet ingestion rate through the developmental period.




Figure 9. Realization of eaglet ingestion rate through the 60-day developmental period

Bald Eagles were the only species of conservation interest at BAFB for which there was potential overlap with PFAS-contaminated habitat. For the case-study assessment using eaglets as the receptor, we assumed that the diet was 80% fish and 20% mammals, although for any given day, the probability that a fish was consumed was determined from a binomial random number generator. The primary PFAS of interest was PFOS because previous field studies identified this PFAS as the most frequently detected and occurring at the highest magnitude in environmental media and in fish tissues (e.g., Lanza et al., 2017, Salice et al. 2018). Moreover, this is the only relevant PFAS for which an avian TRV is available (Newsted et al. 2005). In particular, Cooper Bayou was apparently contaminated via groundwater discharge from two former fire training areas (Salice et al. 2018). The former fire training areas have since been repurposed and concentrations of PFOS in surface soils are minimal thereby reducing likely exposures of Bald Eagles to PFOS-contaminated terrestrial prey items. Lanza et al. (2017) and Lanza (2015) reported fish tissue concentrations for fish collected over the course of a year from Cooper Bayou. These PFOS data in whole fish were used to estimate exposure; a reasonable mean PFOS concentration in whole fish was 2000 ng/g dry weight (sd = 1000). The potential distribution of whole body PFOS concentrations in fish was modeled as a lognormal distribution

and if a foraging eagle obtained a fish from Cooper Bayou, the PFOS concentration in that fish was assigned based on a random draw from the lognormal distribution. In essence, the model was construed such that eaglets consumed sufficient quantities of fish or mammals to meet their daily energy requirements. Eaglets were exposed to PFOS only by consuming PFOS-contaminated fish from Cooper Bayou.

A significant uncertainty in estimating PFOS exposure to Bald Eagles at BAFB lies whether any fish consumed were from PFOS-contaminated Cooper Bayou. A key determinant of where eagles may obtain prey is their home range size or foraging area. There are not data on foraging area for eagles at BAFB. Garrett et al. (1993) conducted a study of eagles on the Columbia River in Oregon and Washington. They fortunately provided estimates of use area for breeding eagles and determined that 25<sup>th</sup>, 50<sup>th</sup>, and 75<sup>th</sup> harmonic contours were 0.4, 1.3, and 4.5 km<sup>2</sup>, respectively, for breeding pairs of eagles. The 95th percentile harmonic contour was almost 22 km<sup>2</sup>. These data indicate that eagles can forage over a very large range but also that a significant proportion of foraging time is spent foraging over a relatively small area (0.4-1.3 km<sup>2</sup>). Moreover, eagle pairs tend to forage more closely to nesting sites. Figure 10 is a map of BAFB with contours placed at 0.5 km intervals. The area extending East of Cooper Bayou is more likely to contain Eagle nests given the lack of anthropogenic activity. Because we do not know the proportion of the Eagle diets that might come from Cooper Bayou, we varied the probability of obtaining fish from Cooper Bayou from 0.10 to 0.9 in increments of 0.1. This was modeled as a binomial such that for any given day of foraging, whether the eagle obtained prey from Cooper Bayou was determined by the outcome of a binomial random number generator. Importantly, for each day of foraging during the developmental period, there was an independent probability of obtaining prey from Cooper Bayou.



Figure 10. BAFB. Contours are 0.5km and extend East from Cooper Bayou. These are used to provide context for modeled probabilistic foraging from Cooper Bayou by Eagles nesting at different locations within BAFB.

Collectively, parents foraging in and around Cooper Bayou determined PFOS exposure to eaglets. For any given day, probabilistic algorithms were used to assign whether foraging eagles obtained fish or mammals, whether fish were from Cooper Bayou, and the PFOS concentration in fish if obtained from Cooper Bayou. An example PFOS exposure profile is provided in Figure 11 for a hypothetical nest that was assigned a probability of obtaining fish from Cooper Bayou of 0.5. In days where there was no PFOS exposure, foraging parents either provided mammalian food to eaglets or obtained a fish from somewhere other than Cooper Bayou (not specified). Exposures are higher early in development because of the relatively higher metabolic demands and low weight of young eaglets. PFOS exposure per individual eaglet was determined by taking the mean PFOS exposure over the 62-day development period. This estimate is akin to dosing or feed studies in which the average dose or approximate daily intake (ADI) is used as the exposure metric (Newsted et al. 2007).




Figure 11. Representative PFOS Dietary Exposure to Post-hatch Eaglet

A preliminary estimate of risk to eaglets at BAFB is summarized in figure 12. Again, the probability of obtaining fish prey from Cooper Bayou was explored as an uncertainty and characterization exercise. At the lowest probability of 0.10, for any given day, there was a 10% chance that prey would be obtained from Cooper Bayou. The 0.10 was included in a function to estimate a binomial (1 or 0) outcome; if a '1' was randomly drawn, the foraging eagle obtained a fish from Cooper Bayou. At the highest probability of 0.90, there was a 90% chance that every day foraging eagles would obtain prey from Cooper Bayou. If a fish was obtained from Cooper Bayou, the PFOS concentration was determined from a lognormal random number generation

the distribution of which was defined by whole body PFOS concentrations in fish from Lanza et al (2017). The ADI of PFOS for simulated eaglets at BAFB exceeded the existing Level IV predatory avian species TRV (Newsted et al. 2005); there were some exceedances even at the lowest probability of obtaining prey from Cooper Bayou (0.10). Above a probability of obtaining prey from Cooper Bayou of 0.20, the TRV was frequently exceeded. Note that the existing Level IV Avian Predator Lowest Observable Adverse Effect Level (LOAEL) was not exceeded even at the highest probability of foraging in Cooper Bayou (0.90).




Figure 12. Estimate of PFOS risk to eaglets at Barksdale Air Force Base

This analysis (Figure 12) suggests that there it is possible for eaglets at BAFB to be exposed to PFOS at levels that exceed existing Level IV predator avian TRV (Newsted et al. 2005). Importantly, the actual likelihood that eagles obtain prey from Cooper Bayou is unknown. Cooper Bayou is a relatively small bayou (3-5 m across) and therefore, by area, likely comprises a small proportion of eagle foraging range. Alternatively, in sampling Cooper Bayou for over a year for fish, large and varied fish species were frequently obtained (Lanza et al. 2017). In

June 2020

EA Engineering, Science, and Technology, Inc., PBC

comparison to nearby Flat River, which was larger, more and larger fish were routinely obtained from Cooper. Hence, while Cooper Bayou may be small, it is possible that piscivorous raptors may preferentially forage there because of the availability of prey. Very detailed data on eagle foraging behaviors are not available but given their wide ranging tendencies, it seems unlikely that there would be a probability higher than 0.20 that eagles would obtain prey from Cooper Bayou on a daily basis.

There are no laboratory toxicity data for the effects of PFOS on avian raptor species. The best available avian toxicity data for PFOS was obtained from studies on standard test species such as bobwhite quail and mallard ducks (Newsted et al. 2007, Newsted et al. 2005). Hence, the existing TRV for raptor species was obtained through the use of adjustment factors and represents, to some extent, an uncertain estimate. Existing field data for avian receptors exposed to PFOS and PFOS + other PFAS provides evidence that exposure and accumulation of PFAS occur in adults (e.g., Bustnes et al. 2013, Custer et al. 2014) and that PFAS are offloaded to eggs (e.g., Custer et al. 2014). One study in which plasma samples from bald eagles in the Midwest U.S. were analyzed for a variety of organic contaminants showed that PFOS had the highest mean and maximum measured concentrations compared to eighteen other organic contaminants (Elliott et al. 2019). Plasma concentrations of PFOS were several hundred times greater than most other measured organic contaminants and the maximum was about a thousand times greater than most other organic contaminants (Elliott et al. 2019). The PFOS plasma concentrations reported by Elliott et al. (2019) can be compared to the Level IV Avian Predator serum TRV that is 3.9 and 0.24 µg PFOS/ml for male and female avian predators, respectively (Newsted et al. 2005). The mean PFOS in plasma of Bald Eagles was 0.335 ug PFOS/ml which exceeds the avian TRV for females. The maximum plasma PFOS concentration was 4.2 µg PFOS/ml which exceeds both male and female avian predator TRVs. This study suggests strongly that bald eagles are exposed to organic contaminants like PFOS, potentially at concentrations high enough to exceed existing TRVs. Of note is that the plasma samples were obtained from eaglets from 159 eagle nests covering a wide area of the Midwest over the course of several years. No nests were located near known former (or current) fire training areas.

The current modeling construct is not spatially explicit in the sense that individual eagles are modeled as moving within a virtual landscape and then using habitat areas following specified habitat use rules. Johnson et al. (2007) implemented a spatially-explicit exposure and risk estimation study using the Spatially-Explicit Exposure Model (SEEM) for songbirds in which habitat suitability values were assigned to patches within the landscape and these were used to create occupancy rules for simulated birds. In this case, the occupancy rules that guided receptor habitat use had an impact on exposure to Pb at former small arms ranges (Johnson et al. 2007). In a follow-up study (Williams et al. 2014), Johnson and colleagues conducted a similar analysis using SEEM for small mammals. Interestingly, in that effort they concluded that the spatially-explicit approach did not yield estimates that were considered "better" than deterministic or point estimates because the forage area of small mammals was not large enough to encompass salient heterogeneities such as habitat or contaminant differences across the landscape. Hence, the net benefit of a full, spatially-explicit assessment depends on specifics of the receptor and contaminant distribution. Eagles forage over a wide area and are somewhat opportunistic often consuming carrion, when available. A detailed habitat suitability index that could be used to set

June 2020

EA Engineering, Science, and Technology, Inc., PBC

spatial use rules for eagles would be unlikely to improve exposure estimates. In this case, and for similar species, a broader assessment approach that includes the probability of encountering PFAS-contaminated prey may be more appropriate. Our analysis suggests that even a 0.1 probability of obtaining prey from PFAS-contaminated Cooper Bayou may result in exposures to eaglets that exceed the existing avian TRV (Newsted et al. 2005). For eaglets occupying nests in or around BAFB, some field studies may be warranted to establish whether eaglets are exposed. Comparisons between field data and model results is a powerful means of understanding how and when to best apply probabilistic and spatially-explicit exposure models (Johnson et al. 2007, Williams et al. 2014).

This current assessment focused on PFOS although we acknowledge that there are many PFAS that ecological receptors could be exposed to. The field data from BAFB and analyses of surface water samples from many US Air Force Base Installations (East et al., in review) suggest that PFOS is the dominant PFAS in surface waters followed by PFHxS. Any site-specific assessment for protected species must consider the site-specific data as much as possible to aid in identifying likely relevant PFAS. As it stands, an a-priori hypothesis is that most DoD installations that have had fire training or suppression activities are likely contaminated with PFOS. Moreover, thus far, PFOS appears to be the most toxic and among the most bioaccumulative. Future research and assessment efforts should more explicitly consider other PFAS, particularly, PFHxS.

The results of this assessment suggest that it is possible that eaglets may be exposed to PFOS at levels that exceed existing TRVs at BAFB. Our risk estimates would decrease if eagles did not obtain prey items from Cooper Bayou, which is possible. Additional, site- and species-specific studies are warranted to better understand potential risk of PFAS to protected raptor species. In particular, studies similar to Elliott et al. (2019) for raptor nests near DoD installations would prove extremely insightful.

### 5. CONCLUSIONS AND IMPLICATIONS FOR FUTURE RESEARCH

Development of the habitat exposure potential using GIS spatial analysis provides a ranking of installation and protected-species specific PFAS exposure potentials. The goal of this research was to provide a framework for determining risk to protected species, which is a function of hazard and exposure. Given that hazard data are limited for PFAS and especially with regard to protected species, our approach sought to focus on identifying protected species that may be exposed to PFAS and then estimate exposure as an initial step in determining if there is a potential risk. This approach is useful in prioritizing protected species (and wildlife taxa, in general) for further assessment should the potential for PFAS exposure be high. It should be noted that this analysis is suitable for evaluation of exposure potentials at specific sites and that broad categorizations should not be made. For example, species identified as having high exposure potentials were not the same across the three bases at JBMDL as habitat is variable on this large installation and thus, species occupancy is variable. Furthermore, AFFF release sites and estimated drainage areas are not correlated with any specific landscape.

June 2020

EA Engineering, Science, and Technology, Inc., PBC

Applying the spatial analysis to three case-study installations we found that the quality of GIS landcover layers contributes to the overall species-specific exposure potential. At JBMDL the high quality of the spatial data allowed for a comprehensive analysis and ranking of protected species based on estimated PFAS exposure potential. There were few updates made to the GIS landscape layer and therefore, little to no changes resulted in species exposure potentials. Although, exposure potential did not change drastically for grouped taxa from the updated GIS landscape layers at Naval Air Station Patuxent River, it did modify species-specific exposure potential (i.e. freshwater species). The GIS landcover layers at Barksdale Air Force Base (BAFB) also suffered from poor quality; however, this was a result of low resolution, not missing landscapes as seen for NAS Patuxent River. There was only one protected species (bald eagle) at BAFB so a comparison of exposure potentials in pre- and post-GIS updates could not be made.

Once a list of species with exposure potential was generated, two species were chosen from the installations to further develop and demonstrate exposure estimation methods that may be useful for future risk assessments. The northern pine snake (Pituophis m. melanoleucus) and the bald eagle (Haliaeetus leucocephalus) were identified as receptors of interest. The pine snake was identified as having moderate-high exposure potential at McGuire and reptilian taxa frequently occur on DoD sites. The bald eagle is a protected species that occupies all three installations and has moderate exposure potential at JBMDL and NAS Patuxent River. These species were chosen based on likelihood of exposure (habitat overlap with PFAS contamination), conservation interest, and because they presented several unique challenges with regard to exposure estimation. The results of the case-study exposure assessments indicated a very low likelihood that the pine snake would be exposed to PFOS at levels that would exceed existing toxicity thresholds but that eaglets may be exposed to PFOS at levels that exceed existing avian Level IV predator TRVs. The assessments for both species highlighted important uncertainties. For example, for pine snakes a significant uncertainty was the estimation of PFOS concentrations in whole rodents based on soil PFOS concentrations. For eagles, more detailed data on foraging behaviors within BAFB would help refine exposure estimates.

There are some excellent opportunities for future research that have become apparent during this effort. To address uncertainties in extrapolated PFAS migration within the landscape, environmental sampling along a gradient, or at the expected surface water termination would be beneficial for proof-of-concept. Protected species are ranked based on their exposure potential, which is driven by habitat overlap with PFAS contamination. This is a critical first step in identifying the species that warrant further consideration. We acknowledge there are limitations to this based on historical sampling of known AFFF release locations because sampling is often from specific, dispersed locations. More detailed field sampling over a wider range of space and habitats would be helpful as a proof-of-concept to inform how PFAS concentrations vary over a contaminated landscape. This could then inform approaches in GIS used to extrapolate across future site assessments.

In addition to environmental sampling, <u>a critical need</u> in understanding true ecological receptor PFAS exposure scenarios is sampling of biota in concert with environmental compartment sampling. This is especially true for the reptilian model as there are very few (if any) data

June 2020

EA Engineering, Science, and Technology, Inc., PBC

available for PFAS tissue concentrations in reptiles, or importantly, in their food items. For example, there are no available BSAF for terrestrial species. This includes terrestrial invertebrates that may be critical for exposures to reptilian and avian insectivores. Our model suggests the northern pine snake may occupy a PFOS impacted site and/or habitat but that exposure is not likely to elicit an adverse apical response. Sampling blood or tissue of a surrogate terrestrial reptilian species would be notably informative to this model. Importantly, these data should be collected in conjunction with animal tagging so robust PFAS exposure can be ascertained based on a temporal and spatial scale. These data would then be incorporated into the probabilistic snake model developed under this statement of need to refine expected reptile PFAS exposure scenarios. This combined modeling and field-based research can provide considerable insights into the actual and estimated PFAS exposure and risk (e.g., Johnson et al. 2017, Williams et al. 2014).

Finally, and perhaps most important would be developing a field component to determining PFOS exposures to bald eagle eaglets. As the current probabilistic model suggests and as some existing field data indicates, there may be risk to eaglets through exposures to dietary items with high PFAS concentrations. We assume proximity of nest location to PFAS impacted sites plays a role in the eaglet exposure profile and would suggest sampling nests along a spatial gradient from near to further distances from impacted sites. One non-lethal approach to determining PFOS exposure to eaglets would be to sample feathers that have been shed, this approach also results in minimum disturbance to the animals. In addition to feather sampling, droppings may be sampled for PFOS, the concern with this approach alone is that fecal material may be from parent or the eaglet whereas feathers can be more easily identified as adult or juvenile. Plasma samples of eaglets would be more informative and can be compared to existing plasma-based TRVs although obtaining plasma would be more labor intensive.

The abovementioned additional effort could serve as a 'ground truth' to the models developed under this statement of need. Moreover, the risk assessment community frequently requests more information on observed field effects to those modeled and/or inferred by environmental sampling alone.

### 6. LITERATURE CITED

Aerostar SES LLC, prepared for U.S. Army Corps of Engineers. April 2018. Site Investigation Report of Fire Fighting Foam Usage at Joint Base McGuire-Dix-Lakehurst Burlington and Ocean Counties, New Jersey. Final Report, Contract No. W912HN-15-C-0022

Apelberg, B. J., Witter, F. R., Herbstman, J. B., Calafat, A. M., Halden, R. U., Needham, L. L., Goldman, L. R. 2007. Cord serum concentrations of perfluorooctane sulfonate (PFOS) and perfluorooctanoate (PFOA) in relation to weight and size at birth. Environmental health perspectives, 115(11): 1670-1676.

Awad, E., Zhang, X., Bhavsar, S. P., Petro, S., Crozier, P. W., Reiner, E. J., Fletcher, R., Tittlemier, S.A., Braekevelt, E. (2011). Long-term environmental fate of perfluorinated compounds after accidental release at Toronto airport. Environmental science and technology, 45(19): 8081-8089.

Bangma, J.T, Ragland, J.M., Rainwater, T.R., Bowden, J.A., Gibbons, J.W., Reiner, J.L. 2019. Perfluoroalkyl substances in diamondback terrapins (*Malaclemys terrapin*) in coastal South Carolina. Chemosphere 215: 305 – 312.

Besser, J. M., Wang, N., Dwyer, F. J., Mayer, F. L., Ingersoll, C. G. 2005. Assessing contaminant sensitivity of endangered and threatened aquatic species: Part II. Chronic toxicity of copper and pentachlorophenol to two endangered species and two surrogate species. Archives of environmental contamination and toxicology, 48(2): 155-165.

Beyer, W. N., Connor, E. E., Gerould, S. 1994. Estimates of soil ingestion by wildlife. The journal of wildlife management, 375-382.

Bogdanska, J., Borg, D., Sundström, M., Bergström, U., Halldin, K., Abedi-Valugerdi, M., Bergman, A., Nelson, B., DePierre, J., Nobel, S. 2011. Tissue distribution of 35S-labelled perfluorooctane sulfonate in adult mice after oral exposure to a low environmentally relevant dose or a high experimental dose. Toxicology, 284(1-3): 54-62.

Bortolotti, G.R. 1984. Physical development of nestling bald eagles with emphasis on the timing of growth events. The Wilson Bulletin 96: 524 - 542.

Buck, R. C., Franklin, J., Berger, U., Conder, J. M., Cousins, I. T., De Voogt, P., Jensen, A.A., Kannan, K., Mabury, S.A., van Leeuwen, S. P. 2011. Perfluoroalkyl and polyfluoroalkyl substances in the environment: terminology, classification, and origins. Integrated environmental assessment and management, 7(4): 513-541.

Buehler, David A. (2000). Bald Eagle (*Haliaeetus leucocephalus*), version 2.0. In The Birds of North America (P. G. Rodewald, editor). Cornell Lab of Ornithology, Ithaca, New York, USA.

#### EA Engineering, Science, and Technology, Inc., PBC

Burger, J and R.T. Zappalorti. 1986. Nest Site Selection by Pine Snakes (*Pituophis melanoleucus*) in the New Jersey Pine Barrens. Copeia 1:116-121. Burger, J and R.T. Zappalorti. 1992. Philopatry and nesting phenology of pine snakes *Pituophis melanoleucus* in the New Jersey Pine Barrens. Behavioral Ecology and Sociobiology 30:331-336.

Chang, S.C., Noker, P.E., Gorman, G.S., Hart, Gibson. S.J., Ehresman, J.A., Butenhoff, D. J., 2012. Comparative pharmacokinetics of perfluorooctanesulfonate (PFOS) in rats, mice, and monkeys. Reproductive toxicology, 33: 428–440. doi:http://dx.doi.org/10.1016/j.reprotox.2011.07.002.

Conder, J.M., Hoke, R.A., Wolf, W.D., Russell, M.H., Buck, R.C. 2007. Are PFCAs Bioaccumulative? A critical review and comparison with regulatory criteria and persistent lipophilic compounds. Environmental Science and Technology 42: 995 – 1003.

Conder, J., Arblaster, J., Larson, E., Brown, J., Higgins, C. 2019. Guidance for assessing the ecological risks of PFAS to threatened and endangered species at aqueous film forming foam – impacted sites. SERDP Project ER18-1614.

Custer, C. M., Custer, T. W., Dummer, P. M., Etterson, M. A., Thogmartin, W. E., Wu, Q., Kannan, K., Trowbridge, A., McKann, P. C. 2014. Exposure and effects of perfluoroalkyl substances in tree swallows nesting in Minnesota and Wisconsin, USA. Archives of environmental contamination and toxicology, 66(1): 120-138.

D'Hollander, W., De Bruyn, L., Hagenaars, A., de Voogt, P., Bervoets, L. 2014. Characterisation of perfluorooctane sulfonate (PFOS) in a terrestrial ecosystem near a fluorochemical plant in Flanders, Belgium. Environmental science and pollution research, 21(20): 11856-11866.

Darwin, R.L. 2004. Estimated Quantities of Aqueous Film Forming Foam (AFFF) in the United States. Fire Fighting Foam Coalition, Arlington, VA.

De Silva, A. O., Spencer, C., Scott, B. F., Backus, S., Muir, D. C. G. 2011. Detection of a cyclic perfluorinated acid, perfluoroethylcyclohexane sulfonate, in the Great Lakes of North America. *Environ. Sci. Technol.* 19:8060-8066

Ding, G., Zhang, J., Chen, Y., Wang, L., Wang, M., Xiong, D., Sun, Y. (2013). Combined effects of PFOS and PFOA on zebrafish (*Danio rerio*) embryos. Archives of environmental contamination and toxicology, 64(4): 668-675.

Petersen, C.E., Lovich, R.E., Stallings, S.A. 2017. Amphibian and Reptile Biodiversity on United States Department of Defense Installations. Department of Defense, Legacy Resource Management Program, Final Report Aug 2017

Dwyer, F. J., Hardesty, D. K., Henke, C. E., Ingersoll, C. G., Whites, D. W., Augspurger, T.,

Canfield, T.J., Mount, D.R., Mayer, F. L. 2005. Assessing contaminant sensitivity of endangered and threatened aquatic species: Part III. Effluent toxicity tests. Archives of environmental contamination and toxicology, 48(2): 174-183.

CH2M. 2018. prepared for Department of the Navy. Preliminary Assessment Report for Potential Releases of Per- and Polyfluoroalkyl Substances (PFAS). Contract No. N62470-11-D-8012

Falandysz, J., Taniyasu, S., Yamashita, N., Rostkowski, P., Zalewski, K., Kannan, K. 2007. Perfluorinated compounds in some terrestrial and aquatic wildlife species from Poland. Journal of environmental science and health, Part A, 42(6): 715-719.

Geisy, J.P., Kannan, K. 2001. Global distribution of perfluorooctane sulfonate in wildlife. Environmental science and technology, 35: 1339-1342.

Giesy, J. P., Naile, J. E., Khim, J. S., Jones, P. D., Newsted, J. L. 2010. Aquatic toxicology of perfluorinated chemicals. In Reviews of environmental contamination and toxicology (pp. 1-52). Springer, New York, NY.

Gewurtz, S. B., Backus, S. M., De Silva, A. O., Ahrens, L., Armellin, A., Evans, M., Fraser, S.,

Golden, D.M., P. Winkler, P. Woerner, G. Fowles, W. Pitts, and D. Jenkins. 2009. Status Assessment of the Northern Pine Snake (*Pituophis m. melanoleucus*) in New Jersey: An Evaluation of Trends and Threats. New Jersey Department of Environmental Protection, Trenton, NJ. 53pp.

Hekster, F. M., Laane, R. W., de Voogt, P. 2003. Environmental and toxicity effects of perfluoroalkylated substances. In Reviews of environmental contamination and toxicology (pp. 99-121). Springer, New York, NY.

Higgins, C.P. Luthy, R.G. 2007. Modeling sorption of anionic surfactants onto sediment materials: An a priori approach for perfluoralkyl surfactants and linear alkylbenxene sulfonates. Environmental science and technology, 44: 3254-3261.

Hoff, P. T., Van Campenhout, K., Van de Vijver, K., Covaci, A., Bervoets, L., Moens, L., Huyskens, G., Goemans, G., Belpaire, C., Blust, R., De Coen, W. 2005. Perfluorooctane sulfonic acid and organohalogen pollutants in liver of three freshwater fish species in Flanders (Belgium): relationships with biochemical and organismal effects. Environmental pollution, 137(2): 324-333.

Hope, B.K. 2000. Generating probabilistic spatially-explicit individual and population exposure estimates for ecological risk assessment. Risk Analysis, 20(5):573-589.

Houtz, E.F., Higgins, C.P., Field, J.A., Sedlak, D.L. 2013. Persistence of Perfluoroalkyl acid

precursors in AFFF-impacted groundwater and soil. Environmental science and technology, 47: 8187-8195.

Hu, X.C., Andrews, D.Q., Lindstrom, A.B., Bruton, T.A., Schaider, L.A., Grandjean, P.,

HydroGeoLogic, Inc. 2015a prepared for Air Force Civil Engineer Center. August 2015. Final Preliminary Assessment Report for Perfluorinated Compounds at Joint Base McGuire-Dix-Lakehurst, New Jersey. Contract No. FA8903-08-D-8772 Task Order 0065

HydroGeoLogic, Inc. 2015b prepared for Air Force Civil Engineer Center. September 2015. Final Preliminary Assessment Report for Perfluorinated Compounds at Barksdale Air Force Base Bossier City, Louisiana. Contract No. FA8903-08-D-8772, Task Order 0065

Lohmann, R., Carignan, C.C., Blum, A., Balan, S.A., Higgins, C.P., 2016. Detection of poly-and perfluoroalkyl substances (PFASs) in US drinking water linked to industrial sites, military fire training areas, and wastewater treatment plants. Environmental science and technology letters, 3: 344 – 350.

Kannan, K., Franson, J.C., Boweman, W.W., Hansen, K.J., Jones, P.D., Geisy. J.P. 2001. Perfluorooctane sulfonate in fish eating water birds including bald eagles and albatrosses. Environmental science and technology, 35: 3065-3070.

Johnson, M.S., Wickwire, W.T., Quinn, M.J., Ziolkowski, D.J., Burmistrov, D.A., Menzie, C.A., Geraghty, C., Minnich, M., Parsons, P.J. 2007. Are songbirds at risk from lead at small arms ranges? An application of the spatially explicit exposure model. Environmental Toxicology and Chemistry 26(10):2215-2225.

Kannan, K., Corsolini, S., Falandysz, J., Fillmann, G., Kumar, K. S., Loganathan, B. G., Mohd, M.A., Olivero, J., Van Wouwe, N., Yang, J.H., Aldous, K. M. 2004. Perfluorooctanesulfonate and related fluorochemicals in human blood from several countries. Environmental science and technology, 38(17): 4489-4495.

Kannan, K., Perrotta, E., Thomas, N.J. 2006. Association between Perfluorinated compounds and pathological conditions in Southern Sea Otters. Environmental Science and Technology 40: 4943 – 4948

Kärrman, A., Elgh-Dalgren, K., Lafossas, C. Møskeland, T. 2011. Environmental levels and distribution of structural isomers of perfluoroalkyl acids after aqueous fire-fighting foam (AFFF) contamination. Environmental chemistry, 8: 372–380.

Kozie, K.D., Anderson, R.K. 1991. Productivity, diet, and environmental contaminants in bald eagles nesting near the Wisconsin shoreline of Lake Superior. Archives of environmental contamination and toxicology, 20: 41–48

Lanza, H.A. 2015. Perfluorinated Compound Residues in Biota Collected near Barksdale Air Force Base, Bossier City, Louisiana, USA. Master's Thesis, Texas Tech University.

Lanza, H. A., Cochran, R. S., Mudge, J. F., Olson, A. D., Blackwell, B. R., Maul, J. D., Salice, C.J. Anderson, T. A. 2017. Temporal monitoring of perfluorooctane sulfonate accumulation in aquatic biota downstream of historical aqueous film forming foam use areas. Environmental toxicology and chemistry, 36(8): 2022-2029.

Larson, E. S., Conder, J. M., Arblaster, J. A. 2018. Modeling avian exposures to perfluoroalkyl substances in aquatic habitats impacted by historical aqueous film forming foam releases. Chemosphere, 201, 335-341.

Long, G.C. Porter, R.C. 2015. USAF investigation of non-fire training area AFFF releases. Presentation at the DoD Environment, Safety and Occupational Health Network and Information Exchange (DENIX), Environmental Data Quality Workgroup Workshop. April 2015

Lovallo, M.J. Anderson, E.M. 1996. Bobcat (*lynx rufus*) Home range size and habitat use in Northwest Wisconsin. The American midland naturalist, 135: 241-252.

Moody, C. A., Martin, J. W., Kwan, W. C., Muir, D. C., Mabury, S. A. 2002. Monitoring perfluorinated surfactants in biota and surface water samples following an accidental release of fire-fighting foam into Etobicoke Creek. Environmental science and technology, 36(4): 545-551.

Newsted, J. L., Jones, P. D., Coady, K., Giesy, J. P. 2005. Avian toxicity reference values for perfluorooctane sulfonate. Environmental science and technology, 39(23): 9357-9362.

Nyffeler, M., Şekercioğlu, Ç. H., Whelan, C. J. 2018. Insectivorous birds consume an estimated 400-500 million tons of prey annually. Die Naturwissenschaften, 105(7-8): 47.

Oakes KD, Benskin JP, Martin JW, Ings JS, Heinrichs JY, Dixon DG, Servos MR. 2010. Biomonitoring of perfluorochemicals and toxicity to the downstream fish community of Etobicoke Creek following deployment of aqueous film-forming foam. Aquatic toxicology, 98: 120–129.

Petersen, C.E., Lovich, R.E, Stallings, S.A. 2017. Amphibian and reptile biodiversity on United States Department of Defense Installations. Department of Defense, Legacy Resource Management Program. Final Report, August 2017

Quesnelle, P.E., L. Fahrig and K.E. Lindsay. 2013. Effects of habitat loss, habitat configuration and matrix composition on declining wetland species. Biological Conservation. 160: 200 - 208.

Rumbold, D. G. 2005. A probabilistic risk assessment of the effects of methylmercury on great egrets and bald eagles foraging at a constructed wetland in South Florida relative to the Everglades. Human and ecological risk assessment, 11(2): 365-388.

Salice, C. J., Anderson, T. A., Anderson, R. H., Olson, A. D. 2018. Ecological risk assessment of perfluoroctane sulfonate to aquatic fauna from a bayou adjacent to former fire training areas at a US Air Force installation. Environmental toxicology and chemistry, 37(8): 2198-2209.

Schultz, C. B. 1998. Dispersal behavior and its implications for reserve design in a rare Oregon butterfly. Conservation biology, 12(2): 284-292.

Tennant, A and RD Barlett. 2000. Snakes of North America: Eastern and Central Regions. Gulf Publishing Company, Houston, Texas.

US EPA. 1993. WILDLIFE EXPOSURE FACTORS HANDBOOK. U.S. Environmental Protection Agency, Washington, D.C., EPA/600/R-93/187.

US EPA. 1997. Ecological Risk Assessment Guidance for Superfund: Process for Designing and Conducting Ecological Risk Assessments. Interim Final. EPA 540-R-97-006/OSWER 9285.7-25/PB97-963211. US Environmental Protection Agency, Solid Waste, and Emergency Response, Washington, DC.

US EPA. 2003. Assessment Factors, Science Policy Council, U.S. Environmental Protection Agency, Washington, D.C., EPA/100/B-03/001.

Weir, S.M., Suski, J.G., Salice, C.J. 2010. Ecological risk of anthropogenic pollutants to reptiles: Evaluating assumptions of sensitivity and exposure. Environmental pollution, 158: 3596 – 3606

Williams, M.A., Wickwire, T., Quinn, M.J., Johnson, M.S. 2014. Improvement, Verification, and Refinement of Spatially-explicit Exposure Models in Risk Assessment. Spatially-Explicit Exposure Model (SEEM) Demonstrations. Final Report. ESTCP PROJECT No. ER-0917.

Zappalorti, RT, MP McCort, DW Burkett, and DM Golden. 2009. 2008 Annual Report of Northern Pine Snake Management and Conservation at Stafford Business Park, Stafford Township, Ocean County, New Jersey. *Unpublished Report* 



Table A-1. Estimated exposure potential of landscapes at Joint Base McGuire Dix Lakehurst

		McGuire			Dix			Lakehur	st
							Habitat	Habitat	
	Habitat Impacted	Habitat Overall	Percent	Habitat	Habitat Overall	Percent	Impacted	Overall	Percent
Landscape Feature	(ac)	(ac)	Impacted	Impacted (ac)	(ac)	Impacted	(ac)	(ac)	Impacted
CEMETERY	0.00	0.01	0.00	0.00	3.47	0.00	0.00	0.00	0.00
RESIDENTIAL, HIGH DENSITY OR MULTIPLE DWELLING		0.00	0.00	0.00	0.00	0.00	0.00	20.73	0.00
RESIDENTIAL, RURAL, SINGLE UNIT	0.00	0.00	0.00	0.00	9.13	0.00	0.00	0.00	0.00
RESIDENTIAL, SINGLE UNIT, LOW DENSITY	0.00	0.00	0.00	0.00	0.20	0.00	0.00		0.00
RESIDENTIAL, SINGLE UNIT, MEDIUM DENSITY	0.00	0.00	0.00	0.00	1.05	0.00	0.00		0.00
COMMERCIAL/SERVICES	0.00	1.31	0.00	0.00	0.42	0.00	0.00	2.62	0.00
MILITARY INSTALLATIONS	178.12	1,317.90	13.52	31.35	3,132.47	1.00	94.59		10.47
TRANSPORTATION/COMMUNICATION/UTILITIES	0.05	5.11	0.89	1.36	220.36	0.62	0.00		0.00
UPLAND RIGHTS-OF-WAY UNDEVELOPED	0.00	3.10	0.00	0.00	81.09	0.00	0.00		0.00
STORMWATER BASIN	0.70	6.95	10.05	0.00	33.27	0.00	0.00		0.00
OTHER URBAN OR BUILT-UP LAND	86.57	1,194.66	7.25	16.61	1,250.32	1.33	75.42	1,729.81	4.36
MANAGED WETLAND IN MAINTAINED LAWN									
GREENSPACE	0.00	4.78	0.00	0.00	4.17	0.00	0.04		0.43
ATHLETIC FIELDS (SCHOOLS)	0.00	11.30	0.00	0.00	2.51	0.00	0.00		0.00
RECREATIONAL LAND	6.03	249.30	2.42	114.81	429.81	26.71	0.00		
AGRICULTURAL WETLANDS (MODIFIED)	0.00	0.00	0.00	92.84	93.97	98.80	0.00		0.00
CROPLAND AND PASTURELAND	0.00	0.01	0.00	0.00	14.17	0.00	0.00		0.00
DECIDUOUS FOREST (10-50% CROWN CLOSURE)	2.67	23.08	11.58	2.46	297.32	0.83	0.00		0.00
DECIDUOUS FOREST (>50% CROWN CLOSURE)	0.00	114.79	0.00	39.15	1,522.35	2.57	0.03		0.02
CONIFEROUS FOREST (10-50% CROWN CLOSURE)	0.04	2.72	1.52	0.00	3,476.78	0.00	0.00		0.00
CONIFEROUS FOREST (>50% CROWN CLOSURE)	0.00	10.32	0.00	7.15	6,216.02	0.12	45.68		3.31
PLANTATION	0.00	0.00	0.00	10.95	341.48	3.21	0.00	104.26	0.00
MIXED FOREST (>50% CONIFEROUS WITH 10-50%									
CROWN CLOSURE)	0.00	2.01	0.00	0.00	209.19	0.00	0.00	10.91	0.00
MIXED FOREST (>50% CONIFEROUS WITH >50%							45.40		
CROWN CLOSURE)	0.00	5.17	0.00	34.29	1,819.31	1.88	15.13	904.40	1.67
MIXED FOREST (>50% DECIDUOUS WITH 10-50%	0.00	2.45	0.00	0.00	100.22	0.00	0.00	14.63	0.00
CROWN CLOSURE)	0.00	3.45	0.00	0.00	100.33	0.00	0.00	14.62	0.00
MIXED FOREST (>50% DECIDUOUS WITH >50%	0.00	2.57	0.00	10.03	005.22	1.20	1 20	275 12	0.27
CROWN CLOSURE)	0.00 5.72	2.57 30.56	0.00	10.93	865.23	1.26			0.37
OLD FIELD (< 25% BRUSH COVERED) DECIDUOUS BRUSH/SHRUBLAND	1.41	13.30	18.73 10.59	1.19 0.00	830.21 78.60	0.14	0.36		
CONIFEROUS BRUSH/SHRUBLAND	0.00	0.00	0.00	0.00	679.92	0.00	5.44		2.71
MIXED DECIDUOUS/CONIFEROUS	0.00	0.00	0.00	0.00	073.32	0.00	3.44	200.54	2.71
BRUSH/SHRUBLAND	0.00	24.11	0.00	5.77	734.89	0.79	3.92	80.91	4.85
STREAMS AND CANALS	3.43	45.00	7.63	6.73	452.34	1.49	4.45	67.79	6.56
NATURAL LAKES	0.00	0.00	0.00	0.00	4.54	0.00	0.00		0.00
ARTIFICIAL LAKES	0.37	1.88	19.47	7.17	396.28	1.81	0.07	91.55	0.08
DECIDUOUS WOODED WETLANDS	2.26	75.92	2.97	9.80	2,334.37	0.42	2.58		4.05
CONIFEROUS WOODED WETLANDS	0.62	4.15	14.84	0.00	1,607.30	0.00	2.06		0.71
ATLANTIC WHITE CEDAR WETLANDS	0.00	0.00	0.00	0.00	298.97	0.00	0.00		0.00
DECIDUOUS SCRUB/SHRUB WETLANDS	0.00	0.64	0.00	0.00	371.69	0.00			4.02
CONIFEROUS SCRUB/SHRUB WETLANDS	0.00					0.00			
MIXED SCRUB/SHRUB WETLANDS (DECIDUOUS									
DOM.)	0.00	2.75	0.00	0.00	635.15	0.00	0.65	60.14	1.08
MIXED SCRUB/SHRUB WETLANDS (CONIFEROUS									
DOM.)	0.00	0.00	0.00	0.00	101.91	0.00	0.00	63.10	0.00
HERBACEOUS WETLANDS	54.96	379.58	14.48	0.74	843.63	0.09	0.00		
MIXED WOODED WETLANDS (DECIDUOUS DOM.)	0.00	5.56	0.00	12.13	523.83	2.32	1.44	47.03	
MIXED WOODED WETLANDS (CONIFEROUS DOM.)	0.00	0.23	0.00	0.00	635.50	0.00	0.00		0.00
DISTURBED WETLANDS (MODIFIED)	1.46	13.80	10.56	0.00	47.03	0.00	0.79		2.44
TRANSITIONAL AREAS	0.84	3.91	21.41	27.51	166.61	16.51	0.00		
UNDIFFERENTIATED BARREN LANDS	0.00	0.00	0.00	0.00		0.00			
PHRAGMITES DOMINANT INTERIOR WETLANDS	0.00			0.00		0.00			

Table A-2. Estimated exposure potential of landscapes at NAS Patuxent River

Landscape Feature	Habitat Impacted (ac)	Habitat Overall (ac)	Percent Impacted
Residential	0.00	19.80	0.00
Institutional	21.68	723.50	3.00
Transportation	10.13	503.97	2.01
Open Urban Land	37.57	1,917.60	1.96
Agriculture	0.00	149.09	0.00
Deciduous Forest	0.02	633.81	0.00
Evergreen Forest	8.90	99.75	8.92
Mixed Forest	24.78	1,323.13	1.87
Old Field	5.26	221.84	2.37
Brush	5.26	127.32	4.13
Nontidal Water	0.98	42.79	2.29
Tidal Water	0.00	47.41	0.00
Emergent Wetland	0.00	109.79	0.00
Forested Wetland	5.53	270.12	2.05
Sandy Shoreline	0.00	18.28	0.00

Table A-3. Estimated exposure potential of landscapes at Barksdale AFB

Landscape Feature	Habitat Impacted (ac)	Habitat Overall (ac)	Percent Impacted
Developed	26.32	3,160.15	0.83
Open Urban Land	4.39	1,750.40	0.25
Cultivated Crops	0.00	24.98	0.00
Pasture/Hay	0.00	27.70	0.00
Deciduous Forest	0.00	78.67	0.00
Evergreen Forest	0.00	6,572.84	0.00
Mixed Forest	0.00	606.28	0.00
Shrub/Scrub	0.00	461.84	0.00
Grassland/Herbaceous	0.00	436.50	0.00
Lakes and Ponds	1.25	737.58	0.17
Streams and Canals	56.56	722.51	7.83
Emergent Herbaceous Wetlands	0.00	25.73	0.00
Woody Wetlands	21.31	7,381.27	0.29




Table B-1. Spatial Exposure Potential of Protected Species at McGuire of Joint Base McGuire-Dix-Lakehurst

Feature Description	Habitat Status	Area (Ac)	Clemmy guttata	Pseudemys rubriventris (Juvenile)	Pseudemys rubriventris (Adult)	Hyla andersonii	Hyla chrysocelis
Cemetery	Impacted	0.00	NA	NA	NA	NA	NA
cemetery	Available	0.01	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Impacted	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Impacted	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA NA	NA 	NA	NA
Residential, Single Unit, Low Density	Impacted	0.00	NA	NA NA	NA	NA	NA
· · · · · · · · · · · · · · · · · · ·	Available	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Residential, Single Unit, Medium Density	Impacted	0.00	NA NA	NA NA	NA NA	NA NA	NA
	Available	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Commercial/Services	Impacted Available	1.31	NA NA	NA NA	NA NA	NA NA	NA NA
		178.12	NA NA	NA NA	NA NA	NA NA	NA NA
Military Installations	Impacted Available	1,317.90	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted	0.05	NA NA	NA NA	NA NA	NA NA	NA NA
Transportation/Communication/Utilities	Available	5.11	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Upland Right-of-Way Undeveloped	Available	3.10	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted	0.70	NA NA	NA NA	NA NA	NA NA	0.70
Stormwater Basin	Available	6.95	NA	NA NA	NA NA	NA	6.95
	Impacted	86.57	NA NA	NA NA	NA NA	NA NA	NA
Other Urban or Built-up Land	Available	1,194.66	NA	NA NA	NA NA	NA	NA
	Impacted	0.00	NA	0.00	NA NA	NA	0.00
Managed Wetland in Maintained Lawn Greenspace	Available	4.78	NA	4.78	NA	NA	4.78
	Impacted	0.00	NA	NA	NA	NA	NA
Athletic Fields (Schools)	Available	11.30	NA	NA	NA	NA	NA
	Impacted	6.03	NA	NA	NA	NA	NA
Recreational Land	Available	249.30	NA	NA	NA	NA	NA
A	Impacted	0.00	NA	0.00	NA	NA	0.00
Agricultural Wetlands (Modified)	Available	0.00	NA	0.00	NA	NA	0.00
Cropland and Pastureland	Impacted	0.00	NA	0.00	NA	NA	0.00
Cropiana ana Pastureiana	Available	0.01	NA	0.01	NA	NA	0.01
Deciduous Forest (10-50% Crown Closure)	Impacted	2.67	NA	NA	NA	NA	NA
Deciduous Forest (10-50% Crown Closure)	Available	23.08	NA	NA	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
Deciduous Forest (>30% Crown Closure)	Available	114.79	NA	NA	NA	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted	0.04	NA	NA	NA	NA	NA
connectous rolest (10 30% crown closure)	Available	2.72	NA	NA	NA	NA	NA
Coniferous Forest (>50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
Connections ( Connection Closure)	Available	10.32	NA	NA	NA	NA	NA
Plantation	Impacted	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
,	Available	2.01	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
· ',	Available	5.17	NA NA	NA NA	NA	NA NA	NA NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted	0.00	NA	NA NA	NA	NA	NA
, , , , , , , , , , , , , , , , , , ,	Available	3.45	NA NA	NA NA	NA NA	NA NA	NA NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	2.57	NA NA	NA NA	NA NA	NA NA	NA NA
Old Field (< 25% Brush Covered)	Impacted	5.72	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	30.56	NA NA	NA NA	NA NA	NA NA	NA NA
Deciduous Brush/Shrubland	Impacted	1.41	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	13.30	NA NA	NA NA	NA NA	NA NA	NA NA
Coniferous Brush/Shrubland	Impacted	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	0.00	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Clemmy guttata	Pseudemys rubriventris (Juvenile)	Pseudemys rubriventris (Adult)	Hyla andersonii	Hyla chrysocelis
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
Iwikea Deciduous/Connerous Brush/Shrubhana	Available	24.11	NA	NA	NA	NA	NA
Streams and Canala	Impacted	3.43	3.43	NA	3.43	NA	NA
Streams and Canals	Available	45.00	45.00	NA	45.00	NA	NA
Nietuwel Lekse	Impacted	0.00	0.00	0.00	0.00	NA	NA
Natural Lakes	Available	0.00	0.00	0.00	0.00	NA	NA
Artificial Lakes	Impacted	0.37	0.37	0.37	0.37	NA	NA
Artificial takes	Available	1.88	1.88	1.88	1.88	NA	NA
Deciduous Wooded Wetlands	Impacted	2.26	2.26	2.26	NA	NA	2.26
Deciduous wooded wetlands	Available	75.92	75.92	75.92	NA	NA	75.92
Coniferous Wooded Wetlands	Impacted	0.62	0.62	0.62	NA	0.62	0.62
Connerous wooded wetlands	Available	4.15	4.15	4.15	NA	4.15	4.15
Atlantic White Coder Wetlands	Impacted	0.00	0.00	0.00	NA	0.00	0.00
Atlantic White Cedar Wetlands	Available	0.00	0.00	0.00	NA	0.00	0.00
Dociduous Conub (Chaub Matlanda	Impacted	0.00	0.00	0.00	NA	0.00	0.00
Deciduous Scrub/Shrub Wetlands	Available	0.64	0.64	0.64	NA	0.64	0.64
Coniferous Scrub/Shrub Wetlands	Impacted	0.00	0.00	0.00	NA	0.00	0.00
Connerous Scrub/Sirrub Wetlands	Available	0.00	0.00	0.00	NA	0.00	0.00
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	0.00	0.00	NA	0.00	0.00
ivilxed Scrub/Siliub Wetlands (Deciduous Dolli.)	Available	2.75	2.75	2.75	NA	2.75	2.75
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	0.00	0.00	NA	0.00	0.00
wettands (Connerous Doni.)	Available	0.00	0.00	0.00	NA	0.00	0.00
Herbaceous Wetlands	Impacted	54.96	54.96	54.96	NA	54.96	54.96
Tierbaceous wectarius	Available	379.58	379.58	379.58	NA	379.58	379.58
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted	0.00	0.00	0.00	NA	0.00	0.00
TVINCE TVOCACE TVCItalias (Decladous Dollis)	Available	5.56	5.56	5.56	NA	5.56	5.56
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted	0.00	0.00	0.00	NA	0.00	0.00
Timed trooded treatailes (connerous boilin)	Available	0.23	0.23	0.23	NA	0.23	0.23
Disturbed Wetlands (Modified)	Impacted	1.46	NA	1.46	NA	NA	1.46
- Islandea recitantes (meanica)	Available	13.80	NA	13.80	NA	NA	13.80
Transitional Areas	Impacted	0.84	NA	NA	NA	NA	NA
	Available	3.91	NA	NA	NA	NA	NA
Undifferentiated Barren Lands	Impacted	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
Phragmites Dominant Interior Wetlands	Impacted	0.00	NA	NA	NA	NA	NA
•	Available	0.00	NA	NA	NA	NA	NA
Species Habitat Impacted (ac)			61.63	59.66	3.80	55.58	59.99
Species Habitat Available Overall (ac)			515.70	489.30	46.88	392.91	494.37
Percent Impacted Over Available			11.9512%	12.1925%	8.1015%	14.1452%	12.1348%
	Percent Available,	Not Impacted	88.0488%	87.8075%	91.8985%	85.8548%	87.8652%

Notes:

ac = acres

NA = not applicable

Feature Description	Habitat Status	Area (Ac)	Pseudotriton montanus	Accipiter gentilis	Ammodramus henslowii	Ammodramus savannarum	Asio flammeus
Comotory	Impacted	0.00	NA	NA	NA	NA	0.00
Cemetery	Available	0.01	NA	NA	NA	NA	0.01
Residential, High Density or Multiple Dwelling	Impacted	0.00	NA	NA	NA	NA	NA
Residential, high Density of Multiple Dweiling	Available	0.00	NA	NA	NA	NA	NA
Desidential Donal Circle Heit	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Commercial/Services	Available	1.31	NA	NA	NA	NA	NA
	Impacted	178.12	NA NA	NA	NA NA	NA NA	NA NA
Military Installations	Available	1,317.90	NA	NA NA	NA NA	NA NA	NA
		0.05	NA NA	NA NA	NA NA	NA NA	NA NA
Transportation/Communication/Utilities	Impacted						
	Available	5.11	NA NA	NA NA	NA 0.00	NA 0.00	NA NA
Upland Right-of-Way Undeveloped	Impacted	0.00	NA NA	NA NA	0.00	0.00	NA NA
	Available	3.10	NA NA	NA NA	3.10	3.10	NA NA
Stormwater Basin	Impacted	0.70	NA	NA	NA	NA	NA
	Available	6.95	NA	NA	NA	NA	NA
Other Urban or Built-up Land	Impacted	86.57	NA	NA	NA	NA	86.57
<u> </u>	Available	1,194.66	NA	NA	NA	NA	1,194.66
Managed Wetland in Maintained Lawn Greenspace	Impacted	0.00	NA	NA	NA	NA	NA
	Available	4.78	NA	NA	NA	NA	NA
Athletic Fields (Schools)	Impacted	0.00	NA	NA	NA	NA	NA
Attrietic Fields (Schools)	Available	11.30	NA	NA	NA	NA	NA
Recreational Land	Impacted	6.03	NA	NA	NA	NA	NA
Recreational Land	Available	249.30	NA	NA	NA	NA	NA
A:   \Al-al   \Al-al   \Al-al   \Al-al	Impacted	0.00	NA	NA	NA	NA	NA
Agricultural Wetlands (Modified)	Available	0.00	NA	NA	NA	NA	NA
Construct Book of Book	Impacted	0.00	NA	NA	NA	NA	0.00
Cropland and Pastureland	Available	0.01	NA	NA	NA	NA	0.01
	Impacted	2.67	NA	NA	NA	NA	NA
Deciduous Forest (10-50% Crown Closure)	Available	23.08	NA	NA	NA	NA	NA
	Impacted	0.00	NA	0.00	NA	NA NA	NA
Deciduous Forest (>50% Crown Closure)	Available	114.79	NA	114.79	NA	NA	NA
	Impacted	0.04	NA NA	NA	NA NA	NA NA	NA NA
Coniferous Forest (10-50% Crown Closure)	Available	2.72	NA	NA	NA NA	NA NA	NA
	Impacted	0.00	NA NA	0.00	NA NA	NA NA	NA NA
Coniferous Forest (>50% Crown Closure)	Available	10.32	NA NA	10.32	NA NA	NA NA	NA NA
				NA	NA NA	NA NA	NA NA
Plantation	Impacted	0.00	NA NA				
	Available	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
<u> </u>	Available	2.01	NA NA	NA 2.22	NA NA	NA NA	NA NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted	0.00	NA	0.00	NA	NA	NA
7	Available	5.17	NA	5.17	NA	NA	NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
2. 2.2	Available	3.45	NA	NA	NA	NA	NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted	0.00	NA	0.00	NA	NA	NA
	Available	2.57	NA	2.57	NA	NA	NA
Old Field (< 25% Brush Covered)	Impacted	5.72	NA	NA	5.72	5.72	5.72
Tota Field (> 25% Brusii Coverea)	Available	30.56	NA	NA	30.56	30.56	30.56
Desiduence Brush /Shamble d	Impacted	1.41	NA	NA	NA	NA	NA
Deciduous Brush/Shrubland	Available	13.30	NA	NA	NA	NA	NA
n if n 1 (a) 11 1	Impacted	0.00	NA	NA	NA	NA	NA
Coniferous Brush/Shrubland	Available	0.00	NA	NA	NA	NA NA	NA
		5.50		1 197.	1 '''''	1	

Feature Description	Habitat Status	Area (Ac)	Pseudotriton montanus	Accipiter gentilis	Ammodramus henslowii	Ammodramus savannarum	Asio flammeus
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
lviixed Deciduous/Connerous Brush/Shrubland	Available	24.11	NA	NA	NA	NA	NA
Streams and Canals	Impacted	3.43	NA	NA	NA	NA	NA
Streams and Canais	Available	45.00	NA	NA	NA	NA	NA
Natural Lakes	Impacted	0.00	0.00	NA	NA	NA	NA
ivaturar Lakes	Available	0.00	0.00	NA	NA	NA	NA
Artificial Lakes	Impacted	0.37	0.37	NA	NA	NA	NA
Artificial Lakes	Available	1.88	1.88	NA	NA	NA	NA
Deciduous Wooded Wetlands	Impacted	2.26	NA	2.26	NA	NA	NA
Deciduous wooded wettands	Available	75.92	NA	75.92	NA	NA	NA
Coniferous Wooded Wetlands	Impacted	0.62	NA	0.62	NA	NA	NA
Connerous wooded wedands	Available	4.15	NA	4.15	NA	NA	NA
Atlantic White Cedar Wetlands	Impacted	0.00	0.00	0.00	NA	NA	NA
Atlantic white Cedar Wetlands	Available	0.00	0.00	0.00	NA	NA	NA
Deciduous Scrub/Shrub Wetlands	Impacted	0.00	NA	NA	NA	NA	NA
Deciduous Scrub/Siliub Wetlands	Available	0.64	NA	NA	NA	NA	NA
Coniferous Scrub/Shrub Wetlands	Impacted	0.00	NA	NA	NA	NA	NA
connerous scrub/smub wetlands	Available	0.00	NA	NA	NA	NA	NA
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA	NA	NA	NA
Wince Scrub/Siliub Wetlands (Decidadus Dolli.)	Available	2.75	NA	NA	NA	NA	NA
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	NA	NA	NA
Trinca serab/sirab tretianas (connercus bonn)	Available	0.00	NA	NA	NA	NA	NA
Herbaceous Wetlands	Impacted	54.96	54.96	NA	54.96	NA	NA
	Available	379.58	379.58	NA	379.58	NA	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted	0.00	NA	0.00	NA	NA	NA
,	Available	5.56	NA	5.56	NA	NA	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted	0.00	NA	0.00	NA	NA	NA
,	Available	0.23	NA	0.23	NA	NA	NA
Disturbed Wetlands (Modified)	Impacted	1.46	1.46	NA	NA	NA	NA
	Available	13.80	13.80	NA 	NA	NA 	NA NA
Transitional Areas	Impacted	0.84	NA	NA NA	NA	NA NA	NA NA
	Available	3.91	NA	NA NA	NA 0.00	NA 0.00	NA NA
Undifferentiated Barren Lands	Impacted	0.00	NA	NA NA	0.00	0.00	NA NA
	Available	0.00	NA NA	NA NA	0.00	0.00	NA 0.00
Phragmites Dominant Interior Wetlands	Impacted	0.00	NA NA	NA NA	NA NA	NA NA	0.00
	Available 0.00  Species Habitat Impacted (ac)		NA 56.79	NA 2.87	NA 60.69	NA 5.72	0.00 92.29
	Species Habitat Availab		56.78 395.26	2.87	60.68 413.24	5.72 33.66	1,225.24
	Percent Impacted (		14.3665%	1.3138%	14.6853%	17.0029%	7.5325%
	Percent Available,						
	reiteiit Available,	ivot iiiipacted	85.6335%	98.6862%	85.3147%	82.9971%	92.4675%

ac = acres

Feature Description	Habitat Status	Area (Ac)	Asio otus	Batramia longicauda	Botaurus lentiginosos	Buteo lineatus	Circus cyaneus
Comotory	Impacted	0.00	0.00	0.00	NA	NA	0.00
Cemetery	Available	0.01	0.01	0.01	NA	NA	0.01
Residential, High Density or Multiple Dwelling	Impacted	0.00	NA	NA	NA	NA	NA
Residential, High Density of Multiple Dwelling	Available	0.00	NA	NA	NA	NA	NA
Basidantial Bural Cinala Hait	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Available	0.00	NA	NA	NA	NA	NA
Basidantial Circle Halt Law Basita	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Available	0.00	NA	NA	NA	NA	NA
Build and Contain the State of	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Available	0.00	NA	NA	NA	NA	NA
0	Impacted	0.00	NA	NA	NA	NA	NA
Commercial/Services	Available	1.31	NA	NA	NA	NA	NA
	Impacted	178.12	NA	NA	NA	NA	NA
Military Installations	Available	1,317.90	NA	NA	NA	NA	NA
	Impacted	0.05	NA	NA	NA	NA	NA
Transportation/Communication/Utilities	Available	5.11	NA	NA	NA	NA	NA
	Impacted	0.00	NA	0.00	NA NA	NA NA	0.00
Upland Right-of-Way Undeveloped	Available	3.10	NA	3.10	NA	NA NA	3.10
	Impacted	0.70	NA NA	NA	NA NA	NA NA	NA
Stormwater Basin	Available	6.95	NA	NA NA	NA NA	NA NA	NA NA
	Impacted	86.57	NA NA	NA NA	NA NA	NA NA	86.57
Other Urban or Built-up Land	Available	1,194.66	NA	NA NA	NA NA	NA NA	1,194.66
	Impacted	0.00	NA NA	NA NA	NA NA	NA NA	0.00
Managed Wetland in Maintained Lawn Greenspace	Available	4.78	NA NA	NA NA	NA NA	NA NA	4.78
		0.00	NA NA	NA NA	NA NA	NA NA	0.00
Athletic Fields (Schools)	Impacted						
	Available	11.30	NA NA	NA NA	NA NA	NA NA	11.30
Recreational Land	Impacted	6.03	NA	NA NA	NA NA	NA NA	NA NA
	Available	249.30	NA NA	NA NA	NA NA	NA NA	NA 0.00
Agricultural Wetlands (Modified)	Impacted	0.00	NA	NA NA	NA NA	NA NA	0.00
	Available	0.00	NA 2.22	NA 2.22	NA NA	NA NA	0.00
Cropland and Pastureland	Impacted	0.00	0.00	0.00	NA	NA NA	0.00
	Available	0.01	0.01	0.01	NA NA	NA NA	0.01
Deciduous Forest (10-50% Crown Closure)	Impacted	2.67	2.67	NA 	NA	NA NA	NA
	Available	23.08	23.08	NA	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted	0.00	0.00	NA	NA	0.00	NA
	Available	114.79	114.79	NA	NA	114.79	NA
Coniferous Forest (10-50% Crown Closure)	Impacted	0.04	0.04	NA	NA	NA	NA
	Available	2.72	2.72	NA 	NA	NA	NA
Coniferous Forest (>50% Crown Closure)	Impacted	0.00	0.00	NA	NA	0.00	NA
· ·	Available	10.32	10.32	NA	NA	10.32	NA
Plantation	Impacted	0.00	NA	0.00	NA	NA	NA
	Available	0.00	NA	0.00	NA	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted	0.00	0.00	NA	NA	NA	NA
2 200 200 200 200 200 200 200 200 200 2	Available	2.01	2.01	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted	0.00	0.00	NA	NA	0.00	NA
	Available	5.17	5.17	NA	NA	5.17	NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted	0.00	0.00	NA	NA	NA	NA
200000000000000000000000000000000000000	Available	3.45	3.45	NA	NA	NA	NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted	0.00	0.00	NA	NA	0.00	NA
	Available	2.57	2.57	NA	NA	2.57	NA
Old Field (< 25% Brush Covered)	Impacted	5.72	5.72	5.72	NA	NA	5.72
old Field (> 23/0 brush covered)	Available	30.56	30.56	30.56	NA	NA	30.56
Deciduous Brush/Shrubland	Impacted	1.41	NA	NA	NA	NA	NA
Deciduous bi usii/ siii ubialiu	Available	13.30	NA	NA	NA	NA	NA
Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
Connectors brush/shirubidhu	Available	0.00	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Asio otus	Batramia longicauda	Botaurus lentiginosos	Buteo lineatus	Circus cyaneus
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
lwiixed Deciduous/Confierous Brusii/Sifrubiand	Available	24.11	NA	NA	NA	NA	NA
Streams and Canals	Impacted	3.43	NA	NA	NA	NA	NA
Streams and Canais	Available	45.00	NA	NA	NA	NA	NA
Natural Lakes	Impacted	0.00	NA	NA	NA	NA	NA
inaturar Lakes	Available	0.00	NA	NA	NA	NA	NA
Artificial Lakes	Impacted	0.37	NA	NA	NA	NA	NA
Artificial Lakes	Available	1.88	NA	NA	NA	NA	NA
Deciduous Wooded Wetlands	Impacted	2.26	NA	NA	NA	2.26	NA
Deciduous wooded wettands	Available	75.92	NA	NA	NA	75.92	NA
Coniferous Wooded Wetlands	Impacted	0.62	NA	NA	NA	0.62	NA
Connerous wooded wedands	Available	4.15	NA	NA	NA	4.15	NA
Atlantic White Cedar Wetlands	Impacted	0.00	NA	NA	NA	0.00	NA
Atlantic white Cedar Wetlands	Available	0.00	NA	NA	NA	0.00	NA
Deciduous Scrub/Shrub Wetlands	Impacted	0.00	NA	NA	0.00	NA	0.00
Deciduous Scrub/Siliub Wetlands	Available	0.64	NA	NA	0.64	NA	0.64
Coniferous Scrub/Shrub Wetlands	Impacted	0.00	NA	NA	0.00	NA	0.00
connerous scrub/smub wetlands	Available	0.00	NA	NA	0.00	NA	0.00
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA	0.00	NA	0.00
wincu scrub/siirub wetianus (bediaudus boiii.)	Available	2.75	NA	NA	2.75	NA	2.75
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	0.00	NA	0.00
Trinca salabysinab tretianas (connelous bonn)	Available	0.00	NA	NA	0.00	NA	0.00
Herbaceous Wetlands	Impacted	54.96	NA	NA	54.96	NA	54.96
	Available	379.58	NA	NA	379.58	NA	379.58
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA	NA	0.00	NA
,	Available	5.56	NA	NA	NA	5.56	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	NA	0.00	NA
	Available	0.23	NA	NA	NA	0.23	NA
Disturbed Wetlands (Modified)	Impacted	1.46	NA	NA	1.46	NA	NA
	Available	13.80	NA	NA NA	13.80	NA NA	NA 0.04
Transitional Areas	Impacted	0.84	NA	NA NA	NA NA	NA	0.84
	Available	3.91	NA 0.00	NA 0.00	NA NA	NA NA	3.91
Undifferentiated Barren Lands	Impacted	0.00	0.00	0.00	NA NA	NA NA	0.00
	Available	0.00	0.00 NA	0.00 NA	NA NA	NA NA	0.00
Phragmites Dominant Interior Wetlands	Impacted		NA NA	NA NA	NA NA	NA NA	0.00
	Available 0.00  Species Habitat Impacted (ac)		8.44	5.72	56.42	2.87	148.09
	Species Habitat Availab		8.44 194.69	33.68	396.77	2.87	1,631.30
	Percent Impacted (		4.3339%	16.9901%	14.2197%	1.3138%	9.0781%
	Percent Available,		4.5559% 95.6661%	83.0099%	85.7803%	98.6862%	90.9219%
	r er cent Avanable,	ot impacted	33.0001/0	03.003370	03./003/0	JO.UOUZ /0	JU.JL1J/0

ac = acres

Feature Description	Habitat Status	Area (Ac)	Cistothorus platensis	Dolichonyx oryzivorus	Falco sparverius	Haliaeetus leucocephalus	Lanius ludovicianus
Comptony	Impacted	0.00	NA	NA	0.00	NA	0.00
Cemetery	Available	0.01	NA	NA	0.01	NA	0.01
Residential, High Density or Multiple Dwelling	Impacted	0.00	NA	NA	NA	NA	NA
Residential, high Density of Multiple Dweiling	Available	0.00	NA	NA	NA	NA	NA
Desidential Donal Circle Heit	Impacted	0.00	NA	NA	0.00	NA	NA
Residential, Rural, Single Unit	Available	0.00	NA	NA	0.00	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Commercial/Services	Available	1.31	NA	NA	NA	NA	NA
	Impacted	178.12	NA	NA	NA	NA NA	NA
Military Installations	Available	1,317.90	NA	NA	NA NA	NA	NA
	Impacted	0.05	NA NA	NA NA	NA NA	NA NA	NA NA
Transportation/Communication/Utilities	Available	5.11	NA	NA NA	NA NA	NA NA	NA
	Impacted	0.00	NA NA	NA NA	0.00	NA NA	0.00
Upland Right-of-Way Undeveloped	Available	3.10		NA NA	3.10	NA NA	3.10
	†		NA NA	NA NA	3.10 NA	NA NA	3.10 NA
Stormwater Basin	Impacted	0.70	NA NA				
	Available	6.95	NA NA	NA NA	NA NA	NA NA	NA NA
Other Urban or Built-up Land	Impacted	86.57	NA	NA	NA	NA NA	NA
	Available	1,194.66	NA	NA	NA	NA	NA
Managed Wetland in Maintained Lawn Greenspace	Impacted	0.00	NA	NA	0.00	NA	NA
	Available	4.78	NA	NA	4.78	NA	NA
Athletic Fields (Schools)	Impacted	0.00	NA	NA	0.00	NA	NA
	Available	11.30	NA	NA	11.30	NA	NA
Recreational Land	Impacted	6.03	NA	NA	6.03	NA	NA
	Available	249.30	NA	NA	249.30	NA	NA
Agricultural Wetlands (Modified)	Impacted	0.00	NA	0.00	0.00	NA	NA
- S. Idantara: Tronamas (Indamosa)	Available	0.00	NA	0.00	0.00	NA	NA
Cropland and Pastureland	Impacted	0.00	NA	0.00	0.00	NA	0.00
eropiana ana rastarelana	Available	0.01	NA	0.01	0.01	NA	0.01
Deciduous Forest (10-50% Crown Closure)	Impacted	2.67	NA	NA	NA	NA	NA
Deciduous Forest (10-30% Crown Closure)	Available	23.08	NA	NA	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
Deciduous Polest (>50% Clown Closule)	Available	114.79	NA	NA	NA	NA	NA
Coniferent Forest (10 F0% Crown Closure)	Impacted	0.04	NA	NA	NA	NA	NA
Coniferous Forest (10-50% Crown Closure)	Available	2.72	NA	NA	NA	NA	NA
Constanting Founds (> FOO( Constanting Classics)	Impacted	0.00	NA	NA	NA	NA	NA
Coniferous Forest (>50% Crown Closure)	Available	10.32	NA	NA	NA	NA	NA
Diameteria	Impacted	0.00	NA	NA	0.00	NA	NA
Plantation	Available	0.00	NA	NA	0.00	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Available	2.01	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA NA	NA	NA NA	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Available	5.17	NA	NA NA	NA NA	NA	NA
	Impacted	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Available	3.45	NA	NA NA	NA NA	NA NA	NA
	Impacted	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Available	2.57	NA NA	NA NA	NA NA	NA NA	NA NA
		+					
Old Field (< 25% Brush Covered)	Impacted	5.72	NA NA	5.72	5.72	NA NA	5.72
	Available	30.56	NA NA	30.56	30.56	NA NA	30.56
Deciduous Brush/Shrubland	Impacted	1.41	NA	NA NA	NA NA	NA NA	1.41
	Available	13.30	NA NA	NA NA	NA NA	NA NA	13.30
Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA 	0.00
	Available	0.00	NA	NA	NA	NA	0.00

Feature Description	Habitat Status	Area (Ac)	Cistothorus platensis	Dolichonyx oryzivorus	Falco sparverius	Haliaeetus leucocephalus	Lanius ludovicianus
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	0.00
Ivilked Deciduous/Connerous Brushi/Sin abiana	Available	24.11	NA	NA	NA	NA	24.11
Streams and Canals	Impacted	3.43	NA	NA	NA	3.43	NA
Streams and Canais	Available	45.00	NA	NA	NA	45.00	NA
Natural Lakes	Impacted	0.00	NA	NA	NA	0.00	NA
ivaturai Lakes	Available	0.00	NA	NA	NA	0.00	NA
Artificial Lakes	Impacted	0.37	NA	NA	NA	0.37	NA
Artificial Lakes	Available	1.88	NA	NA	NA	1.88	NA
Deciduous Wooded Wetlands	Impacted	2.26	NA	NA	NA	2.26	NA
Deciduous wooded wettands	Available	75.92	NA	NA	NA	75.92	NA
Coniferous Wooded Wetlands	Impacted	0.62	NA	NA	NA	0.62	NA
Connerous wooded wedands	Available	4.15	NA	NA	NA	4.15	NA
Atlantic White Codar Wetlands	Impacted	0.00	NA	NA	NA	0.00	NA
Atlantic White Cedar Wetlands	Available	0.00	NA	NA	NA	0.00	NA
Deciduous Scrub/Shrub Wetlands	Impacted	0.00	0.00	NA	NA	0.00	0.00
Deciduous Scrub/Siliub Wetlands	Available	0.64	0.64	NA	NA	0.64	0.64
Coniferous Scrub/Shrub Wetlands	Impacted	0.00	0.00	NA	NA	0.00	0.00
connerous scrub/smub wetianus	Available	0.00	0.00	NA	NA	0.00	0.00
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA	NA	0.00	0.00
ivincu serub/siriub vvetianus (beetuuous boin.)	Available	2.75	NA	NA	NA	2.75	2.75
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	NA	0.00	0.00
Trinca seras/siras tretianas (connersas serin)	Available	0.00	NA	NA	NA	0.00	0.00
Herbaceous Wetlands	Impacted	54.96	54.96	NA	NA	54.96	NA
	Available	379.58	379.58	NA	NA	379.58	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA	NA	0.00	NA
,	Available	5.56	NA	NA	NA	5.56	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	NA	0.00	NA
,	Available	0.23	NA	NA	NA	0.23	NA
Disturbed Wetlands (Modified)	Impacted	1.46	1.46	NA	NA	1.46	1.46
	Available	13.80	13.80	NA	NA	13.80	13.80
Transitional Areas	Impacted	0.84	NA	NA	0.84	NA NA	NA
	Available	3.91	NA NA	NA 0.00	3.91	NA NA	NA NA
Undifferentiated Barren Lands	Impacted	0.00	NA	0.00	0.00	NA NA	NA NA
	Available	0.00	NA NA	0.00 NA	0.00	NA 0.00	NA NA
Phragmites Dominant Interior Wetlands	Impacted	0.00 0.00	NA NA		NA NA	0.00	NA NA
	Available Species Habitat		NA	NA 5.72	NA 12.50	0.00	NA 8 FO
	Species Habitat Availab		56.42	5.72	12.59	63.09	8.59 88.29
	Percent Impacted C	, ,	394.02	30.57	302.97	529.51	9.7288%
	Percent Available,		14.3190%	18.7205%	4.1555%	11.9149%	
	Percent Available,	not impacted	85.6810%	81.2795%	95.8445%	88.0851%	90.2712%

ac = acres

Feature Description	Habitat Status	Area (Ac)	Melanerpes erythrocephalus	Nyctanassa violaceus	Nycticorax nycticorax	Pandion haliaetus	Passerculus sandwichensis
Cometery	Impacted	0.00	NA	NA	NA	NA	0.00
Cemetery	Available	0.01	NA	NA	NA	NA	0.01
Residential, High Density or Multiple Dwelling	Impacted	0.00	NA	NA	NA	NA	NA
Residential, high Density of Multiple Dwelling	Available	0.00	NA	NA	NA	NA	NA
Desidential Donal Circle Heit	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Commercial/Services	Available	1.31	NA	NA	NA	NA	NA
	Impacted	178.12	NA NA	NA NA	NA NA	NA NA	NA NA
Military Installations	Available	1,317.90	NA NA	NA NA	NA NA	NA NA	NA NA
Transportation/Communication/Utilities	Impacted	0.05	NA	NA NA	NA	NA NA	NA
	Available	5.11	NA NA	NA NA	NA NA	NA NA	NA 0.00
Upland Right-of-Way Undeveloped	Impacted	0.00	NA	NA 	NA	NA 	0.00
	Available	3.10	NA	NA NA	NA	NA	3.10
Stormwater Basin	Impacted	0.70	NA	NA	NA	NA	NA
	Available	6.95	NA	NA	NA	NA	NA
Other Urban or Built-up Land	Impacted	86.57	NA	NA	NA	NA	NA
Other Orban or Bunt-up Land	Available	1,194.66	NA	NA	NA	NA	NA
Managed Watland in Maintained Louis Consumer	Impacted	0.00	NA	NA	NA	NA	NA
Managed Wetland in Maintained Lawn Greenspace	Available	4.78	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Athletic Fields (Schools)	Available	11.30	NA	NA	NA	NA	NA
	Impacted	6.03	NA	NA NA	NA	NA	6.03
Recreational Land	Available	249.30	NA	NA NA	NA NA	NA NA	249.30
		0.00	NA NA	NA NA	NA NA	NA NA	NA
Agricultural Wetlands (Modified)	Impacted Available			NA NA			
		0.00	NA NA		NA NA	NA NA	NA NA
Cropland and Pastureland	Impacted	0.00	NA	NA NA	NA	NA	NA
	Available	0.01	NA	NA	NA	NA	NA
Deciduous Forest (10-50% Crown Closure)	Impacted	2.67	NA	NA	NA	NA	NA
	Available	23.08	NA	NA	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted	0.00	0.00	NA	NA	NA	NA
Decidadas Forest (* 50% Grown Glosare)	Available	114.79	114.79	NA	NA	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted	0.04	NA	NA	NA	NA	NA
Connerous Forest (10-30% Crown Closure)	Available	2.72	NA	NA	NA	NA	NA
Consideration Female (5 500/ Consum Classical)	Impacted	0.00	0.00	NA	NA	NA	NA
Coniferous Forest (>50% Crown Closure)	Available	10.32	10.32	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Plantation	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA NA	NA NA	NA NA	NA	NA NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Available	2.01	NA NA	NA NA	NA NA	NA NA	NA NA
		0.00	0.00	NA NA	NA NA	NA NA	NA NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted						
	Available	5.17	5.17	NA NA	NA NA	NA NA	NA NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted	0.00	NA	NA 	NA	NA NA	NA
,	Available	3.45	NA	NA NA	NA	NA	NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted	0.00	0.00	NA	NA	NA	NA
	Available	2.57	2.57	NA	NA	NA	NA
Old Field (< 25% Brush Covered)	Impacted	5.72	NA	NA	NA	NA	5.72
olu Fielu (> 25% bi usii Coveleu)	Available	30.56	NA	NA	NA	NA	30.56
D. 11 D 1 (c) 1	Impacted	1.41	NA	NA	NA	NA	NA
Deciduous Brush/Shrubland	Available	13.30	NA	NA	NA	NA	NA
	Impacted	0.00	NA NA	NA NA	NA NA	NA	NA NA
Coniferous Brush/Shrubland	Available	0.00	NA	NA NA	NA NA	NA NA	NA NA
	Available	0.00	IVA	IVA	IVA	INA	INA

Feature Description	Habitat Status	Area (Ac)	Melanerpes erythrocephalus	Nyctanassa violaceus	Nycticorax nycticorax	Pandion haliaetus	Passerculus sandwichensis
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
Iwixed Deciduous/Conferous Brush/Shrubhand	Available	24.11	NA	NA	NA	NA	NA
Streams and Canals	Impacted	3.43	NA	3.43	3.43	NA	NA
Streams and Canais	Available	45.00	NA	45.00	45.00	NA	NA
Natural Lakes	Impacted	0.00	NA	0.00	0.00	0.00	NA
ivaturar Lakes	Available	0.00	NA	0.00	0.00	0.00	NA
Artificial Lakes	Impacted	0.37	NA	0.37	0.37	0.37	NA
Artificial Lakes	Available	1.88	NA	1.88	1.88	1.88	NA
Deciduous Wooded Wetlands	Impacted	2.26	2.26	2.26	2.26	NA	NA
Deciduous wooded wettands	Available	75.92	75.92	75.92	75.92	NA	NA
Coniferous Wooded Wetlands	Impacted	0.62	0.62	0.62	0.62	NA	NA
Connerous wooded wedands	Available	4.15	4.15	4.15	4.15	NA	NA
Atlantic White Codar Wetlands	Impacted	0.00	0.00	NA	0.00	NA	NA
Atlantic White Cedar Wetlands	Available	0.00	0.00	NA	0.00	NA	NA
Deciduous Scrub/Shrub Wetlands	Impacted	0.00	NA	0.00	0.00	NA	NA
Deciduous Scrub/Siliub Wetlands	Available	0.64	NA	0.64	0.64	NA	NA
Coniferous Scrub/Shrub Wetlands	Impacted	0.00	NA	0.00	0.00	NA	NA
Connerous Scrub/Sinub Wetianus	Available	0.00	NA	0.00	0.00	NA	NA
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	NA	0.00	0.00	NA	NA
wince serus/sirius wettands (beetadous boin.)	Available	2.75	NA	2.75	2.75	NA	NA
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	NA	0.00	0.00	NA	NA
Trinca serab/sirab tretianas (connersus bonn)	Available	0.00	NA	0.00	0.00	NA	NA
Herbaceous Wetlands	Impacted	54.96	NA	54.96	54.96	NA	NA
	Available	379.58	NA	379.58	379.58	NA	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted	0.00	0.00	0.00	0.00	NA	NA
,	Available	5.56	5.56	5.56	5.56	NA	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted	0.00	0.00	0.00	0.00	NA	NA
,	Available	0.23	0.23	0.23	0.23	NA	NA
Disturbed Wetlands (Modified)	Impacted	1.46	NA	1.46	1.46	NA	NA
	Available	13.80	NA	13.80	13.80	NA	NA
Transitional Areas	Impacted	0.84	NA	NA NA	NA	NA	NA
	Available	3.91	NA NA	NA NA	NA NA	NA NA	NA 0.00
Undifferentiated Barren Lands	Impacted	0.00	NA	NA NA	NA NA	NA NA	0.00
	Available	0.00	NA NA	NA 0.00	NA 0.00	NA NA	0.00
Phragmites Dominant Interior Wetlands	Impacted	0.00 0.00	NA NA	0.00		NA NA	NA NA
	Available Species Habitat		NA 2.87	0.00	0.00	NA 0.27	NA 11.75
	Species Habitat Availab		2.87 218.70	63.09 529.51	63.09 529.51	0.37 1.88	11.75 282.97
	Percent Impacted (	, -	1.3138%	11.9149%	11.9149%	1.88 19.4651%	4.1533%
	Percent Available,						
	reiteiit Available,	Mor impacted	98.6862%	88.0851%	88.0851%	80.5349%	95.8467%

ac = acres

Feature Description	Habitat Status	Area (Ac)	Podilymbus podiceps	Pooecetes gramineus	Strix varia	Atrytone arogos arogos	Bolaria selene myrina
Comotory	Impacted	0.00	NA	NA	NA	NA	NA
Cemetery	Available	0.01	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Impacted	0.00	NA	NA	NA	NA	NA
Residential, high bensity of Multiple Dwelling	Available	0.00	NA	NA	NA	NA	NA
Desidential Dural Cinale Unit	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Available	0.00	NA	NA	NA	NA	NA
Built-off Code to the Built	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Commercial/Services	Available	1.31	NA	NA	NA	NA	NA
	Impacted	178.12	NA	NA	NA	NA	NA
Military Installations	Available	1,317.90	NA	NA	NA	NA	NA
	Impacted	0.05	NA	NA	NA	NA	NA
Transportation/Communication/Utilities	Available	5.11	NA	NA	NA	NA	NA NA
	Impacted	0.00	NA NA	NA NA	NA NA	0.00	NA NA
Upland Right-of-Way Undeveloped	Available	3.10	NA	NA NA	NA NA	3.10	NA NA
	Impacted	0.70	NA NA	NA NA	NA NA	NA	NA NA
Stormwater Basin	Available	6.95	NA	NA NA	NA NA	NA NA	NA NA
		86.57	NA NA	NA NA	NA NA	NA NA	NA NA
Other Urban or Built-up Land	Impacted Available			NA NA			
		1,194.66	NA NA		NA NA	NA NA	NA NA
Managed Wetland in Maintained Lawn Greenspace	Impacted	0.00	NA	NA NA	NA NA	NA NA	NA NA
	Available	4.78	NA NA	NA NA	NA NA	NA NA	NA NA
Athletic Fields (Schools)	Impacted	0.00	NA	NA	NA	NA	NA
	Available	11.30	NA	NA	NA	NA	NA NA
Recreational Land	Impacted	6.03	NA	NA	NA	NA	NA
	Available	249.30	NA	NA	NA	NA	NA
Agricultural Wetlands (Modified)	Impacted	0.00	NA	NA	NA	NA	NA
, ,	Available	0.00	NA	NA	NA	NA	NA
Cropland and Pastureland	Impacted	0.00	NA	0.00	NA	NA	NA
'	Available	0.01	NA	0.01	NA	NA	NA
Deciduous Forest (10-50% Crown Closure)	Impacted	2.67	NA	NA	NA	NA	NA
	Available	23.08	NA	NA	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted	0.00	NA	NA	0.00	NA	NA
200000000000000000000000000000000000000	Available	114.79	NA	NA	114.79	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted	0.04	NA	NA	NA	NA	NA
Connectors to rest (20 50% crown crossine)	Available	2.72	NA	NA	NA	NA	NA
Coniferous Forest (>50% Crown Closure)	Impacted	0.00	NA	NA	0.00	NA	NA
Communication (1 5075 Crottin Closure)	Available	10.32	NA	NA	10.32	NA	NA
Plantation	Impacted	0.00	NA	0.00	NA	NA	NA
	Available	0.00	NA	0.00	NA	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
INITIALE POLEST (> 30% CONNECTORS WITH TO-30% CLOWN CHOSULE)	Available	2.01	NA	NA	NA	NA	NA
Mixed Forest (> E00/ Coniference with > E00/ Creams (1-200)	Impacted	0.00	NA	NA	0.00	NA	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Available	5.17	NA	NA	5.17	NA	NA
Mind Found (S FOO) Decidence with 40 FOO( Committee of the committee of th	Impacted	0.00	NA	NA	NA	NA	NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Available	3.45	NA	NA	NA	NA	NA
AND A STATE OF THE	Impacted	0.00	NA	NA	0.00	NA	NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Available	2.57	NA	NA	2.57	NA	NA
	Impacted	5.72	NA	5.72	NA NA	5.72	NA NA
Old Field (< 25% Brush Covered)	Available	30.56	NA	30.56	NA	30.56	NA NA
	Impacted	1.41	NA NA	NA	NA NA	NA	NA NA
Deciduous Brush/Shrubland	Available	13.30	NA	NA NA	NA NA	NA NA	NA NA
	Impacted	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Coniferous Brush/Shrubland	· ·						
	Available	0.00	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Podilymbus podiceps	Pooecetes gramineus	Strix varia	Atrytone arogos arogos	Bolaria selene myrina
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
Ivilxed Deciduous/Connerous Brushly Shrubland	Available	24.11	NA	NA	NA	NA	NA
Streams and Canals	Impacted	3.43	NA	NA	NA	NA	NA
Streams and Canais	Available	45.00	NA	NA	NA	NA	NA
Natural Lakes	Impacted	0.00	0.00	NA	NA	NA	NA
ivaturar Lakes	Available	0.00	0.00	NA	NA	NA	NA
Artificial Lakes	Impacted	0.37	0.37	NA	NA	NA	NA
Artificial Lakes	Available	1.88	1.88	NA	NA	NA	NA
Deciduous Wooded Wetlands	Impacted	2.26	2.26	NA	NA	NA	NA
Deciduous wooded wetlands	Available	75.92	75.92	NA	NA	NA	NA
Coniferous Wooded Wetlands	Impacted	0.62	0.62	NA	NA	NA	NA
Connerous wooded wedands	Available	4.15	4.15	NA	NA	NA	NA
Atlantic White Cedar Wetlands	Impacted	0.00	0.00	NA	NA	NA	NA
Additic write cedar wedands	Available	0.00	0.00	NA	NA	NA	NA
Deciduous Scrub/Shrub Wetlands	Impacted	0.00	0.00	NA	NA	NA	0.00
Deciduous Scrub/Siliub Wetlands	Available	0.64	0.64	NA	NA	NA	0.64
Coniferous Scrub/Shrub Wetlands	Impacted	0.00		NA	NA	NA	0.00
Connerous Scrub/Siliub Wetlands	Available	0.00		NA	NA	NA	0.00
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	0.00	NA	NA	NA	0.00
Wettands (Decidadus Donn)	Available	2.75	2.75	NA	NA	NA	2.75
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	0.00	NA	NA	NA	0.00
Timed Sci as John as Wellands (Colliner ods Scinity	Available	0.00	0.00	NA	NA	NA	0.00
Herbaceous Wetlands	Impacted	54.96	54.96	NA	NA	NA	54.96
	Available	379.58	379.58	NA	NA	NA	379.58
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA	0.00	NA	NA
	Available	5.56	NA	NA	5.56	NA	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	0.00	NA	NA
,	Available	0.23	NA	NA	0.23	NA	NA
Disturbed Wetlands (Modified)	Impacted	1.46	NA	NA	NA	NA	NA
	Available	13.80	NA	NA NA	NA	NA	NA
Transitional Areas	Impacted	0.84	NA	NA	NA	NA	NA
	Available	3.91	NA NA	NA 2.22	NA NA	NA	NA
Undifferentiated Barren Lands	Impacted	0.00	NA	0.00	NA NA	NA NA	NA NA
	Available	0.00	NA 0.00	0.00	NA NA	NA NA	NA NA
Phragmites Dominant Interior Wetlands	Impacted	0.00	0.00	NA NA	NA NA	NA NA	NA NA
	Available 0.00  Species Habitat Impacted (ac)		0.00	NA F 72	NA 0.00	NA 5.72	NA 54.06
			58.20	5.72	0.00	5.72	54.96
	Species Habitat Availab Percent Impacted (	• •	464.91	30.57	138.63	33.66	382.96
			12.5186%	18.7205%	0.0000%	17.0029%	14.3516%
	Percent Available,	Not impacted	87.4814%	81.2795%	100.0000%	82.9971%	85.6484%

ac = acres

Feature Description	Habitat Status	Area (Ac)	Callophrys irus	Lynx rufus	Clemmys insculpta	Clemmys muhlenbergii	Crotalus horridus
Cemetery	Impacted	0.00	NA	NA	NA	NA	NA
Cemetery	Available	0.01	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Impacted	0.00	NA	NA	NA	NA	NA
Residential, high Density of Multiple Dwelling	Available	0.00	NA	NA	NA	NA	NA
Desidential Donal Circle Heit	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Available	0.00	NA	NA	NA	NA	NA
	Impacted	0.00	NA	NA	NA	NA	NA
Commercial/Services	Available	1.31	NA	NA	NA	NA	NA
	Impacted	178.12	NA	NA	NA	NA	NA
Military Installations	Available	1,317.90	NA	NA NA	NA NA	NA	NA
	Impacted	0.05	NA NA	NA NA	NA NA	NA NA	NA NA
Transportation/Communication/Utilities	Available	5.11	NA	NA NA	NA NA	NA NA	NA NA
	Impacted	0.00	0.00	NA NA	NA NA	NA NA	NA NA
Upland Right-of-Way Undeveloped	Available	3.10	3.10	NA NA	NA NA	NA NA	NA NA
		0.70		NA NA	NA NA	NA NA	NA NA
Stormwater Basin	Impacted		NA NA				
	Available	6.95	NA NA	NA NA	NA NA	NA NA	NA NA
Other Urban or Built-up Land	Impacted	86.57	NA	NA NA	NA	NA NA	NA NA
	Available	1,194.66	NA	NA	NA	NA	NA
Managed Wetland in Maintained Lawn Greenspace	Impacted	0.00	NA	NA	NA	NA	NA
	Available	4.78	NA	NA	NA	NA	NA
Athletic Fields (Schools)	Impacted	0.00	NA	NA	NA	NA	NA
	Available	11.30	NA	NA	NA	NA	NA
Recreational Land	Impacted	6.03	NA	NA	NA	NA	NA
	Available	249.30	NA	NA	NA	NA	NA
Agricultural Wetlands (Modified)	Impacted	0.00	NA	NA	NA	NA	NA
- Butananan menanas (meanica)	Available	0.00	NA	NA	NA	NA	NA
Cropland and Pastureland	Impacted	0.00	NA	0.00	NA	NA	NA
eropiana ana rastarelana	Available	0.01	NA	0.01	NA	NA	NA
Deciduous Forest (10-50% Crown Closure)	Impacted	2.67	NA	NA	NA	NA	NA
Deciduous Forest (10-30% Crown Closure)	Available	23.08	NA	NA	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Available	114.79	NA	NA	NA	NA	NA
0 - 16	Impacted	0.04	NA	0.04	NA	NA	0.04
Coniferous Forest (10-50% Crown Closure)	Available	2.72	NA	2.72	NA	NA	2.72
0	Impacted	0.00	NA	0.00	NA	NA	0.00
Coniferous Forest (>50% Crown Closure)	Available	10.32	NA	10.32	NA	NA	10.32
	Impacted	0.00	NA	NA NA	NA	NA NA	NA
Plantation	Available	0.00	NA	NA	NA NA	NA NA	NA
	Impacted	0.00	NA NA	0.00	NA NA	NA NA	0.00
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Available	2.01	NA NA	2.01	NA NA	NA NA	2.01
	Impacted	0.00	NA NA	0.00	NA NA	NA NA	0.00
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Available	5.17	NA NA	5.17	NA NA	NA NA	5.17
		0.00	NA NA	0.00	NA NA	NA NA	0.00
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted Available	3.45		3.45			3.45
			NA NA		NA NA	NA NA	
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted	0.00	NA	0.00	NA NA	NA NA	0.00
	Available	2.57	NA 5.72	2.57	NA NA	NA NA	2.57
Old Field (< 25% Brush Covered)	Impacted	5.72	5.72	NA NA	NA	NA NA	NA NA
•	Available	30.56	30.56	NA	NA	NA 	NA
Deciduous Brush/Shrubland	Impacted	1.41	NA	NA	NA	NA	NA
,	Available	13.30	NA	NA	NA	NA	NA
Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Callophrys irus	Lynx rufus	Clemmys insculpta	Clemmys muhlenbergii	Crotalus horridus
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
Ivilked Deciduous/Connerous Brushi/Sin ubiand	Available	24.11	NA	NA	NA	NA	NA
Streams and Canals	Impacted	3.43	NA	NA	3.43	NA	NA
Streams and Canais	Available	45.00	NA	NA	45.00	NA	NA
Natural Lakes	Impacted	0.00	NA	NA	0.00	NA	NA
inaturar Lakes	Available	0.00	NA	NA	0.00	NA	NA
Artificial Lakes	Impacted	0.37	NA	NA	NA	NA	NA
Artificial takes	Available	1.88	NA	NA	NA	NA	NA
Deciduous Wooded Wetlands	Impacted	2.26	NA	NA	2.26	NA	NA
Deciduous wooded wettands	Available	75.92	NA	NA	75.92	NA	NA
Coniferous Wooded Wetlands	Impacted	0.62	NA	NA	0.62	NA	0.62
Connerous wooded wedands	Available	4.15	NA	NA	4.15	NA	4.15
Atlantic White Cedar Wetlands	Impacted	0.00	NA	NA	NA	NA	0.00
Additic white cedar weddings	Available	0.00	NA	NA	NA	NA	0.00
Deciduous Scrub/Shrub Wetlands	Impacted	0.00	NA	NA	0.00	0.00	NA
Deciduous Scrub/Siliub Wetlands	Available	0.64	NA	NA	0.64	0.64	NA
Coniferous Scrub/Shrub Wetlands	Impacted	0.00	NA	NA	0.00	0.00	NA
connerous scrub/smub wetlands	Available	0.00	NA	NA	0.00	0.00	NA
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA	0.00	0.00	NA
Wincu Scrub/Siliub Wetianus (Deciduous Doili.)	Available	2.75	NA	NA	2.75	2.75	NA
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	0.00	0.00	NA
Trinca serab/sirab tretianas (connercus bonn)	Available	0.00	NA	NA	0.00	0.00	NA
Herbaceous Wetlands	Impacted	54.96	NA	NA	54.96	54.96	NA
	Available	379.58	NA	NA	379.58	379.58	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA	0.00	NA	NA
,	Available	5.56	NA	NA	5.56	NA	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	0.00	NA	0.00
,	Available	0.23	NA	NA	0.23	NA	0.23
Disturbed Wetlands (Modified)	Impacted	1.46	NA	NA	NA	NA	NA
	Available	13.80	NA	NA	NA	NA	NA 
Transitional Areas	Impacted	0.84	NA	NA NA	NA	NA NA	NA NA
	Available	3.91	NA NA	NA NA	NA NA	NA NA	NA NA
Undifferentiated Barren Lands	Impacted	0.00	NA	NA NA	NA NA	NA NA	NA NA
	Available	0.00	NA NA	NA NA	NA 0.00	NA 0.00	NA NA
Phragmites Dominant Interior Wetlands	Impacted	0.00 0.00	NA NA	NA NA		0.00	NA NA
	Available Species Habitat		NA 5.72	NA 0.04	0.00	0.00	NA 0.66
	Species Habitat Availab		5.72	0.04	61.27	54.96	0.66
	Percent Impacted (		33.66 17.0029%	26.25 0.1572%	513.83	382.96	30.62 2.1468%
	Percent Impacted C				11.9238%	14.3516%	
	reitent Available,	not impacted	82.9971%	99.8428%	88.0762%	85.6484%	97.8532%

ac = acres

Feature Description	Habitat Status	Area (Ac)	Elaphe guttata	Pituophis m melanoleucus
Compatonic	Impacted	0.00	NA	NA
Cemetery	Available	0.01	NA	NA
Residential, High Density or Multiple Dwelling	Impacted	0.00	NA	NA
Residential, Fight Density of Multiple Dweiling	Available	0.00	NA	NA
Residential, Rural, Single Unit	Impacted	0.00	NA	NA
Residential, Rural, Single Offic	Available	0.00	NA	NA
Residential, Single Unit, Low Density	Impacted	0.00	NA	NA
nesidential, single only zon bensity	Available	0.00	NA	NA
Residential, Single Unit, Medium Density	Impacted	0.00	NA	NA
	Available	0.00	NA	NA
Commercial/Services	Impacted	0.00	NA	NA
	Available	1.31	NA	NA
Military Installations	Impacted	178.12	NA NA	NA NA
	Available	1,317.90	NA NA	NA NA
Transportation/Communication/Utilities	Impacted	0.05	NA NA	NA NA
	Available	5.11 0.00	NA NA	NA NA
Upland Right-of-Way Undeveloped	Impacted Available	3.10	NA NA	NA NA
	Impacted	0.70	NA NA	NA NA
Stormwater Basin	Available	6.95	NA NA	NA NA
	Impacted	86.57	NA NA	NA NA
Other Urban or Built-up Land	Available	1,194.66	NA NA	NA NA
	Impacted	0.00	NA NA	NA NA
Managed Wetland in Maintained Lawn Greenspace	Available	4.78	NA	NA NA
	Impacted	0.00	NA NA	NA NA
Athletic Fields (Schools)	Available	11.30	NA NA	NA
	Impacted	6.03	NA	NA NA
Recreational Land	Available	249.30	NA	NA
	Impacted	0.00	NA	NA
Agricultural Wetlands (Modified)	Available	0.00	NA	NA
Cuantand and Bastowaland	Impacted	0.00	NA	NA
Cropland and Pastureland	Available	0.01	NA	NA
Deciduous Forest (10-50% Crown Closure)	Impacted	2.67	NA	2.67
Decidious Polest (10-30% Clowii Closule)	Available	23.08	NA	23.08
Deciduous Forest (>50% Crown Closure)	Impacted	0.00	NA	NA
Decidadus Forest (>30% Crown Closure)	Available	114.79	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted	0.04	NA	0.04
	Available	2.72	NA	2.72
Coniferous Forest (>50% Crown Closure)	Impacted	0.00	0.00	NA
,	Available	10.32	10.32	NA 
Plantation	Impacted	0.00	NA NA	NA NA
	Available	0.00	NA NA	NA 0.00
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted	0.00	NA NA	0.00
	Available	2.01	NA 0.00	2.01
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted	0.00	0.00 5.17	NA NA
	Available	5.17 0.00	5.17 NA	NA 0.00
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted Available	3.45	NA NA	3.45
	Impacted	0.00	NA NA	3.45 NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Available	2.57	NA NA	NA NA
	Impacted	5.72	NA NA	5.72
Old Field (< 25% Brush Covered)	Available	30.56	NA NA	30.56
	Impacted	1.41	NA NA	NA
Deciduous Brush/Shrubland	Available	13.30	NA NA	NA NA
	Impacted	0.00	NA	NA NA
Coniferous Brush/Shrubland	Available	0.00	NA	NA NA
	Available	0.00	IVA	I IVA

Feature Description	Habitat Status	Area (Ac)	Elaphe guttata	Pituophis m melanoleucus		
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	0.00	NA	NA		
ivilxed Deciduous/Confletous Brush/Shrubland	Available	24.11	NA	NA		
Change and Canala	Impacted	3.43	NA	NA		
Streams and Canals	Available	45.00	NA	NA		
Natural Lakes	Impacted	0.00	NA	NA		
Natural Lakes	Available	0.00	NA	NA		
Autificial Labora	Impacted	0.37	NA	NA		
Artificial Lakes	Available	1.88	NA	NA		
Desidueus Weeded Wetlende	Impacted	2.26	NA	NA		
Deciduous Wooded Wetlands	Available	75.92	NA	NA		
Coulfornia Mandad Matlanda	Impacted	0.62	0.62	NA		
Coniferous Wooded Wetlands	Available	4.15	4.15	NA		
Address: Nathita Cardan Maddan da	Impacted	0.00	0.00	NA		
Atlantic White Cedar Wetlands	Available	0.00	0.00	NA		
Desidueus Comub (Chaub Matlanda	Impacted	0.00	NA	NA		
Deciduous Scrub/Shrub Wetlands	Available	0.64	NA	NA		
Coniferous Scrub/Shrub Wetlands	Impacted	0.00	0.00	NA		
Connerous Scrub/Siliub Wetlands	Available	0.00	0.00	NA		
Mixed Scrub/Shrub Wetlands (Deciduous Dom )	Impacted	0.00	NA	NA		
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Available	2.75	NA	NA		
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	0.00	NA		
Wince Screen Street Westernes (Connerous Bonn)	Available	0.00	0.00	NA		
Herbaceous Wetlands	Impacted	54.96	NA	NA		
	Available	379.58	NA	NA		
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA		
,	Available	5.56	NA	NA		
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted	0.00	0.00	NA		
,	Available	0.23	0.23	NA NA		
Disturbed Wetlands (Modified)	Impacted	1.46	NA 	NA 		
	Available	13.80	NA NA	NA NA		
Transitional Areas	Impacted	0.84	NA NA	NA NA		
	Available	3.91	NA 0.00	NA 0.00		
Undifferentiated Barren Lands	Impacted	0.00	0.00	0.00		
	Available	0.00	0.00	0.00		
Phragmites Dominant Interior Wetlands	Impacted Available		NA NA	NA NA		
	Species Habitat I	0.00	NA 0.62	NA 9.44		
	Species Habitat Available	•		8.44 61.83		
	•			13.6478%		
	Percent Impacted Over Available Percent Available, Not Impacted					
	reiteilt Avallable, l	iot iiipacted	96.8998%	86.3522%		

ac = acres

Table B-2. Spatial Exposure Potential of Protected Species at Dix of Joint Base McGuire-Dix-Lakehurst

Feature Description	Habitat Status	Area (Ac)	Clemmy guttata	Pseudemys rubriventris (Juvenile)	Pseudemys rubriventris (Adult)	Hyla andersonii	Hyla chrysocelis
Comotory	Impacted	0.00	NA	NA	NA	NA	NA
Cemetery	Available	3.47	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Impacted	0.00	NA	NA	NA	NA	NA
residential, riight bensity of Martiple bwening	Available	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Impacted	0.00	NA	NA	NA	NA	NA
nesidential, naral, single sinc	Available	9.13	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Impacted	0.00	NA	NA	NA	NA	NA
The state of the second of the	Available	0.20	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Impacted	0.00	NA	NA	NA	NA	NA
	Available	1.05	NA	NA	NA	NA	NA
Commercial/Services	Impacted	0.00	NA	NA	NA	NA	NA
	Available	0.42	NA	NA	NA	NA	NA
Military Installations	Impacted	31.35	NA	NA	NA	NA	NA
,	Available	3,132.47	NA	NA	NA	NA	NA
Transportation/Communication/Utilities	Impacted	1.36	NA	NA	NA	NA	NA
	Available	220.36	NA	NA	NA	NA	NA
Upland Right-of-Way Undeveloped	Impacted	0.00	NA	NA	NA	NA	NA
	Available	81.09	NA	NA	NA	NA	NA
Stormwater Basin	Impacted	0.00	NA	NA	NA	NA	0.00
	Available	33.27	NA	NA	NA	NA	33.27
Other Urban or Built-up Land	Impacted	16.61	NA	NA	NA	NA	NA
	Available	1,250.32	NA	NA	NA	NA	NA
Managed Wetland in Maintained Lawn Greenspace	Impacted	0.00	NA	0.00	NA	NA	0.00
	Available	4.17	NA	4.17	NA	NA	4.17
Athletic Fields (Schools)	Impacted	0.00	NA	NA	NA	NA	NA
	Available	2.51	NA	NA	NA	NA	NA
Recreational Land	Impacted	114.81	NA	NA	NA	NA	NA
	Available	429.81	NA	NA	NA	NA	NA
Agricultural Wetlands (Modified)	Impacted	92.84	NA	92.84	NA	NA	92.84
<b>3</b> ,	Available	93.97	NA	93.97	NA	NA	93.97
Cropland and Pastureland	Impacted	0.00	NA	0.00	NA	NA	0.00
	Available	14.17	NA	14.17	NA	NA	14.17
Deciduous Forest (10-50% Crown Closure)	Impacted	2.46	NA	NA	NA	NA	NA
,	Available	297.32	NA	NA	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted	39.15	NA	NA	NA	NA	NA
,	Available	1,522.35	NA	NA	NA	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
	Available	3,476.78	NA	NA	NA	NA	NA
Coniferous Forest (>50% Crown Closure)	Impacted	7.15	NA	NA	NA	NA	NA
	Available	6,216.02	NA	NA	NA	NA	NA
Plantation	Impacted	10.95	NA	NA	NA	NA	NA
	Available	341.48	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
,	Available	209.19	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted	34.29	NA	NA	NA	NA	NA
,	Available	1,819.31	NA	NA	NA	NA	NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
	Available	100.33	NA	NA	NA	NA	NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted	10.93	NA	NA	NA	NA	NA
	Available	865.23	NA	NA	NA	NA	NA
Old Field (< 25% Brush Covered)	Impacted	1.19	NA	NA	NA	NA	NA
	Available	830.21	NA	NA	NA	NA	NA
Deciduous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
	Available	78.60	NA	NA	NA	NA	NA
Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
	Available	679.92	NA	NA	NA	NA	NA
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	5.77	NA	NA	NA	NA	NA
Trinca Decidadas, connerous Drasil, Sili abiana	Available	734.89	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Clemmy guttata	Pseudemys rubriventris (Juvenile)	Pseudemys rubriventris (Adult)	Hyla andersonii	Hyla chrysocelis
Streams and Canals	Impacted	6.73	6.73	NA	6.73	NA	NA
Streams and Canais	Available	452.34	452.34	NA	452.34	NA	NA
Natural Lakes	Impacted	0.00	0.00	0.00	0.00	NA	NA
ivaturar Lakes	Available	4.54	4.54	4.54	4.54	NA	NA
Artificial Lakes	Impacted	7.17	7.17	7.17	7.17	NA	NA
Al tilitial takes	Available	396.28	396.28	396.28	396.28	NA	NA
Deciduous Wooded Wetlands	Impacted	9.80	9.80	9.80	NA	NA	9.80
Deciduous wooded wetiailus	Available	2,334.37	2,334.37	2,334.37	NA	NA	2,334.37
Coniferous Wooded Wetlands	Impacted	0.00	0.00	0.00	NA	0.00	0.00
Connerous wooded wetlands	Available	1,607.30	1,607.30	1,607.30	NA	1,607.30	1,607.30
Atlantic White Cedar Wetlands	Impacted	0.00	0.00	0.00	NA	0.00	0.00
Atlantic White Cedar Wetlands	Available	298.97	298.97	298.97	NA	298.97	298.97
Deciduous Scrub/Shrub Wetlands	Impacted	0.00	0.00	0.00	NA	0.00	0.00
Deciduous Scrub/Shrub Wetlands	Available	371.69	371.69	371.69	NA	371.69	371.69
Coniference Servich /Shruch Westlands	Impacted	0.00	0.00	0.00	NA	0.00	0.00
Coniferous Scrub/Shrub Wetlands	Available	292.71	292.71	292.71	NA	292.71	292.71
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	0.00	0.00	NA	0.00	0.00
ivilxed Scrub/Silrub Wetlands (Deciduous Doni.)	Available	635.15	635.15	635.15	NA	635.15	635.15
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	0.00	0.00	NA	0.00	0.00
ivilked Scrub/Siliub Wetlands (Confferous Dom.)	Available	101.91	101.91	101.91	NA	101.91	101.91
Herbaceous Wetlands	Impacted	0.74	0.74	0.74	NA	0.74	0.74
Herbaceous wetlanus	Available	843.63	843.63	843.63	NA	843.63	843.63
Mixed Wooded Wetlands (Desidueus Dem.)	Impacted	12.13	12.13	12.13	NA	12.13	12.13
Mixed Wooded Wetlands (Deciduous Dom.)	Available	523.83	523.83	523.83	NA	523.83	523.83
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted	0.00	0.00	0.00	NA	0.00	0.00
winked wooded wettands (Conferous Dom.)	Available	635.50	635.50	635.50	NA	635.50	635.50
Disturbed Wetlands (Modified)	Impacted	0.00	NA	0.00	NA	NA	0.00
Disturbed Wetlands (Modified)	Available	47.03	NA	47.03	NA	NA	47.03
Transitional Areas	Impacted	27.51	NA	NA	NA	NA	NA
Transitional Areas	Available	166.61	NA	NA	NA	NA	NA
Undifferentiated Barren Lands	Impacted	0.00	NA	NA	NA	NA	NA
Ondinerentiated Barren Lands	Available	4.34	NA	NA	NA	NA	NA
Phragmites Dominant Interior Wetlands	Impacted	0.00	NA	NA	NA	NA	NA
rmagnines Dominant interior wetlands	Available	2.67	NA	NA	NA	NA	NA
	Species Habitat In	npacted (ac)	36.58	122.69	13.90	12.88	115.52
	Species Habitat Available	Overall (ac)	8,498.20	8,205.19	853.17	5,310.67	7,837.64
	Percent Impacted Ov	er Available	0.4304%	1.4952%	1.6296%	0.2424%	1.4739%
	Percent Available, N	ot Impacted	99.5696%	98.5048%	98.3704%	99.7576%	98.5261%

ac = acres

Feature Description	Habitat Status	Area (Ac)	Pseudotriton montanus	Accipiter gentilis	Ammodramus henslowii	Ammodramus savannarum	Asio flammeus
Comotory	Impacted	0.00	NA	NA	NA	NA	0.00
Cemetery	Available	3.47	NA	NA	NA	NA	3.47
Residential, High Density or Multiple Dwelling	Impacted	0.00	NA	NA	NA	NA	NA
Residential, High Density of Multiple Dwelling	Available	0.00	NA	NA	NA	NA	NA
Desidential Dural Cingle Unit	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Available	9.13	NA	NA	NA	NA	NA
Posidontial Single Unit Law Density	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Available	0.20	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Onit, Medium Density	Available	1.05	NA	NA	NA	NA	NA
Commercial/Services	Impacted	0.00	NA	NA	NA	NA	NA
Commercial/Services	Available	0.42	NA	NA	NA	NA	NA
Military Installations	Impacted	31.35	NA	NA	NA	NA	NA
ivilitary installations	Available	3,132.47	NA	NA	NA	NA	NA
Transportation/Communication/Utilities	Impacted	1.36	NA	NA	NA	NA	NA
Transportation/Communication/Othities	Available	220.36	NA	NA	NA	NA	NA
Upland Right-of-Way Undeveloped	Impacted	0.00	NA	NA	0.00	0.00	NA
Opiana Right-or-way Ondeveloped	Available	81.09	NA	NA	81.09	81.09	NA
Stormwater Basin	Impacted	0.00	NA	NA	NA	NA	NA
Stormwater Dasin	Available	33.27	NA	NA	NA	NA	NA
Other Urban or Built-up Land	Impacted	16.61	NA	NA	NA	NA	16.61
Other Orban or Built-up Land	Available	1,250.32	NA	NA	NA	NA	1,250.32
Managed Wetland in Maintained Lawn Greenspace	Impacted	0.00	NA	NA	NA	NA	NA
Managed Wetland III Manitanied Lawn Greenspace	Available	4.17	NA	NA	NA	NA	NA
Athletic Fields (Schools)	Impacted	0.00	NA	NA	NA	NA	NA
Athletic Fields (Schools)	Available	2.51	NA	NA	NA	NA	NA
Recreational Land	Impacted	114.81	NA	NA	NA	NA	NA
Necreational Land	Available	429.81	NA	NA	NA	NA	NA
Agricultural Wetlands (Modified)	Impacted	92.84	NA	NA	NA	NA	NA
Agricultural Wetlands (Woulfied)	Available	93.97	NA	NA	NA	NA	NA
Cropland and Pastureland	Impacted	0.00	NA	NA	NA	NA	0.00
cropiana ana rastarciana	Available	14.17	NA	NA	NA	NA	14.17
Deciduous Forest (10-50% Crown Closure)	Impacted	2.46	NA	NA	NA	NA	NA
Deciduous Forest (10-50% Crown Closure)	Available	297.32	NA	NA	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted	39.15	NA	39.15	NA	NA	NA
Deciduous Forest (>30% crown closure)	Available	1,522.35	NA	1,522.35	NA	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
connectous forest (10-30% crown closure)	Available	3,476.78	NA	NA	NA	NA	NA
Coniferous Forest (>50% Crown Closure)	Impacted	7.15	NA	7.15	NA	NA	NA
connerous rorest (230% crown closure)	Available	6,216.02	NA	6,216.02	NA	NA	NA
Plantation	Impacted	10.95	NA	NA	NA	NA	NA
Tuntation	Available	341.48	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
Trinked Forest (* 50% Commercus With 10 50% Crown Closure)	Available	209.19	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted	34.29	NA	34.29	NA	NA	NA
The state of the s	Available	1,819.31	NA	1,819.31	NA	NA	NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
	Available	100.33	NA	NA	NA	NA	NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted	10.93	NA	10.93	NA	NA	NA
	Available	865.23	NA	865.23	NA	NA	NA
Old Field (< 25% Brush Covered)	Impacted	1.19	NA	NA	1.19	1.19	1.19
	Available	830.21	NA	NA	830.21	830.21	830.21
Deciduous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
	Available	78.60	NA	NA	NA	NA	NA
Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
Come. Sas Diasily Sili abidila	Available	679.92	NA	NA	NA	NA	NA
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	5.77	NA	NA	NA	NA	NA
minea bediadous, comicious brasily simubiana	Available	734.89	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Pseudotriton montanus	Accipiter gentilis	Ammodramus henslowii	Ammodramus savannarum	Asio flammeus
Streams and Canals	Impacted	6.73	NA	NA	NA	NA	NA
Streams and Canais	Available	452.34	NA	NA	NA	NA	NA
Natural Lakes	Impacted	0.00	0.00	NA	NA	NA	NA
ivaturar Lakes	Available	4.54	4.54	NA	NA	NA	NA
Artificial Lakes	Impacted	7.17	7.17	NA	NA	NA	NA
Ai tiliciai Lakes	Available	396.28	396.28	NA	NA	NA	NA
Deciduous Wooded Wetlands	Impacted	9.80	NA	9.80	NA	NA	NA
Decidadas Froducti Freduitas	Available	2,334.37	NA	2,334.37	NA	NA	NA
Coniferous Wooded Wetlands	Impacted	0.00	NA	0.00	NA	NA	NA
Connerous wooded wetlands	Available	1,607.30	NA	1,607.30	NA	NA	NA
Atlantic White Cedar Wetlands	Impacted	0.00	0.00	0.00	NA	NA	NA
Atlantic White Cedar Wetlands	Available	298.97	298.97	298.97	NA	NA	NA
Deciduous Scrub/Shrub Wetlands	Impacted	0.00	NA	NA	NA	NA	NA
Deciduous Scrub/Siliub Wetialius	Available	371.69	NA	NA	NA	NA	NA
Coniferous Scrub/Shrub Wetlands	Impacted	0.00	NA	NA	NA	NA	NA
Connerous scrub/shrub Wetlands	Available	292.71	NA	NA	NA	NA	NA
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA	NA	NA	NA
Wilked Scrab/Siliab Wetlands (Deciduous Dolli.)	Available	635.15	NA	NA	NA	NA	NA
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	NA	NA	NA
wettands (Connerous Donn.)	Available	101.91	NA	NA	NA	NA	NA
Harbacoous Wotlands	Impacted	0.74	0.74	NA	0.74	NA	NA
herbaceous wetianus	Available	843.63	843.63	NA	843.63	NA	NA
Mixed Wooded Wetlands (Posiduous Dom.)	Impacted	12.13	NA	12.13	NA	NA	NA
wilked wooded wetlands (Deciduous Doni.)	Available	523.83	NA	523.83	NA	NA	NA
Miyad Waadad Watlands (Canifornia Dom.)	Impacted	0.00	NA	0.00	NA	NA	NA
ixed Wooded Wetlands (Deciduous Dom.) ixed Wooded Wetlands (Coniferous Dom.) sturbed Wetlands (Modified)	Available	635.50	NA	635.50	NA	NA	NA
Disturbed Wetlands (Medified)	Impacted	0.00	0.00	NA	NA	NA	NA
Disturbed Wetianus (Modined)	Available	47.03	47.03	NA	NA	NA	NA
Transitional Areas	Impacted	27.51	NA	NA	NA	NA	NA
Transitional Areas	Available	166.61	NA	NA	NA	NA	NA
Undifferentiated Barren Lands	Impacted	0.00	NA	NA	0.00	0.00	NA
Ondinerentiated Barren Lands	Available	4.34	NA	NA	4.34	4.34	NA
Phragmitos Dominant Interior Wetlands	Impacted	0.00	NA	NA	NA	NA	0.00
Phragmites Dominant Interior Wetlands	Available	2.67	NA	NA	NA	NA	2.67
	Species Habitat I	mpacted (ac)	7.91	113.45	1.93	1.19	17.80
	Species Habitat Available		1,590.44	15,822.86	1,759.26	915.64	2,100.84
	Percent Impacted O	ver Available	0.4976%	0.7170%	0.1097%	0.1297%	0.8474%
	Percent Available, N	Not Impacted	99.5024%	99.2830%	99.8903%	99.8703%	99.1526%

ac = acres

Feature Description	Habitat Status	Area (Ac)	Asio otus	Batramia longicauda	Botaurus lentiginosos	Buteo lineatus	Circus cyaneus
Cometany	Impacted	0.00	0.00	0.00	NA	NA	0.00
Cemetery	Available	3.47	3.47	3.47	NA	NA	3.47
Residential, High Density or Multiple Dwelling	Impacted	0.00	NA	NA	NA	NA	NA
nesidential, nigh behisty of Multiple bwelling	Available	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Impacted	0.00	NA	NA	NA	NA	NA
nesidential, rural, single offit	Available	9.13	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Offic, Low Density	Available	0.20	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Impacted	0.00	NA	NA	NA	NA	NA
nestachtal, single sint, mediani sensity	Available	1.05	NA	NA	NA	NA	NA
Commercial/Services	Impacted	0.00	NA	NA	NA	NA	NA
	Available	0.42	NA	NA	NA	NA	NA
Military Installations	Impacted	31.35	NA	NA	NA	NA	NA
,	Available	3,132.47	NA	NA	NA	NA	NA
Transportation/Communication/Utilities	Impacted	1.36	NA	NA	NA	NA	NA
, ,	Available	220.36	NA	NA	NA	NA	NA
Upland Right-of-Way Undeveloped	Impacted	0.00	NA	0.00	NA	NA	0.00
, , , , , , ,	Available	81.09	NA	81.09	NA	NA	81.09
Stormwater Basin	Impacted	0.00	NA	NA	NA 	NA	NA 
	Available	33.27	NA	NA	NA	NA	NA
Other Urban or Built-up Land	Impacted	16.61	NA	NA	NA	NA	16.61
	Available	1,250.32	NA	NA	NA	NA	1,250.32
Managed Wetland in Maintained Lawn Greenspace	Impacted	0.00	NA	NA	NA	NA	0.00
	Available	4.17	NA	NA	NA	NA	4.17
Athletic Fields (Schools)	Impacted	0.00	NA	NA	NA	NA	0.00
,	Available	2.51	NA	NA	NA	NA	2.51
Recreational Land	Impacted	114.81	NA	NA	NA	NA	NA 
	Available	429.81	NA	NA	NA	NA	NA
Agricultural Wetlands (Modified)	Impacted	92.84	NA	NA	NA 	NA	92.84
	Available	93.97	NA	NA 0.00	NA NA	NA NA	93.97
Cropland and Pastureland	Impacted	0.00	0.00	0.00	NA NA	NA	0.00
	Available	14.17	14.17	14.17	NA NA	NA NA	14.17
Deciduous Forest (10-50% Crown Closure)	Impacted	2.46	2.46	NA NA	NA NA	NA	NA NA
	Available	297.32	297.32	NA NA	NA NA	NA 20.15	NA NA
Deciduous Forest (>50% Crown Closure)	Impacted	39.15	39.15	NA NA	NA NA	39.15	NA NA
	Available	1,522.35 0.00	1,522.35 0.00	NA NA	NA NA	1,522.35 NA	NA NA
Coniferous Forest (10-50% Crown Closure)	Impacted Available						
		3,476.78 7.15	3,476.78 7.15	NA NA	NA NA	NA 7.15	NA NA
Coniferous Forest (>50% Crown Closure)	Impacted Available	6,216.02	6,216.02	NA NA	NA NA	6,216.02	NA NA
	Impacted	10.95	0,216.02 NA	10.95	NA NA	0,210.02 NA	NA NA
Plantation	Available	341.48	NA NA	341.48	NA NA	NA NA	NA NA
	Impacted	0.00	0.00	NA	NA NA	NA NA	NA NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Available	209.19	209.19	NA NA	NA NA	NA NA	NA NA
	Impacted	34.29	34.29	NA NA	NA NA	34.29	NA NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Available	1,819.31	1,819.31	NA NA	NA NA	1,819.31	NA NA
	Impacted	0.00	0.00	NA NA	NA NA	1,819.31 NA	NA NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Available	100.33	100.33	NA NA	NA NA	NA NA	NA NA
	Impacted	10.93	10.93	NA NA	NA NA	10.93	NA NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Available	865.23	865.23	NA NA	NA NA	865.23	NA NA
	Impacted	1.19	1.19	1.19	NA NA	NA	1.19
Old Field (< 25% Brush Covered)	Available	830.21	830.21	830.21	NA NA	NA	830.21
	Impacted	0.00	NA	NA	NA NA	NA NA	NA
Deciduous Brush/Shrubland	Available	78.60	NA	NA NA	NA NA	NA	NA NA
	Impacted	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Coniferous Brush/Shrubland	Available	679.92	NA NA	NA	NA NA	NA NA	NA
	Impacted	5.77	NA	NA NA	NA NA	NA NA	NA NA
Mixed Deciduous/Coniferous Brush/Shrubland	Available	734.89	NA	NA	NA	NA	NA
	Available	/34.89	IVA	INA	INA NA	IVA	I NA

Feature Description	Habitat Status	Area (Ac)	Asio otus	Batramia longicauda	Botaurus lentiginosos	Buteo lineatus	Circus cyaneus
Streams and Canals	Impacted	6.73	NA	NA	NA	NA	NA
Streams and Canais	Available	452.34	NA	NA	NA	NA	NA
Natural Lakes	Impacted	0.00	NA	NA	NA	NA	NA
Natural Lakes	Available	4.54	NA	NA	NA	NA	NA
Artificial Lakes	Impacted	7.17	NA	NA	NA	NA	NA
Altificial Lakes	Available	396.28	NA	NA	NA	NA	NA
Deciduous Wooded Wetlands	Impacted	9.80	NA	NA	NA	9.80	NA
Deciduous wooded wetlands	Available	2,334.37	NA	NA	NA	2,334.37	NA
Coniferous Wooded Wetlands	Impacted	0.00	NA	NA	NA	0.00	NA
Connerous wooded wettands	Available	1,607.30	NA	NA	NA	1,607.30	NA
Atlantic White Cedar Wetlands	Impacted	0.00	NA	NA	NA	0.00	NA
Atlantic write Cedar Wetlands	Available	298.97	NA	NA	NA	298.97	NA
Desidueus Comub /Chmub Westlands	Impacted	0.00	NA	NA	0.00	NA	0.00
Deciduous Scrub/Shrub Wetlands	Available	371.69	NA	NA	371.69	NA	371.69
Coniferous Scrub/Shrub Wetlands	Impacted	0.00	NA	NA	0.00	NA	0.00
Connerous Scrub/Shrub Wetlands	Available	292.71	NA	NA	292.71	NA	292.71
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA	0.00	NA	0.00
witked Scrub/Siliub Wetlands (Deciduous Dom.)	Available	635.15	NA	NA	635.15	NA	635.15
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	0.00	NA	0.00
wilked Scrub/Shrub Wetlands (Confilerous Doin.)	Available	101.91	NA	NA	101.91	NA	101.91
Herbaceous Wetlands	Impacted	0.74	NA	NA	0.74	NA	0.74
nerbaceous wedanus	Available	843.63	NA	NA	843.63	NA	843.63
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted	12.13	NA	NA	NA	12.13	NA
wilked wooded wetlands (Deciduous Dom.)	Available	523.83	NA	NA	NA	523.83	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	NA	0.00	NA
witked wooded wettands (conferous born.)	Available	635.50	NA	NA	NA	635.50	NA
Disturbed Wetlands (Modified)	Impacted	0.00	NA	NA	0.00	NA	NA
Disturbed Wetianus (Modined)	Available	47.03	NA	NA	47.03	NA	NA
Transitional Areas	Impacted	27.51	NA	NA	NA	NA	27.51
Transitional Areas	Available	166.61	NA	NA	NA	NA	166.61
Undifferentiated Barren Lands	Impacted	0.00	0.00	0.00	NA	NA	0.00
Ondinerentiated Barren Lanus	Available	4.34	4.34	4.34	NA	NA	4.34
Phragmites Dominant Interior Westlands	Impacted	0.00	NA	NA	NA	NA	0.00
Phragmites Dominant Interior Wetlands	Available	2.67	NA	NA	NA	NA	2.67
	Species Habitat	Impacted (ac)	95.16	12.14	0.74	113.45	138.90
	Species Habitat Availabl		15,358.72	1,274.76	2,292.11	15,822.86	4,698.61
	Percent Impacted O	ver Available	0.6196%	0.9524%	0.0324%	0.7170%	2.9561%
	Percent Available,	Not Impacted	99.3804%	99.0476%	99.9676%	99.2830%	97.0439%

ac = acres

Feature Description	Habitat Status	Area (Ac)	Cistothorus platensis	Dolichonyx oryzivorus	Falco sparverius	Haliaeetus leucocephalus	Lanius ludovicianus
Comptony	Impacted	0.00	NA NA	NA	0.00	NA NA	0.00
Cemetery	Available	3.47	NA	NA	3.47	NA	3.47
Posidontial High Density or Multiple Duralling	Impacted	0.00	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Available	0.00	NA	NA	NA	NA	NA
Desidential Dural Cingle Unit	Impacted	0.00	NA	NA	0.00	NA	NA
Residential, Rural, Single Unit	Available	9.13	NA	NA	9.13	NA	NA
Posidontial Single Unit Law Density	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Available	0.20	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Onit, Medium Density	Available	1.05	NA	NA	NA	NA	NA
Commercial/Services	Impacted	0.00	NA	NA	NA	NA	NA
Commercial/Services	Available	0.42	NA	NA	NA	NA	NA
Military Installations	Impacted	31.35	NA	NA	NA	NA	NA
ivilitary installations	Available	3,132.47	NA	NA	NA	NA	NA
Transportation/Communication/Utilities	Impacted	1.36	NA	NA	NA	NA	NA
Transportation/communication/ othities	Available	220.36	NA	NA	NA	NA	NA
Upland Right-of-Way Undeveloped	Impacted	0.00	NA	NA	0.00	NA	0.00
Opiana Right-or-way Ondeveloped	Available	81.09	NA	NA	81.09	NA	81.09
Stormwater Basin	Impacted	0.00	NA	NA	NA	NA	NA
Storiliwater Basiii	Available	33.27	NA	NA	NA	NA	NA
Other Urban or Built-up Land	Impacted	16.61	NA	NA	NA	NA	NA
Other Orban or Built-up Land	Available	1,250.32	NA	NA	NA	NA	NA
Managed Wetland in Maintained Lawn Greenspace	Impacted	0.00	NA	NA	0.00	NA	NA
IManaged Wetiand in Maintained Lawn Greenspace	Available	4.17	NA	NA	4.17	NA	NA
Athletic Fields (Schools)	Impacted	0.00	NA	NA	0.00	NA	NA
Athletic Fields (Schools)	Available	2.51	NA	NA	2.51	NA	NA
Recreational Land	Impacted	114.81	NA	NA	114.81	NA	NA
Recreational Land	Available	429.81	NA	NA	429.81	NA	NA
Agricultural Wetlands (Modified)	Impacted	92.84	NA	92.84	92.84	NA	NA
Agricultural Wetlands (Woulfied)	Available	93.97	NA	93.97	93.97	NA	NA
Cropland and Pastureland	Impacted	0.00	NA	0.00	0.00	NA	0.00
cropiana ana rastaretana	Available	14.17	NA	14.17	14.17	NA	14.17
Deciduous Forest (10-50% Crown Closure)	Impacted	2.46	NA	NA	NA	NA	NA
Deciduous Forest (10-50% Crown Closure)	Available	297.32	NA	NA	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted	39.15	NA	NA	NA	NA	NA
Deciduous Forest (>30% crown closure)	Available	1,522.35	NA	NA	NA	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
connectous Forest (10-30/0 crown closure)	Available	3,476.78	NA	NA	NA	NA	NA
Coniferous Forest (>50% Crown Closure)	Impacted	7.15	NA	NA	NA	NA	NA
connerous rorest (250% crown closure)	Available	6,216.02	NA	NA	NA	NA	NA
Plantation	Impacted	10.95	NA	NA	10.95	NA	NA
	Available	341.48	NA	NA	341.48	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
Times : orest (* 55% comicious with 10 56% crown closure)	Available	209.19	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted	34.29	NA	NA	NA	NA	NA
Times : orest (* 55% confictions with * 56% crown closure)	Available	1,819.31	NA	NA	NA	NA	NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
Times . Orest (* 55% Sectional With 10 55% Crown Closure)	Available	100.33	NA	NA	NA	NA	NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted	10.93	NA	NA	NA	NA	NA
Times Torest (2 55/5 Sectional With 250/6 Crown Closure)	Available	865.23	NA	NA	NA	NA	NA
Old Field (< 25% Brush Covered)	Impacted	1.19	NA	1.19	1.19	NA	1.19
ola Ficia (* 25% brasil coverea)	Available	830.21	NA	830.21	830.21	NA	830.21
Deciduous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	0.00
Decidadous Di usii/ Siii ubiallu	Available	78.60	NA	NA	NA	NA	78.60
Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	0.00
Connectous brushly shirubianu	Available	679.92	NA	NA	NA	NA	679.92
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	5.77	NA	NA	NA	NA	5.77
ivinca Decidadas Colificidas Diasily Siliabidia	Available	734.89	NA	NA	NA	NA	734.89

Feature Description	Habitat Status	Area (Ac)	Cistothorus platensis	Dolichonyx oryzivorus	Falco sparverius	Haliaeetus leucocephalus	Lanius ludovicianus
Streams and Canals	Impacted	6.73	NA	NA	NA	6.73	NA
Streams and Canais	Available	452.34	NA	NA	NA	452.34	NA
Natural Lakes	Impacted	0.00	NA	NA	NA	0.00	NA
Ivaturai Lakes	Available	4.54	NA	NA	NA	4.54	NA
Artificial Lakes	Impacted	7.17	NA	NA	NA	7.17	NA
Al tilitial takes	Available	396.28	NA	NA	NA	396.28	NA
Deciduous Wooded Wetlands	Impacted	9.80	NA	NA	NA	9.80	NA
Deciduous vvooded vvetialius	Available	2,334.37	NA	NA	NA	2,334.37	NA
Coniferous Wooded Wetlands	Impacted	0.00	NA	NA	NA	0.00	NA
Connerous wooded wetlands	Available	1,607.30	NA	NA	NA	1,607.30	NA
Atlantic White Cedar Wetlands	Impacted	0.00	NA	NA	NA	0.00	NA
Additic Wille Cedai Wedalids	Available	298.97	NA	NA	NA	298.97	NA
Deciduous Scrub/Shrub Wetlands	Impacted	0.00	0.00	NA	NA	0.00	0.00
Deciduous Scrub/Siliub Wetlands	Available	371.69	371.69	NA	NA	371.69	371.69
Coniferous Scrub/Shrub Wetlands	Impacted	0.00	0.00	NA	NA	0.00	0.00
Connerous Scrub/Siliub Wetlands	Available	292.71	292.71	NA	NA	292.71	292.71
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA	NA	0.00	0.00
ivilked Scrub/Siliub Wetlands (Deciduous Dolli.)	Available	635.15	NA	NA	NA	635.15	635.15
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	NA	0.00	0.00
ivilked Scrub/Siliub Wetlands (Confferous Doffi.)	Available	101.91	NA	NA	NA	101.91	101.91
Herbaceous Wetlands	Impacted	0.74	0.74	NA	NA	0.74	NA
nerbaceous wedanus	Available	843.63	843.63	NA	NA	843.63	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted	12.13	NA	NA	NA	12.13	NA
ivilxed wooded wetlands (Deciduous Doin.)	Available	523.83	NA	NA	NA	523.83	NA
Mixed Wooded Watlands (Canifornus Dom )	Impacted	0.00	NA	NA	NA	0.00	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Available	635.50	NA	NA	NA	635.50	NA
Disturbed Wetlands (Modified)	Impacted	0.00	0.00	NA	NA	0.00	0.00
Disturbed Wetlands (Woulfied)	Available	47.03	47.03	NA	NA	47.03	47.03
Transitional Areas	Impacted	27.51	NA	NA	27.51	NA	NA
Transitional Areas	Available	166.61	NA	NA	166.61	NA	NA
Undifferentiated Barren Lands	Impacted	0.00	NA	0.00	0.00	NA	NA
onumerentiateu barren Lanus	Available	4.34	NA	4.34	4.34	NA	NA
Phragmites Dominant Interior Wetlands	Impacted	0.00	NA	NA	NA	0.00	NA
rinaginites Dominant interior Wetlands	Available	2.67	NA	NA	NA	2.67	NA
	Species Habitat Im	npacted (ac)	0.74	94.03	247.31	36.58	6.97
	Species Habitat Available	Overall (ac)	1,555.05	942.69	1,980.96	8,547.90	3,870.82
	Percent Impacted Ove	er Available	0.0478%	9.9744%	12.4842%	0.4280%	0.1800%
	Percent Available, No	ot Impacted	99.9522%	90.0256%	87.5158%	99.5720%	99.8200%

ac = acres

Feature Description	Habitat Status	Area (Ac)	Melanerpes erythrocephalus	Nyctanassa violaceus	Nycticorax nycticorax	Pandion haliaetus	Passerculus sandwichensis
Compton	Impacted	0.00	NA	NA	NA	NA	0.00
Cemetery	Available	3.47	NA	NA	NA	NA	3.47
Decidential High Density on Multiple Develling	Impacted	0.00	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Available	0.00	NA	NA	NA	NA	NA
Desidential Dural Cinale Unit	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Available	9.13	NA	NA	NA	NA	NA
Posidontial Single Unit Law Density	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Available	0.20	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Offit, Mediani Density	Available	1.05	NA	NA	NA	NA	NA
Commercial/Services	Impacted	0.00	NA	NA	NA	NA	NA
Commercial/Services	Available	0.42	NA	NA	NA	NA	NA
Military Installations	Impacted	31.35	NA	NA	NA	NA	NA
ivilital y ilistaliations	Available	3,132.47	NA	NA	NA	NA	NA
Transportation/Communication/Utilities	Impacted	1.36	NA	NA	NA	NA	NA
Transportation/ communication/ offittles	Available	220.36	NA	NA	NA	NA	NA
Upland Right-of-Way Undeveloped	Impacted	0.00	NA	NA	NA	NA	0.00
Opiana Right-or-way Ondeveloped	Available	81.09	NA	NA	NA	NA	81.09
Stormwater Basin	Impacted	0.00	NA	NA	NA	NA	NA
Stormwater basin	Available	33.27	NA	NA	NA	NA	NA
Other Urban or Puilt up Land	Impacted	16.61	NA	NA	NA	NA	NA
Other Urban or Built-up Land	Available	1,250.32	NA	NA	NA	NA	NA
Managad Watland in Maintained Lawn Greenspase	Impacted	0.00	NA	NA	NA	NA	NA
Managed Wetland in Maintained Lawn Greenspace	Available	4.17	NA	NA	NA	NA	NA
Athletic Fields (Schools)	Impacted	0.00	NA	NA	NA	NA	NA
Athletic Fields (Schools)	Available	2.51	NA	NA	NA	NA	NA
Recreational Land	Impacted	114.81	NA	NA	NA	NA	114.81
Recreational Land	Available	429.81	NA	NA	NA	NA	429.81
Agricultural Wetlands (Modified)	Impacted	92.84	NA	NA	NA	NA	NA
Agricultural Wetlands (Wodilled)	Available	93.97	NA	NA	NA	NA	NA
Cropland and Pastureland	Impacted	0.00	NA	NA	NA	NA	NA
Cropiana and Pastureiana	Available	14.17	NA	NA	NA	NA	NA
Deciduous Forest (10-50% Crown Closure)	Impacted	2.46	NA	NA	NA	NA	NA
Deciduous Forest (10-30% Crown Closure)	Available	297.32	NA	NA	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted	39.15	39.15	NA	NA	NA	NA
Deciduous Polest (>30% Clowii Closule)	Available	1,522.35	1,522.35	NA	NA	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
Connerous Forest (10-30% Crown Closure)	Available	3,476.78	NA	NA	NA	NA	NA
Coniferous Forest (>50% Crown Closure)	Impacted	7.15	7.15	NA	NA	NA	NA
Confidences Forest (>50% Crown Closure)	Available	6,216.02	6,216.02	NA	NA	NA	NA
Plantation	Impacted	10.95	NA	NA	NA	NA	NA
Tantation	Available	341.48	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
Thinked Forest (> 50% Connerous with 10-50% Crown Closure)	Available	209.19	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted	34.29	34.29	NA	NA	NA	NA
Trinca i orest (> 50% Connerous with >50% Crown Closure)	Available	1,819.31	1,819.31	NA	NA	NA	NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted	0.00	NA	NA	NA	NA	NA
mined i orest (>30% Deciduous with 10-30% Crown Closure)	Available	100.33	NA	NA	NA	NA	NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted	10.93	10.93	NA	NA	NA	NA
Militar i orest (>30% Deciduous with >30% Crown Closure)	Available	865.23	865.23	NA	NA	NA	NA
Old Field (< 25% Brush Covered)	Impacted	1.19	NA	NA	NA	NA	1.19
Old Field (< 23% brush Covered)	Available	830.21	NA	NA	NA	NA	830.21
Deciduous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
Deciduous Diusii/ Siii ubidiiu	Available	78.60	NA	NA	NA	NA	NA
Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
Connectous brushing sin ubianu	Available	679.92	NA	NA	NA	NA	NA
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	5.77	NA	NA	NA	NA	NA
mixed Deciduous, Connerous Drushi, Shrubianu	Available	734.89	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Melanerpes erythrocephalus	Nyctanassa violaceus	Nycticorax nycticorax	Pandion haliaetus	Passerculus sandwichensis
Streams and Canals	Impacted	6.73	NA	6.73	6.73	NA	NA
Streams and Canais	Available	452.34	NA	452.34	452.34	NA	NA
Natural Lakes	Impacted	0.00	NA	0.00	0.00	0.00	NA
Natural Lakes	Available	4.54	NA	4.54	4.54	4.54	NA
Artificial Lakes	Impacted	7.17	NA	7.17	7.17	7.17	NA
Al tilitial takes	Available	396.28	NA	396.28	396.28	396.28	NA
Deciduous Wooded Wetlands	Impacted	9.80	9.80	9.80	9.80	NA	NA
Deciduous aaooden aaetiailus	Available	2,334.37	2,334.37	2,334.37	2,334.37	NA	NA
Coniferous Wooded Wetlands	Impacted	0.00	0.00	0.00	0.00	NA	NA
Connerous Wooded Wetlands	Available	1,607.30	1,607.30	1,607.30	1,607.30	NA	NA
Atlantic White Cedar Wetlands	Impacted	0.00	0.00	NA	0.00	NA	NA
Atlantic White Cedar Wetlands	Available	298.97	298.97	NA	298.97	NA	NA
Deciduous Scrub/Shrub Wetlands	Impacted	0.00	NA	0.00	0.00	NA	NA
Deciduous Scrub/Siliub Wedailus	Available	371.69	NA	371.69	371.69	NA	NA
Coniferous Scrub/Shrub Wetlands	Impacted	0.00	NA	0.00	0.00	NA	NA
Connerous Scrub/Siliub Wetlands	Available	292.71	NA	292.71	292.71	NA	NA
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	NA	0.00	0.00	NA	NA
ivilked Scrub/Siliub Wetlands (Deciduous Dolli.)	Available	635.15	NA	635.15	635.15	NA	NA
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	NA	0.00	0.00	NA	NA
wettands (connerous bonn.)	Available	101.91	NA	101.91	101.91	NA	NA
Herbaceous Wetlands	Impacted	0.74	NA	0.74	0.74	NA	NA
Tierbaceous wecianus	Available	843.63	NA	843.63	843.63	NA	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted	12.13	12.13	12.13	12.13	NA	NA
wooded wetlands (Deciduous Dom.)	Available	523.83	523.83	523.83	523.83	NA	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted	0.00	0.00	0.00	0.00	NA	NA
wooded wetlands (connerous bonn.)	Available	635.50	635.50	635.50	635.50	NA	NA
Disturbed Wetlands (Modified)	Impacted	0.00	NA	0.00	0.00	NA	NA
Disturbed Wetlands (Woulfied)	Available	47.03	NA	47.03	47.03	NA	NA
Transitional Areas	Impacted	27.51	NA	NA	NA	NA	NA
Transitional Areas	Available	166.61	NA	NA	NA	NA	NA
Undifferentiated Barren Lands	Impacted	0.00	NA	NA	NA	NA	0.00
Chairerentiated Barren Lands	Available	4.34	NA	NA	NA	NA	4.34
Phragmites Dominant Interior Wetlands	Impacted	0.00	NA	0.00	0.00	NA	NA
i magnites bolimant interior wetlands	Available	2.67	NA	2.67	2.67	NA	NA
	Species Habitat	Impacted (ac)	113.45	36.58	36.58	7.17	116.00
	Species Habitat Availabl	e Overall (ac)	15,822.86	8,248.93	8,547.90	400.82	1,348.92
	Percent Impacted C	ver Available	0.7170%	0.4435%	0.4280%	1.7888%	8.5994%
	Percent Available,	Not Impacted	99.2830%	99.5565%	99.5720%	98.2112%	91.4006%

ac = acres

Impacted	NA N
Available   3.47   NA	NA N
Available   0.00	NA N
Available   0.00	NA N
Available   9.13   NA   NA   NA   NA   NA   NA   NA   N	NA N
Available   9.13   NA	NA N
Available   0.20	NA N
Available   0.20	NA N
Available   1.05	NA N
Available   1.05	NA N
Available   0.42	NA N
Available   0.42   NA	NA N
NA	NA
Available   3,132.47	NA NA NA NA NA NA NA NA NA
Available   220.36	NA NA NA NA NA
Available   220.36	NA NA NA NA
Upland Right-of-Way Undeveloped         Available         81.09         NA         NA         NA         NA         81.09           Stormwater Basin         Impacted         0.00         NA	NA NA NA NA
Upland Right-of-Way Undeveloped         Available         81.09         NA         NA         NA         NA         81.09           Stormwater Basin         Impacted         0.00         NA	NA NA NA NA
Stormwater Basin         Impacted Available         0.00 33.27         NA NA NA         NA NA NA         NA NA         NA NA         NA NA         NA NA         NA NA         NA NA         NA NA         NA NA NA         NA NA NA         NA NA NA         NA NA NA         NA NA NA         NA NA NA         NA NA NA         NA NA	NA NA NA
Available 33.27 NA	NA NA
Other Urban or Built-up Land  Impacted 16.61 NA NA NA NA NA NA NA  Available 1,250.32 NA NA NA NA NA NA  Managed Wetland in Maintained Lawn Greenspace  Impacted 0.00 NA NA NA NA NA NA	NA
Available 1,250.32 NA	
Managed Wetland in Maintained Lawn Greenspace Impacted 0.00 NA NA NA NA NA NA	
IManaged Wetland in Maintained Lawn Greenspace	NA
7401001C 1.27 147	NA
Impacted 0.00 NA NA NA NA	NA NA
Athletic Fields (Schools)  Available	NA
Impacted 114.81 NA NA NA NA NA	NA NA
Recreational Land  Available 429.81 NA NA NA NA NA	NA NA
Available 425.61 NA NA NA NA NA NA	NA NA
Agricultural Wetlands (Modified)	
	NA NA
Cropland and PasturelandImpacted0.00NA0.00NANAAvailable14.17NA14.17NANA	NA NA
IDeciduous Forest (10-50% Crown Closure)	NA
Available 297.32 NA	NA NA
Deciduous Forest (>50% Crown Closure)  Impacted 39.15  NA  NA  NA  1.523.25	NA
Available 1,522.35 NA NA 1,522.35 NA	NA
Coniferous Forest (10-50% Crown Closure)  Impacted 0.00 NA	NA
Available 3,476.78 NA NA NA NA NA	NA
Coniferous Forest (>50% Crown Closure)  Impacted 7.15 NA NA 7.15 NA O. 1.15 N	NA
Available 6,216.02 NA NA 6,216.02 NA	NA NA
Plantation NA 10.95 NA	NA
Available 341.48 NA 341.48 NA NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)  Impacted  0.00  NA  NA  NA  NA  NA  NA  NA  NA  NA	NA
Available 209.19 NA NA NA NA	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)  Impacted 34.29  NA  NA  NA  34.29  NA  NA  NA  NA  NA  NA  NA  NA  NA  N	NA
Available 1,819.31 NA NA 1,819.31 NA	NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)  Impacted  0.00  NA  NA  NA  NA  NA  NA  NA  NA  NA	NA
Available 100.33 NA NA NA NA	NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)  Impacted 10.93  NA  NA  NA  10.93  NA  NA  NA  NA  NA  NA  NA  NA  NA  N	NA
Available 865.23 NA NA 865.23 NA	NA
Old Field (< 25% Brush Covered)  Impacted 1.19  NA 1.19  NA 1.19  NA 1.19	NA
Available 830.21 NA 830.21 NA 830.21	NA
Deciduous Brush/Shrubland Impacted 0.00 NA	NA
Available 78.60 NA NA NA NA NA	NA
Coniferous Brush/Shrubland NA NA NA NA NA NA NA	NA
Available 679.92 NA NA NA NA NA	NA
Mixed Deciduous (Coniferous Brush (Shrubland NA NA NA NA NA	NA
Mixed Deciduous/Coniferous Brush/Shrubland  Available 734.89 NA NA NA NA NA	NA

Feature Description	Habitat Status	Area (Ac)	Podilymbus podiceps	Pooecetes gramineus	Strix varia	Atrytone arogos arogos	Bolaria selene myrina
Streams and Canals	Impacted	6.73	NA	NA	NA	NA	NA
Streams and Canais	Available	452.34	NA	NA	NA	NA	NA
Natural Lakes	Impacted	0.00	0.00	NA	NA	NA	NA
ivaturar Lakes	Available	4.54	4.54	NA	NA	NA	NA
Artificial Lakes	Impacted	7.17	7.17	NA	NA	NA	NA
Al tilicial Lakes	Available	396.28	396.28	NA	NA	NA	NA
Deciduous Wooded Wetlands	Impacted	9.80	9.80	NA	NA	NA	NA
Deciduous Wooded Wetlands	Available	2,334.37	2,334.37	NA	NA	NA	NA
Coniferous Wooded Wetlands	Impacted	0.00	0.00	NA	NA	NA	NA
Connerous wooded wetlands	Available	1,607.30	1,607.30	NA	NA	NA	NA
Atlantic White Cedar Wetlands	Impacted	0.00	0.00	NA	NA	NA	NA
Atlantic White Cedar Wetlands	Available	298.97	298.97	NA	NA	NA	NA
Deciduous Scrub/Shrub Wetlands	Impacted	0.00	0.00	NA	NA	NA	0.00
Deciduous Scrub/Siliub Wetlands	Available	371.69	371.69	NA	NA	NA	371.69
Coniferous Scrub/Shrub Wetlands	Impacted	0.00		NA	NA	NA	0.00
	Available	292.71		NA	NA	NA	292.71
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	0.00	NA	NA	NA	0.00
	Available	635.15	635.15	NA	NA	NA	635.15
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	0.00	NA	NA	NA	0.00
wettands (connerous bonn.)	Available	101.91	101.91	NA	NA	NA	101.91
Herbaceous Wetlands	Impacted	0.74	0.74	NA	NA	NA	0.74
nerbaceous wetianus	Available	843.63	843.63	NA	NA	NA	843.63
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted	12.13	NA	NA	12.13	NA	NA
wooded wettands (Deciduous Dom.)	Available	523.83	NA	NA	523.83	NA	NA
Miyad Waadad Watlanda (Coniferens Dam )	Impacted	0.00	NA	NA	0.00	NA	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Available	635.50	NA	NA	635.50	NA	NA
Disturbed Wetlands (Modified)	Impacted	0.00	NA	NA	NA	NA	NA
Disturbed Wetlands (Modified)	Available	47.03	NA	NA	NA	NA	NA
Transitional Areas	Impacted	27.51	NA	NA	NA	NA	NA
Transitional Areas	Available	166.61	NA	NA	NA	NA	NA
Undifferentiated Barren Lands	Impacted	0.00	NA	0.00	NA	NA	NA
Onumerentiateu barren Lanus	Available	4.34	NA	4.34	NA	NA	NA
Phragmitas Dominant Interior Matlands	Impacted	0.00	0.00	NA	NA	NA	NA
Phragmites Dominant Interior Wetlands	Available	2.67	2.67	NA	NA	NA	NA
	Species Habitat Ir	mpacted (ac)	17.71	12.14	103.65	1.19	0.74
	Species Habitat Available		6,596.50	1,190.20	11,582.23	911.30	2,245.08
	Percent Impacted Ov	ver Available	0.2685%	1.0201%	0.8949%	0.1303%	0.0331%
	Percent Available, N	lot Impacted	99.7315%	98.9799%	99.1051%	99.8697%	99.9669%

ac = acres

Feature Description	Habitat Status	Area (Ac)	Callophrys irus	Lynx rufus	Clemmys insculpta	Clemmys muhlenbergii	Crotalus horridus
Comotory	Impacted	0.00	NA	NA	NA	NA	NA
Cemetery	Available	3.47	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Impacted	0.00	NA	NA	NA	NA	NA
Residential, High Density of Multiple Dwelling	Available	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Impacted	0.00	NA	NA	NA	NA	NA
nesidential, Rural, Single Offic	Available	9.13	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Offic, Low Density	Available	0.20	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Impacted	0.00	NA	NA	NA	NA	NA
Residential, Single Offit, Wedidin Density	Available	1.05	NA	NA	NA	NA	NA
Commercial/Services	Impacted	0.00	NA	NA	NA	NA	NA
Commercially Services	Available	0.42	NA	NA	NA	NA	NA
Military Installations	Impacted	31.35	NA	NA	NA	NA	NA
ivilitary installations	Available	3,132.47	NA	NA	NA	NA	NA
Transportation/Communication/Utilities	Impacted	1.36	NA	NA	NA	NA	NA
Transportation, communication, othices	Available	220.36	NA	NA	NA	NA	NA
Upland Right-of-Way Undeveloped	Impacted	0.00	0.00	NA	NA	NA	NA
Opiana Right-or-way Onacyclopea	Available	81.09	81.09	NA	NA	NA	NA
Stormwater Basin	Impacted	0.00	NA	NA	NA	NA	NA
Stormwater basin	Available	33.27	NA	NA	NA	NA	NA
Other Urban or Built-up Land	Impacted	16.61	NA	NA	NA	NA	NA
Other Orban or Built-up Land	Available	1,250.32	NA	NA	NA	NA	NA
Managed Wetland in Maintained Lawn Greenspace	Impacted	0.00	NA	NA	NA	NA	NA
ianageu wenanu in iviaintaineu Lawn Greenspace	Available	4.17	NA	NA	NA	NA	NA
Athletic Fields (Schools)	Impacted	0.00	NA	NA	NA	NA	NA
Admetic Ficial (Schools)	Available	2.51	NA	NA	NA	NA	NA
Recreational Land	Impacted	114.81	NA	NA	NA	NA	NA
Necreational Earla	Available	429.81	NA	NA	NA	NA	NA
gricultural Wetlands (Modified)	Impacted	92.84	NA	NA	NA	NA	NA
Agricultural Wellands (Modifica)	Available	93.97	NA	NA	NA	NA	NA
Cropland and Pastureland	Impacted	0.00	NA	0.00	NA	NA	NA
or opining and it assure tails	Available	14.17	NA	14.17	NA	NA	NA
Deciduous Forest (10-50% Crown Closure)	Impacted	2.46	NA	NA	NA	NA	NA
200,000,000,000,000,000,000,000,000,000	Available	297.32	NA	NA	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted	39.15	NA	NA	NA	NA	NA
200144040101001(7 2070 0101111 0100410)	Available	1,522.35	NA	NA	NA	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted	0.00	NA	0.00	NA	NA	0.00
Connections to test (10 30% crown crossing)	Available	3,476.78	NA	3,476.78	NA	NA	3,476.78
Coniferous Forest (>50% Crown Closure)	Impacted	7.15	NA	7.15	NA	NA	7.15
Connections to rest (* 50% crown crossine)	Available	6,216.02	NA	6,216.02	NA	NA	6,216.02
Plantation	Impacted	10.95	NA	NA	NA	NA	NA
	Available	341.48	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted	0.00	NA	0.00	NA	NA	0.00
, 11 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1 1	Available	209.19	NA	209.19	NA	NA	209.19
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted	34.29	NA	34.29	NA	NA	34.29
	Available	1,819.31	NA	1,819.31	NA	NA	1,819.31
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted	0.00	NA	0.00	NA	NA	0.00
2. 2.2. ( 2.2. 2.2. 2.2. 2.2. 2.2. 2.2.	Available	100.33	NA	100.33	NA	NA	100.33
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted	10.93	NA	10.93	NA	NA	10.93
The state of the s	Available	865.23	NA	865.23	NA	NA	865.23
Old Field (< 25% Brush Covered)	Impacted	1.19	1.19	NA	NA	NA	NA
	Available	830.21	830.21	NA	NA	NA	NA
Deciduous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
	Available	78.60	NA	NA	NA	NA	NA
Coniferous Brush/Shrubland	Impacted	0.00	NA	NA	NA	NA	NA
	Available	679.92	NA	NA	NA	NA	NA
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	5.77	NA	NA	NA	NA	NA
	Available	734.89	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Callophrys irus	Lynx rufus	Clemmys insculpta	Clemmys muhlenbergii	Crotalus horridus
Streams and Canals	Impacted	6.73	NA	NA	6.73	NA	NA
Streams and Canais	Available	452.34	NA	NA	452.34	NA	NA
Natural Lakes	Impacted	0.00	NA	NA	0.00	NA	NA
ivaturar Lakes	Available	4.54	NA	NA	4.54	NA	NA
Artificial Lakes	Impacted	7.17	NA	NA	NA	NA	NA
Artificial Lakes	Available	396.28	NA	NA	NA	NA	NA
Deciduous Wooded Wetlands	Impacted	9.80	NA	NA	9.80	NA	NA
Decidaous Wooded Wetlands	Available	2,334.37	NA	NA	2,334.37	NA	NA
Coniferous Wooded Wetlands	Impacted	0.00	NA	NA	0.00	NA	0.00
Connerous wooded wetlands	Available	1,607.30	NA	NA	1,607.30	NA	1,607.30
Atlantic White Cedar Wetlands	Impacted	0.00	NA	NA	NA	NA	0.00
Atlantic White Cedai Wetlands	Available	298.97	NA	NA	NA	NA	298.97
Deciduous Scrub/Shrub Wetlands	Impacted	0.00	NA	NA	0.00	0.00	NA
Deciduous Scrub/Siliub Wetlands	Available	371.69	NA	NA	371.69	371.69	NA
Coniferous Scrub/Shrub Wetlands	Impacted	0.00	NA	NA	0.00	0.00	NA
	Available	292.71	NA	NA	292.71	292.71	NA
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA	0.00	0.00	NA
	Available	635.15	NA	NA	635.15	635.15	NA
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	0.00	0.00	NA
wettands (Connerous Doni.)	Available	101.91	NA	NA	101.91	101.91	NA
Herbaceous Wetlands	Impacted	0.74	NA	NA	0.74	0.74	NA
nerbaceous wetianus	Available	843.63	NA	NA	843.63	843.63	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted	12.13	NA	NA	12.13	NA	NA
ivilxed wooded wetlands (Decidaous Doin.)	Available	523.83	NA	NA	523.83	NA	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted	0.00	NA	NA	0.00	NA	0.00
ivilxed wooded wetlands (confierous bonn.)	Available	635.50	NA	NA	635.50	NA	635.50
Disturbed Wetlands (Modified)	Impacted	0.00	NA	NA	NA	NA	NA
Disturbed Wetlands (Modified)	Available	47.03	NA	NA	NA	NA	NA
Transitional Areas	Impacted	27.51	NA	NA	NA	NA	NA
Transitional Areas	Available	166.61	NA	NA	NA	NA	NA
Undifferentiated Barren Lands	Impacted	0.00	NA	NA	NA	NA	NA
ondinerentiated barren Lands	Available	4.34	NA	NA	NA	NA	NA
Phragmites Dominant Interior Wetlands	Impacted	0.00	NA	NA	0.00	0.00	NA
rmagnites Dominant interior Wetlands	Available	2.67	NA	NA	2.67	2.67	NA
	Species Habitat Ir	mpacted (ac)	1.19	52.36	29.41	0.74	52.36
	Species Habitat Available	Overall (ac)	911.30	12,701.03	7,805.62	2,247.75	15,228.62
	Percent Impacted Ov	er Available	0.1303%	0.4123%	0.3768%	0.0330%	0.3438%
	Percent Available, N	lot Impacted	99.8697%	99.5877%	99.6232%	99.9670%	99.6562%

ac = acres

Feature Description	Habitat Status	Area (Ac)	Elaphe guttata	Pituophis m melanoleucus
	Impacted	0.00	, NA	NA
Cemetery	Available	3.47	NA	NA
Residential, High Density or Multiple Dwelling	Impacted	0.00	NA	NA
nesidential, riigh bensity of Multiple bwening	Available	0.00	NA	NA
Residential, Rural, Single Unit	Impacted	0.00	NA	NA
, , ,	Available	9.13	NA	NA NA
Residential, Single Unit, Low Density	Impacted Available	0.00 0.20	NA NA	NA NA
	Impacted	0.20	NA NA	NA NA
Residential, Single Unit, Medium Density	Available	1.05	NA NA	NA NA
	Impacted	0.00	NA	NA NA
Commercial/Services	Available	0.42	NA	NA
Military Installations	Impacted	31.35	NA	NA
wintary instanations	Available	3,132.47	NA	NA
Transportation/Communication/Utilities	Impacted	1.36	NA	NA
	Available	220.36	NA	NA 
Upland Right-of-Way Undeveloped	Impacted	0.00	NA NA	NA NA
	Available Impacted	81.09 0.00	NA NA	NA NA
Stormwater Basin	Available	33.27	NA NA	NA NA
	Impacted	16.61	NA	NA NA
Other Urban or Built-up Land	Available	1,250.32	NA	NA
Managad Watland in Maintained Laws Greensnass	Impacted	0.00	NA	NA
Managed Wetland in Maintained Lawn Greenspace	Available	4.17	NA	NA
Athletic Fields (Schools)	Impacted	0.00	NA	NA
	Available	2.51	NA	NA
Recreational Land	Impacted	114.81	NA	NA NA
	Available	429.81 92.84	NA NA	NA NA
Agricultural Wetlands (Modified)	Impacted Available	93.97	NA NA	NA NA
	Impacted	0.00	NA NA	NA NA
Cropland and Pastureland	Available	14.17	NA	NA NA
Desidueus Ferrest (10 F00) Creum Cleaums)	Impacted	2.46	NA	2.46
Deciduous Forest (10-50% Crown Closure)	Available	297.32	NA	297.32
Deciduous Forest (>50% Crown Closure)	Impacted	39.15	NA	NA
Decidadas Forest (7 30% drown drosare)	Available	1,522.35	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted	0.00	NA	0.00
	Available	3,476.78	NA 7.15	3,476.78
Coniferous Forest (>50% Crown Closure)	Impacted Available	7.15 6,216.02	7.15 6,216.02	NA NA
	Impacted	10.95	NA	NA NA
Plantation	Available	341.48	NA	NA NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted	0.00	NA	0.00
white Porest (> 50% Connerous with 10-50% Crown Closure)	Available	209.19	NA	209.19
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted	34.29	34.29	NA
	Available	1,819.31	1,819.31	NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted	0.00	NA NA	0.00
	Available Impacted	100.33 10.93	NA NA	100.33 NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Available	865.23	NA NA	NA NA
	Impacted	1.19	NA NA	1.19
Old Field (< 25% Brush Covered)	Available	830.21	NA NA	830.21
Dociduous Brush/Shruhland	Impacted	0.00	NA	NA
Deciduous Brush/Shrubland	Available	78.60	NA	NA
Coniferous Brush/Shrubland	Impacted	0.00	NA	NA
	Available	679.92	NA	NA
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted	5.77	NA NA	NA NA
	Available	734.89	NA	NA

Feature Description	Habitat Status	Area (Ac)	Elaphe guttata	Pituophis m melanoleucus
Streams and Canals	Impacted	6.73	NA	NA
Streams and Canais	Available	452.34	NA	NA
Noticed Lakes	Impacted	0.00	NA	NA
Natural Lakes	Available	4.54	NA	NA
Artificial Lakes	Impacted	7.17	NA	NA
AI UIICIdi Lakes	Available	396.28	NA	NA
Deciduous Wooded Wetlands	Impacted	9.80	NA	NA
Deciduous wooded wettailus	Available	2,334.37	NA	NA
Coniferous Wooded Wetlands	Impacted	0.00	0.00	NA
Connerous wooded wetlands	Available	1,607.30	1,607.30	NA
Atlantic White Cedar Wetlands	Impacted	0.00	0.00	NA
Atlantic White Cedar Wetlands	Available	298.97	298.97	NA
Desidueus Comul /Chaula Metlands	Impacted	0.00	NA	NA
Deciduous Scrub/Shrub Wetlands	Available	371.69	NA	NA
Coniference County (Church Westlands	Impacted	0.00	0.00	NA
Coniferous Scrub/Shrub Wetlands	Available	292.71	292.71	NA
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted	0.00	NA	NA
ivilxed scrub/shrub Wetlands (Deciduous Dom.)	Available	635.15	NA	NA
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted	0.00	0.00	NA
weitands (conferous Dom.)	Available	101.91	101.91	NA
Hauba aa aya Maklanda	Impacted	0.74	NA	NA
Herbaceous Wetlands	Available	843.63	NA	NA
Missed Meeded Meklanda (Desidueus Dess)	Impacted	12.13	NA	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Available	523.83	NA	NA
Baiwad Mandad Maklanda (Canifarawa Dana)	Impacted	0.00	0.00	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Available	635.50	635.50	NA
Distructed Mathemate (Mandified)	Impacted	0.00	NA	NA
Disturbed Wetlands (Modified)	Available	47.03	NA	NA
Transitional Areas	Impacted	27.51	NA	NA
ITalisitional Areas	Available	166.61	NA	NA
Undifferentiated Barren Lands	Impacted	0.00	0.00	0.00
Ondinerentiated Barren Lands	Available	4.34	4.34	4.34
Physical Dominant Interior Mattends	Impacted	0.00	NA	NA
Phragmites Dominant Interior Wetlands	Available	2.67	NA	NA
	Species Habitat	Impacted (ac)	41.44	3.65
	Species Habitat Availab	le Overall (ac)	10,976.05	4,918.17
	Percent Impacted	Over Available	0.3775%	0.0742%
	Percent Available,	Not Impacted	99.6225%	99.9258%

ac = acres

Table B-3. Spatial Exposure Potential of Protected Species at Lakehurst of Joint Base McGuire-Dix-Lakehurst

Feature Description	Habitat Status	Area (Ac)	Clemmy guttata	Pseudemys rubriventris (Juvenile)	Pseudemys rubriventris (Adult)	Hyla andersonii	Hyla chrysocelis
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Cemetery	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	20.73	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
,	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Commercial/Services	Impacted (AFFF Area)	0.00	NA	NA NA	NA	NA	NA
•	Available	2.62	NA	NA NA	NA	NA	NA
	Impacted (AFFF Drainage)	72.52	NA NA	NA NA	NA	NA	NA NA
Military Installations	Impacted (AFFF Area)	22.07	NA	NA NA	NA NA	NA	NA NA
initally installations	Available	903.70	NA	NA NA	NA	NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Transportation/Communication/Utilities	Impacted (AFFF Area)	0.00	NA	NA NA	NA	NA	NA NA
	Available	2.43	NA NA	NA NA	NA	NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA	NA NA
Upland Right-of-Way Undeveloped	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA	NA NA
, ,	Available	3.42	NA	NA NA	NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA	NA NA	NA	NA	0.00
Stormwater Basin	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	0.00
	Available	23.49	NA	NA NA	NA	NA	23.49
	Impacted (AFFF Drainage)	67.20	NA NA	NA NA	NA NA	NA NA	NA
Other Urban or Built-up Land	Impacted (AFFF Area)	8.22	NA	NA NA	NA	NA	NA
other orban or bank up tand	Available	1,729.81	NA	NA NA	NA NA	NA	NA NA
	Impacted (AFFF Drainage)	0.04	NA NA	0.04	NA NA	NA NA	0.04
Managed Wetland in Maintained Lawn Greenspace	Impacted (AFFF Area)	0.00	NA NA	0.00	NA NA	NA	0.00
Wallaged Westalia in Wallitanied Edwir Greenspace	Available	10.38	NA NA	10.38	NA NA	NA	10.38
	Impacted (AFFF Drainage)	0.00	NA NA	NA	NA NA	NA NA	NA
Athletic Fields (Schools)	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Attrictic Ficius (Schools)	Available	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Recreational Land	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Recreational Land	Available	82.29	NA NA	NA NA	NA NA		NA NA
	Impacted (AFFF Drainage)		NA NA	0.00	NA NA	NA NA	0.00
Agricultural Wetlands (Modified)		0.00					
Agricultulal Wetialius (Wiouillea)	Impacted (AFFF Area)	0.00	NA NA	0.00	NA NA	NA NA	0.00
	Available	0.00	NA NA	0.00	NA NA	NA NA	0.00
Cropland and Dastivialized	Impacted (AFFF Drainage)	0.00	NA NA	0.00	NA NA	NA NA	0.00
Cropland and Pastureland	Impacted (AFFF Area)	0.00	NA NA	0.00	NA NA	NA NA	0.00
	Available	0.00	NA NA	0.00	NA NA	NA NA	0.00
Desidueus Ferent /10 FOO/ Cuerum Clearum	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA	NA
Deciduous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA	NA	NA
	Available	0.73	NA NA	NA NA	NA NA	NA NA	NA
Double of Francisco	Impacted (AFFF Drainage)	0.01	NA	NA NA	NA 	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	NA	NA	NA	NA	NA
	Available	146.32	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	184.15	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Clemmy guttata	Pseudemys rubriventris (Juvenile)	Pseudemys rubriventris (Adult)	Hyla andersonii	Hyla chrysocelis
	Impacted (AFFF Drainage)	45.67	NA	NA	NA	NA	NA
Coniferous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	NA	NA	NA	NA	NA
·	Available	1,381.20	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Plantation	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	104.26	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA	NA	NA NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)		0.00	NA	NA	NA NA	NA	NA NA
	Available	10.91	NA NA	NA NA	NA NA	NA	NA NA
	Impacted (AFFF Drainage)	15.12	NA NA	NA NA	NA NA	NA NA	NA NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted (AFFF Area)	0.01	NA NA	NA NA	NA NA	NA NA	NA NA
ivilized Forest (> 30% confirerous with >30% crown closure)	Available	904.40	NA NA	NA NA	NA NA	NA NA	NA NA
					NA NA		
Missed Ferrest (> F00/ Periduana with 10 F00/ Creases Cleanes)	Impacted (AFFF Drainage)	0.00	NA NA	NA NA		NA	NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	· '	0.00	NA	NA NA	NA	NA	NA
	Available	14.62	NA	NA 	NA	NA	NA
Adjust Farrest (SPROV Death as the PROV C. C.	Impacted (AFFF Drainage)	1.32	NA NA	NA NA	NA	NA	NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted (AFFF Area)	0.06	NA	NA 	NA	NA	NA
	Available	375.12	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.30	NA	NA	NA	NA	NA
Old Field (< 25% Brush Covered)	Impacted (AFFF Area)	0.06	NA	NA	NA	NA	NA
	Available	124.81	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Deciduous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	1.15	NA	NA	NA	NA	NA
Coniferous Brush/Shrubland	Impacted (AFFF Drainage)	5.44	NA	NA	NA	NA	NA
	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	200.94	NA	NA	NA	NA	NA
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted (AFFF Drainage)	3.82	NA	NA	NA	NA	NA
	Impacted (AFFF Area)	0.10	NA	NA	NA	NA	NA
	Available	80.91	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	4.45	4.45	NA	4.45	NA	NA
Streams and Canals	Impacted (AFFF Area)	0.00	0.00	NA	0.00	NA	NA
	Available	67.79	67.79	NA	67.79	NA	NA
	Impacted (AFFF Drainage)	0.00	0.00	0.00	0.00	NA	NA
Natural Lakes	Impacted (AFFF Area)	0.00	0.00	0.00	0.00	NA	NA
	Available	0.95	0.95	0.95	0.95	NA	NA NA
	Impacted (AFFF Drainage)	0.07	0.07	0.07	0.07	NA	NA NA
Artificial Lakes	Impacted (AFFF Area)	0.00	0.00	0.00	0.00	NA	NA NA
The tribut Edites	Available	91.55	91.55	91.55	91.55	NA	NA NA
	Impacted (AFFF Drainage)	2.58	2.58	2.58	NA	NA NA	2.58
Deciduous Wooded Wetlands	Impacted (AFFF Area)	0.00	0.00	0.00	NA NA	NA NA	0.00
beciadous vvooded vvetidiids	Available	63.59	63.59	63.59	NA NA	NA NA	63.59
	Impacted (AFFF Drainage)	2.06	2.06	2.06	NA NA	2.06	2.06
Coniferous Wooded Wetlands		0.00	0.00	0.00	NA NA	0.00	0.00
Connerous vvooueu vvetialius	Impacted (AFFF Area)						
	Available	289.14	289.14	289.14	NA NA	289.14	289.14
Atlantia White Cadau Malada	Impacted (AFFF Drainage)	0.00	0.00	0.00	NA NA	0.00	0.00
Atlantic White Cedar Wetlands	Impacted (AFFF Area)	0.00	0.00	0.00	NA NA	0.00	0.00
	Available	109.14	109.14	109.14	NA NA	109.14	109.14
	Impacted (AFFF Drainage)	3.34	3.34	3.34	NA	3.34	3.34
Deciduous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	0.00	0.00	NA	0.00	0.00
	Available	83.03	83.03	83.03	NA	83.03	83.03
		0.00	0.00	0.00	NA	0.00	0.00
	Impacted (AFFF Drainage)	0.00					
Coniferous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	0.00	0.00	NA	0.00	0.00
Coniferous Scrub/Shrub Wetlands					NA NA	0.00 51.98	
Coniferous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	0.00	0.00			0.00
Coniferous Scrub/Shrub Wetlands  Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted (AFFF Area) Available	0.00 51.98	0.00 51.98	0.00 51.98	NA	51.98	0.00 51.98

Feature Description	Habitat Status	Area (Ac)	Clemmy guttata	Pseudemys rubriventris (Juvenile)	Pseudemys rubriventris (Adult)	Hyla andersonii	Hyla chrysocelis
	Impacted (AFFF Drainage)	0.00	0.00	0.00	NA	0.00	0.00
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	0.00	0.00	NA	0.00	0.00
	Available	63.10	63.10	63.10	NA	63.10	63.10
	Impacted (AFFF Drainage)	0.00	0.00	0.00	NA	0.00	0.00
Herbaceous Wetlands	Impacted (AFFF Area)	0.00	0.00	0.00	NA	0.00	0.00
	Available	21.06	21.06	21.06	NA	21.06	21.06
	Impacted (AFFF Drainage)	1.44	1.44	1.44	NA	1.44	1.44
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	0.00	0.00	NA	0.00	0.00
	Available	47.03	47.03	47.03	NA	47.03	47.03
	Impacted (AFFF Drainage)	0.00	0.00	0.00	NA	0.00	0.00
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	0.00	0.00	NA	0.00	0.00
·	Available	78.97	78.97	78.97	NA	78.97	78.97
Disturbed Wetlands (Modified)	Impacted (AFFF Drainage)	0.51	NA	0.51	NA	NA	0.51
	Impacted (AFFF Area)	0.28	NA	0.28	NA	NA	0.28
	Available	32.13	NA	32.13	NA	NA	32.13
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Transitional Areas	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	14.34	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Undifferentiated Barren Lands	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	34.78	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.38	NA	NA	NA	NA	NA
Phragmites Dominant Interior Wetlands	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	7.23	NA	NA	NA	NA	NA
	Species Habitat Impacted, AFI	FF Drainage (ac)	14.58	10.68	4.52	7.49	10.61
	Species Habitat Impacted	l, AFFF Area (ac)	0.00	0.28	0.00	0.00	0.28
	Species Habitat Availa	able Overall (ac)	1,027.47	1,002.20	160.29	803.60	933.19
	Percent Impacted Over Available	, AFFF Drainage	1.4190%	1.0660%	2.8169%	0.9317%	1.1374%
	Percent Impacted Over Avail	lable, AFFF Area	0.0000%	0.0279%	0.0000%	0.0000%	0.0300%
	Percent Available	e, Not Impacted	98.5810%	98.9061%	97.1831%	99.0683%	98.8326%

ac = acres NA = not applicable

Feature Description	Habitat Status	Area (Ac)	Pseudotriton montanus	Accipiter gentilis	Ammodramus henslowii	Ammodramus savannarum	Asio flammeus
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	0.00
Cemetery	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	0.00
•	Available	0.00	NA	NA	NA	NA	0.00
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	20.73	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
, , ,	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
, , , , , , , , , , , , , , , , , , ,	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA	NA NA	NA NA	NA
Residential, Single Unit, Medium Density	Impacted (AFFF Area)	0.00	NA	NA NA	NA	NA	NA
income in any one give only income in a constraint	Available	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Commercial/Services	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	2.62	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	72.52	NA NA	NA NA	NA NA	NA NA	NA NA
Military Installations	Impacted (AFFF Area)	22.07	NA NA	NA NA	NA NA	NA NA	NA NA
The state of the s	Available	903.70	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Transportation/Communication/Utilities	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Transportation, communication, othities	Available	2.43	NA NA	NA NA	NA NA	NA NA	NA NA
Upland Right-of-Way Undeveloped	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	0.00	0.00	NA NA
	Impacted (AFFF Area)	0.00	NA NA	NA NA	0.00	0.00	NA NA
	Available	3.42	NA NA	NA NA	3.42	3.42	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA	NA	NA NA
Stormwater Basin	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Stormwater basin	Available	23.49	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	67.20	NA NA	NA NA	NA NA	NA NA	67.20
Other Urban or Built-up Land	Impacted (AFFF Area)	8.22	NA NA	NA NA	NA NA	NA NA	8.22
Other Orban or Built-up Land	Available	1,729.81	NA NA	NA NA	NA NA	NA NA	1,729.81
	Impacted (AFFF Drainage)	0.04	NA NA	NA NA	NA NA	NA NA	1,729.81 NA
Managed Wetland in Maintained Lawn Greenspace		0.00	NA NA	NA NA	NA NA	NA NA	NA NA
I wanaged wetland in Manitained Lawn Greenspace	Impacted (AFFF Area) Available	10.38					
			NA NA	NA NA	NA NA	NA NA	NA NA
Athlatic Fields (Schools)	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Athletic Fields (Schools)	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Pocreational Land	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Recreational Land	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	82.29	NA NA	NA NA	NA NA	NA NA	NA NA
Agricultural Motlands (Madified)	Impacted (AFFF Drainage)	0.00	NA	NA NA	NA NA	NA NA	NA NA
Agricultural Wetlands (Modified)	Impacted (AFFF Area)	0.00	NA	NA NA	NA NA	NA NA	NA NA
	Available	0.00	NA NA	NA NA	NA NA	NA NA	NA 0.00
Cupuland and Dastriusland	Impacted (AFFF Drainage)	0.00	NA	NA NA	NA NA	NA NA	0.00
Cropland and Pastureland	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	0.00
	Available	0.00	NA	NA NA	NA NA	NA NA	0.00
D. 11 F /40 TOX 6	Impacted (AFFF Drainage)	0.00	NA	NA NA	NA NA	NA NA	NA
Deciduous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	NA 	NA 	NA 	NA
	Available	0.73	NA	NA	NA NA	NA	NA
	Impacted (AFFF Drainage)	0.01	NA	0.01	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	NA	0.01	NA	NA	NA
	Available	146.32	NA	146.32	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	184.15	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Pseudotriton montanus	Accipiter gentilis	Ammodramus henslowii	Ammodramus savannarum	Asio flammeus
	Impacted (AFFF Drainage)	45.67	NA	45.67	NA	NA	NA
Coniferous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	NA	0.01	NA	NA	NA
	Available	1,381.20	NA	1,381.20	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Plantation	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	104.26	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
· 1	Available	10.91	NA	NA	NA	NA NA	NA
	Impacted (AFFF Drainage)	15.12	NA	15.12	NA	NA	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted (AFFF Area)	0.01	NA	0.01	NA	NA	NA
, , , , , , , , , , , , , , , , , , , ,	Available	904.40	NA NA	904.40	NA	NA NA	NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA	NA NA	NA NA	NA NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	NA NA	NA NA	NA NA	NA
Thinked Forest (F50/0 Beefladous With 10 50/0 crown crosure)	Available	14.62	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	1.32	NA NA	1.32	NA NA	NA NA	NA NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted (AFFF Area)	0.06	NA NA	0.06	NA NA	NA NA	NA NA
	Available	375.12	NA NA	375.12	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.30	NA NA	NA	0.30	0.30	0.30
Old Field (< 25% Brush Covered)	Impacted (AFFF Area)	0.06	NA NA	NA NA	0.06	0.06	0.06
old Field (* 25% Brush covered)	Available	124.81	NA NA	NA NA	124.81	124.81	124.81
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA	NA	NA
Deciduous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Deciduous Brushy Shrubland	Available	1.15	NA NA	NA NA	NA NA	NA NA	NA NA
Coniferous Brush/Shrubland	Impacted (AFFF Drainage)	5.44	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	200.94	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	3.82	NA NA	NA NA	NA NA	NA NA	NA NA
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted (AFFF Area)	0.10	NA NA	NA NA	NA NA	NA NA	NA NA
Mixed Deciduous/Connerous Brushi/Sinabiand	Available				NA NA		
	Impacted (AFFF Drainage)	80.91 4.45	NA NA	NA NA	NA NA	NA NA	NA NA
Streams and Canals	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Streams and Canais	Available	67.79					
	Impacted (AFFF Drainage)	0.00	NA 0.00	NA NA	NA NA	NA NA	NA NA
Natural Lakes	, , ,			NA NA	NA NA	NA NA	NA NA
ivaturar Lakes	Impacted (AFFF Area)	0.00	0.00	NA NA	NA NA	NA NA	NA NA
	Available	0.95	0.95	NA NA	NA NA	NA NA	NA NA
Artificial Lakes	Impacted (AFFF Drainage)	0.07	0.07	NA NA	NA NA	NA NA	NA NA
Al Ulludi Lakes	Impacted (AFFF Area)	0.00	0.00	NA NA	NA NA	NA NA	NA NA
	Available	91.55	91.55	NA 2.59	NA NA	NA NA	NA NA
Deciduous Wooded Wetlands	Impacted (AFFF Drainage)	2.58	NA NA	2.58	NA NA	NA NA	NA NA
Deciduous vyooded vyelidiius	Impacted (AFFF Area) Available	0.00 63.59	NA NA	0.00 63.59	NA NA	NA NA	NA NA
<del> </del>			NA NA	2.06	NA NA	NA NA	NA NA
Coniferous Wooded Wetlands	Impacted (AFFF Drainage)	2.06	NA NA			NA NA	NA NA
Connerous wooden wettalius	Impacted (AFFF Area)	0.00	NA NA	0.00	NA NA	NA NA	NA NA
	Available	289.14	NA 0.00	289.14	NA NA	NA NA	NA NA
Atlantic White Coder Westlands	Impacted (AFFF Drainage)	0.00	0.00	0.00	NA NA	NA NA	NA NA
Atlantic White Cedar Wetlands	Impacted (AFFF Area)	0.00	0.00	0.00	NA NA	NA NA	NA NA
	Available	109.14	109.14	109.14	NA NA	NA NA	NA NA
Desidueus Comph/Chmph Madalarada	Impacted (AFFF Drainage)	3.34	NA NA	NA NA	NA NA	NA NA	NA NA
Deciduous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	83.03	NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA 	NA 	NA
Coniferous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	NA	NA	NA 	NA 	NA
	Available	51.98	NA	NA	NA NA	NA	NA
	Impacted (AFFF Drainage)	0.65	NA	NA	NA	NA	NA
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	60.14	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Pseudotriton montanus	Accipiter gentilis	Ammodramus henslowii	Ammodramus savannarum	Asio flammeus
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	63.10	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	0.00	NA	0.00	NA	NA
Herbaceous Wetlands	Impacted (AFFF Area)	0.00	0.00	NA	0.00	NA	NA
	Available	21.06	21.06	NA	21.06	NA	NA
	Impacted (AFFF Drainage)	1.44	NA	1.44	NA	NA	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	NA	0.00	NA	NA	NA
	Available	47.03	NA	47.03	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	0.00	NA	NA	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	NA	0.00	NA	NA	NA
	Available	78.97	NA	78.97	NA	NA	NA
Disturbed Wetlands (Modified)	Impacted (AFFF Drainage)	0.51	0.51	NA	NA	NA	NA
	Impacted (AFFF Area)	0.28	0.28	NA	NA	NA	NA
	Available	32.13	32.13	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Transitional Areas	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	14.34	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	0.00	0.00	NA
Undifferentiated Barren Lands	Impacted (AFFF Area)	0.00	NA	NA	0.00	0.00	NA
	Available	34.78	NA	NA	34.78	34.78	NA
	Impacted (AFFF Drainage)	0.38	NA	NA	NA	NA	0.38
Phragmites Dominant Interior Wetlands	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	0.00
	Available	7.23	NA	NA	NA	NA	7.23
	Species Habitat Impacted, AFF	F Drainage (ac)	0.57	68.21	0.30	0.30	67.88
	Species Habitat Impacted,	AFFF Area (ac)	0.28	0.09	0.06	0.06	8.28
	Species Habitat Availal	ble Overall (ac)	254.83	3,394.91	184.07	163.01	1,861.84
	Percent Impacted Over Available,	AFFF Drainage	0.2253%	2.0091%	0.1646%	0.1859%	3.6459%
	Percent Impacted Over Availa	able, AFFF Area	0.1099%	0.0027%	0.0327%	0.0369%	0.4446%
	Percent Available	, Not Impacted	99.6648%	97.9882%	99.8027%	99.7772%	95.9094%

ac = acres NA = not applicable

Feature Description	Habitat Status	Area (Ac)	Asio otus	Batramia longicauda	Botaurus lentiginosos	Buteo lineatus	Circus cyaneus
	Impacted (AFFF Drainage)	0.00	0.00	0.00	NA	NA	0.00
Cemetery	Impacted (AFFF Area)	0.00	0.00	0.00	NA	NA	0.00
	Available	0.00	0.00	0.00	NA	NA	0.00
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	20.73	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
,	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Impacted (AFFF Area)	0.00	NA	NA	NA	NA NA	NA
,	Available	0.00	NA NA	NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA NA	NA NA
Commercial/Services	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	2.62	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	72.52	NA NA	NA NA	NA NA	NA NA	NA NA
Military Installations	Impacted (AFFF Area)	22.07	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	903.70	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Transportation/Communication/Utilities	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Transportation, communication, othices	Available	2.43	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	0.00	NA NA	NA NA	0.00
Upland Right-of-Way Undeveloped	Impacted (AFFF Area)	0.00	NA NA	0.00	NA NA	NA NA	0.00
opiana night-oi-way ondeveloped	Available	3.42	NA NA	3.42	NA NA	NA NA	3.42
	Impacted (AFFF Drainage)	0.00	NA NA	NA	NA NA	NA NA	NA
Stormwater Basin	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Stormwater basin	Available	23.49	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	67.20	NA NA	NA NA	NA NA	NA NA	67.20
Other Urban or Built-up Land	Impacted (AFFF Area)	8.22	NA NA	NA NA	NA NA		8.22
Other Orban or Bunt-up Land	Available		NA NA			NA NA	
		1,729.81		NA NA	NA NA	NA NA	1,729.81
Managed Wetland in Maintained Laws Creenance	Impacted (AFFF Drainage)	0.04	NA NA	NA NA	NA NA	NA NA	0.04
Managed Wetland in Maintained Lawn Greenspace	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	0.00
	Available	10.38	NA NA	NA NA	NA NA	NA NA	10.38
Ashlesia Fielda (Calessala)	Impacted (AFFF Drainage)	0.00	NA	NA NA	NA	NA NA	0.00
Athletic Fields (Schools)	Impacted (AFFF Area)	0.00	NA	NA	NA	NA NA	0.00
	Available	0.00	NA NA	NA NA	NA NA	NA NA	0.00
December of Land	Impacted (AFFF Drainage)	0.00	NA NA	NA	NA	NA NA	NA NA
Recreational Land	Impacted (AFFF Area)	0.00	NA NA	NA	NA	NA NA	NA NA
	Available	82.29	NA NA	NA	NA NA	NA NA	NA 0.00
A	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA 	0.00
Agricultural Wetlands (Modified)	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	0.00
	Available	0.00	NA	NA	NA	NA	0.00
	Impacted (AFFF Drainage)	0.00	0.00	0.00	NA	NA	0.00
Cropland and Pastureland	Impacted (AFFF Area)	0.00	0.00	0.00	NA	NA	0.00
	Available	0.00	0.00	0.00	NA	NA	0.00
	Impacted (AFFF Drainage)	0.00	0.00	NA	NA	NA	NA
Deciduous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	0.00	NA	NA	NA	NA
	Available	0.73	0.73	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.01	0.01	NA	NA	0.01	NA
Deciduous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	0.01	NA	NA	0.01	NA
	Available	146.32	146.32	NA	NA	146.32	NA
	Impacted (AFFF Drainage)	0.00	0.00	NA	NA	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	0.00	NA	NA	NA	NA
	Available	184.15	184.15	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Asio otus	Batramia longicauda	Botaurus lentiginosos	Buteo lineatus	Circus cyaneus
	Impacted (AFFF Drainage)	45.67	45.67	NA	NA	45.67	NA
Coniferous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	0.01	NA	NA	0.01	NA
·	Available	1,381.20	1,381.20	NA	NA	1,381.20	NA
	Impacted (AFFF Drainage)	0.00	NA	0.00	NA	NA	NA
Plantation	Impacted (AFFF Area)	0.00	NA	0.00	NA	NA	NA
	Available	104.26	NA	104.26	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	0.00	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)		0.00	0.00	NA	NA	NA	NA
, ,	Available	10.91	10.91	NA	NA	NA	NA
	Impacted (AFFF Drainage)	15.12	15.12	NA	NA	15.12	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted (AFFF Area)	0.01	0.01	NA	NA	0.01	NA
,	Available	904.40	904.40	NA	NA	904.40	NA
	Impacted (AFFF Drainage)	0.00	0.00	NA NA	NA NA	NA NA	NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted (AFFF Area)	0.00	0.00	NA NA	NA NA	NA NA	NA
,	Available	14.62	14.62	NA	NA NA	NA	NA
	Impacted (AFFF Drainage)	1.32	1.32	NA NA	NA NA	1.32	NA NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted (AFFF Area)	0.06	0.06	NA NA	NA NA	0.06	NA NA
	Available	375.12	375.12	NA NA	NA NA	375.12	NA NA
	Impacted (AFFF Drainage)	0.30	0.30	0.30	NA NA	NA	0.30
Old Field (< 25% Brush Covered)	Impacted (AFFF Area)	0.06	0.06	0.06	NA NA	NA NA	0.06
one read ( - 15/5 Stabil Covered)	Available	124.81	124.81	124.81	NA NA	NA NA	124.81
	Impacted (AFFF Drainage)	0.00	NA	NA	NA NA	NA NA	NA
Deciduous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Decidous brushy shrubland	Available	1.15	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	5.44	NA NA		NA NA		
Coniferous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Connerous brushy sili ubianu	Available	200.94	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	3.82	NA NA	NA NA	NA NA		NA NA
Mixed Deciduous/Coniferous Brush/Shrubland		3.82 0.10	NA NA	NA NA	NA NA	NA NA	NA NA
ivilkeu Deciduous/Colilletous Brush/Shrubland	Impacted (AFFF Area)						
	Available	80.91	NA NA	NA NA	NA NA	NA NA	NA NA
Streams and Canals	Impacted (AFFF Drainage)	4.45	NA NA	NA NA	NA NA	NA NA	NA NA
Streams and Canais	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	67.79	NA NA	NA NA	NA NA	NA NA	NA NA
Natural Lakas	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Natural Lakes	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	0.95	NA NA	NA NA	NA NA	NA NA	NA NA
Autificial Labor	Impacted (AFFF Drainage)	0.07	NA	NA NA	NA NA	NA	NA
Artificial Lakes	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA
	Available	91.55	NA NA	NA NA	NA NA	NA 2.50	NA
B. H W J. IW d I	Impacted (AFFF Drainage)	2.58	NA	NA NA	NA NA	2.58	NA
Deciduous Wooded Wetlands	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	0.00	NA
	Available	63.59	NA NA	NA NA	NA NA	63.59	NA
o to a second of the second	Impacted (AFFF Drainage)	2.06	NA	NA NA	NA NA	2.06	NA
Coniferous Wooded Wetlands	Impacted (AFFF Area)	0.00	NA	NA 	NA	0.00	NA
	Available	289.14	NA	NA 	NA	289.14	NA
	Impacted (AFFF Drainage)	0.00	NA	NA 	NA 	0.00	NA
Atlantic White Cedar Wetlands	Impacted (AFFF Area)	0.00	NA	NA	NA	0.00	NA
	Available	109.14	NA	NA NA	NA	109.14	NA
	Impacted (AFFF Drainage)	3.34	NA	NA	3.34	NA	3.34
Deciduous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	0.00
	Available	83.03	NA	NA	83.03	NA	83.03
	Impacted (AFFF Drainage)	0.00	NA	NA	0.00	NA	0.00
Coniferous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	0.00
	Available	51.98	NA	NA	51.98	NA	51.98
	Impacted (AFFF Drainage)	0.65	NA	NA	0.65	NA	0.65
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	0.00
1	Available	60.14	NA	NA	60.14	NA	60.14

Feature Description	Habitat Status	Area (Ac)	Asio otus	Batramia longicauda	Botaurus lentiginosos	Buteo lineatus	Circus cyaneus
	Impacted (AFFF Drainage)	0.00	NA	NA	0.00	NA	0.00
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	0.00
	Available	63.10	NA	NA	63.10	NA	63.10
	Impacted (AFFF Drainage)	0.00	NA	NA	0.00	NA	0.00
Herbaceous Wetlands	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	0.00
	Available	21.06	NA	NA	NA     0.00     NA       NA     63.10     NA       NA     0.00     NA       NA     0.00     NA	21.06	
	Impacted (AFFF Drainage)	1.44	NA	NA	NA	1.44	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	NA	0.00	NA
	Available	47.03	63.10         NA         NA         63.10           0.00         NA         NA         0.00           0.00         NA         NA         0.00           21.06         NA         NA         NA           1.44         NA         NA         NA           0.00         NA         NA         NA           47.03         NA         NA         NA           0.00         NA         NA         NA           0.00         NA         NA         NA           0.00         NA         NA         NA           0.51         NA         NA         NA           0.28         NA         NA         NA           0.01         NA         NA         NA           0.02         NA         NA         NA           0.00         0.00         NA         NA           0.00         NA         NA         NA	47.03	NA		
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	0.00	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	NA	0.00	NA
	Available	78.97	NA	NA	NA	78.97	NA
	Impacted (AFFF Drainage)	0.51	NA	NA	0.51	NA	NA
Disturbed Wetlands (Modified)	Impacted (AFFF Area)	0.28	NA	NA	0.28	NA	NA
	Available	32.13	NA	NA	32.13	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	0.00
Transitional Areas	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	0.00
	Available	14.34	NA	NA	NA	NA	14.34
	Impacted (AFFF Drainage)	0.00	0.00	0.00	NA	NA	0.00
Undifferentiated Barren Lands	Impacted (AFFF Area)	0.00	0.00	0.00	NA	NA	0.00
	Available	34.78	34.78	34.78	NA	NA	34.78
	Impacted (AFFF Drainage)	0.38	NA	NA	NA	NA	0.38
Phragmites Dominant Interior Wetlands	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	0.00
	Available	7.23	NA	NA	NA	NA	7.23
	Species Habitat Impacted, AFFF	Drainage (ac)	62.43	0.30	4.49	68.21	71.91
	Species Habitat Impacted, A	AFFF Area (ac)	0.15	0.06	0.28	0.09	8.28
	Species Habitat Available	le Overall (ac)	3,177.04	267.27	311.44	3,394.91	2,204.07
	Percent Impacted Over Available, A	AFFF Drainage	1.9652%	0.1134%	1.4426%	2.0091%	3.2628%
	Percent Impacted Over Availab	ole, AFFF Area	0.0048%	0.0225%	0.0899%	0.0027%	0.3756%
	Percent Available, I	Not Impacted	98.0301%	99.8641%	98.4675%	97.9882%	96.3616%

Feature Description	Habitat Status	Area (Ac)	Cistothorus platensis	Dolichonyx oryzivorus	Falco sparverius	Haliaeetus leucocephalus	Lanius ludovicianus
	Impacted (AFFF Drainage)	0.00	NA	NA	0.00	NA	0.00
Cemetery	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	0.00
	Available	0.00	NA	NA	0.00	NA	0.00
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	20.73	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	0.00	NA	NA
Residential, Rural, Single Unit	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	NA
	Available	0.00	NA	NA	0.00	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA NA	NA	NA NA	NA
Commercial/Services	Impacted (AFFF Area)	0.00	NA	NA NA	NA	NA NA	NA
	Available	2.62	NA	NA NA	NA	NA NA	NA
	Impacted (AFFF Drainage)	72.52	NA NA	NA NA	NA NA	NA NA	NA
Military Installations	Impacted (AFFF Area)	22.07	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	903.70	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA
Transportation/Communication/Utilities	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	2.43	NA NA	NA NA	NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	0.00	NA NA	0.00
Upland Right-of-Way Undeveloped	Impacted (AFFF Area)	0.00	NA NA	NA NA	0.00	NA NA	0.00
opiana Right-or-way ondeveloped	Available	3.42	NA NA	NA NA	3.42	NA NA	3.42
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA	NA NA	NA
Stormwater Basin	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA	NA NA	NA
Stormwater basin	Available	23.49	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	67.20	NA NA	NA NA	NA NA	NA NA	NA NA
Other Urban or Built-up Land	Impacted (AFFF Area)	8.22	NA NA	NA NA	NA NA	NA NA	
Cities Orban of Built-up Land	Available		NA NA		NA NA		NA NA
<del> </del>		1,729.81		NA NA	0.04	NA NA	NA NA
Managed Wetland in Maintained Laws Creenance	Impacted (AFFF Drainage)	0.04	NA NA	NA NA		NA NA	NA
Managed Wetland in Maintained Lawn Greenspace	Impacted (AFFF Area)	0.00	NA NA	NA NA	0.00	NA NA	NA
	Available	10.38	NA NA	NA NA	10.38	NA NA	NA
Anthonic Biolds (Calcada)	Impacted (AFFF Drainage)	0.00	NA	NA NA	0.00	NA NA	NA
Athletic Fields (Schools)	Impacted (AFFF Area)	0.00	NA	NA NA	0.00	NA	NA
<del> </del>	Available	0.00	NA	NA	0.00	NA	NA
L	Impacted (AFFF Drainage)	0.00	NA	NA 	0.00	NA 	NA
Recreational Land	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	NA
	Available	82.29	NA	NA	82.29	NA	NA
A	Impacted (AFFF Drainage)	0.00	NA	0.00	0.00	NA 	NA
Agricultural Wetlands (Modified)	Impacted (AFFF Area)	0.00	NA	0.00	0.00	NA	NA
<u> </u>	Available	0.00	NA	0.00	0.00	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	0.00	0.00	NA	0.00
Cropland and Pastureland	Impacted (AFFF Area)	0.00	NA	0.00	0.00	NA	0.00
	Available	0.00	NA	0.00	0.00	NA	0.00
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Deciduous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
•	Available	0.73	NA	NA	NA	NA	NA
<u> </u>		0.01	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)			-		· •	
	Impacted (AFFF Drainage) Impacted (AFFF Area)	0.01	NA	NA	NA	NA	NA
			NA NA	NA NA	NA NA	NA NA	NA NA
Deciduous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01 146.32					
	Impacted (AFFF Area) Available	0.01	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Cistothorus platensis	Dolichonyx oryzivorus	Falco sparverius	Haliaeetus leucocephalus	Lanius ludovicianus
	Impacted (AFFF Drainage)	45.67	NA	NA	NA	NA	NA
Coniferous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	NA	NA	NA	NA	NA
	Available	1,381.20	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	0.00	NA	NA
Plantation	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	NA
	Available	104.26	NA	NA	104.26	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)		0.00	NA	NA	NA	NA	NA
	Available	10.91	NA	NA NA	NA	NA NA	NA
	Impacted (AFFF Drainage)	15.12	NA NA	NA NA	NA NA	NA NA	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted (AFFF Area)	0.01	NA	NA NA	NA NA	NA	NA
Trince Forest (* 50% connerous with * 50% crown closure)	Available	904.40	NA NA	NA NA	NA NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)		0.00	NA NA	NA NA	NA NA	NA NA	NA NA
lwiked Forest (>30% Decidaous with 10-30% Crown Closure)	Available	14.62					
			NA NA	NA NA	NA NA	NA NA	NA NA
Mixed Forest (NEOV Desiduous with NEOV Crown Classes)	Impacted (AFFF Drainage)	1.32	NA NA	NA NA	NA NA	NA NA	NA NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted (AFFF Area)	0.06	NA NA	NA NA	NA	NA NA	NA NA
	Available	375.12	NA NA	NA 0.20	NA 0.30	NA NA	NA 0.20
Old Field ( 4.35% Procts On 1991)	Impacted (AFFF Drainage)	0.30	NA	0.30	0.30	NA NA	0.30
Old Field (< 25% Brush Covered)	Impacted (AFFF Area)	0.06	NA	0.06	0.06	NA	0.06
	Available	124.81	NA	124.81	124.81	NA	124.81
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	0.00
Deciduous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	0.00
	Available	1.15	NA	NA	NA	NA	1.15
	Impacted (AFFF Drainage)	5.44	NA	NA	NA	NA	5.44
Coniferous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	0.00
	Available	200.94	NA	NA	NA	NA	200.94
	Impacted (AFFF Drainage)	3.82	NA	NA	NA	NA	3.82
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted (AFFF Area)	0.10	NA	NA	NA	NA	0.10
	Available	80.91	NA	NA	NA	NA	80.91
	Impacted (AFFF Drainage)	4.45	NA	NA	NA	4.45	NA
Streams and Canals	Impacted (AFFF Area)	0.00	NA	NA	NA	0.00	NA
	Available	67.79	NA	NA	NA	67.79	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	0.00	NA
Natural Lakes	Impacted (AFFF Area)	0.00	NA	NA	NA	0.00	NA
	Available	0.95	NA	NA	NA	0.95	NA
	Impacted (AFFF Drainage)	0.07	NA	NA	NA	0.07	NA
Artificial Lakes	Impacted (AFFF Area)	0.00	NA	NA	NA	0.00	NA
	Available	91.55	NA	NA	NA	91.55	NA
	Impacted (AFFF Drainage)	2.58	NA NA	NA NA	NA NA	2.58	NA
Deciduous Wooded Wetlands	Impacted (AFFF Area)	0.00	NA	NA NA	NA NA	0.00	NA
	Available	63.59	NA NA	NA NA	NA NA	63.59	NA NA
	Impacted (AFFF Drainage)	2.06	NA NA	NA NA	NA NA	2.06	NA NA
Coniferous Wooded Wetlands	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	0.00	NA NA
Commercial Westernance	Available	289.14	NA NA	NA NA	NA NA	289.14	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	0.00	NA NA
Atlantic White Cedar Wetlands	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	0.00	NA NA
Addition willing court wedialius	Available	109.14					
			NA 2.24	NA NA	NA NA	109.14	NA 2.24
Dociduous Corub/Shrub Matter de	Impacted (AFFF Drainage)	3.34	3.34	NA NA	NA NA	3.34	3.34
Deciduous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	0.00	NA NA	NA NA	0.00	0.00
	Available	83.03	83.03	NA NA	NA	83.03	83.03
a 16 a 16 a 1 a 1	Impacted (AFFF Drainage)	0.00	0.00	NA	NA	0.00	0.00
Coniferous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	0.00	NA	NA	0.00	0.00
	Available	51.98	51.98	NA	NA	51.98	51.98
	Impacted (AFFF Drainage)	0.65	NA	NA	NA	0.65	0.65
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	NA	0.00	0.00
	Available	60.14	NA	NA	NA	60.14	60.14

Feature Description	Habitat Status A	Area (Ac)	Cistothorus platensis	Dolichonyx oryzivorus	Falco sparverius	Haliaeetus leucocephalus	Lanius ludovicianus
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	0.00	0.00
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	NA	0.00	0.00
	Available	63.10	NA	NA	NA	63.10	63.10
	Impacted (AFFF Drainage)	0.00	0.00	NA	NA	0.00	NA
Herbaceous Wetlands	Impacted (AFFF Area)	0.00	0.00	NA	NA	0.00	NA
	Available	21.06	21.06	NA	NA	0.00 63.10 0.00 0.00 21.06 1.44 0.00 47.03 0.00 0.00 78.97 0.51 0.28 32.13 NA	NA
	Impacted (AFFF Drainage)	1.44	NA	NA	NA	1.44	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	NA	0.00	NA
	Available	47.03	NA	NA	NA	47.03	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	0.00	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	NA	0.00	NA
	Available	78.97	NA	NA	NA	78.97	NA
	Impacted (AFFF Drainage)	0.51	0.51	NA	NA	0.51	0.51
Disturbed Wetlands (Modified)	Impacted (AFFF Area)	0.28	0.28	NA	NA	0.28	0.28
	Available	32.13	32.13	NA	NA	32.13	32.13
	Impacted (AFFF Drainage)	0.00	NA	NA	0.00	NA	NA
Transitional Areas	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	NA
	Available	14.34	NA	NA	14.34	0.00 0.00 63.10 0.00 0.00 0.00 21.06 1.44 0.00 47.03 0.00 0.00 78.97 0.51 0.28 32.13 NA	NA
	Impacted (AFFF Drainage)	0.00	NA	0.00	0.00	NA	NA
Undifferentiated Barren Lands	Impacted (AFFF Area)	0.00	NA	0.00	0.00	NA	NA
	Available	34.78	NA	34.78	34.78	NA	NA
	Impacted (AFFF Drainage)	0.38	NA	NA	NA	0.38	NA
Phragmites Dominant Interior Wetlands	Impacted (AFFF Area)	0.00	NA	NA	NA	0.00	NA
	Available	7.23	NA	NA	NA	7.23	NA
	Species Habitat Impacted, AFFF D	Orainage (ac)	3.84	0.30	0.35	15.47	14.06
	Species Habitat Impacted, AF	_	0.28	0.06	0.06	0.28	0.44
	Species Habitat Available	Overall (ac)	188.20	159.59	374.27	1,066.83	701.59
	Percent Impacted Over Available, AF		2.0414%	0.1899%	0.0928%	1.4497%	2.0038%
	Percent Impacted Over Available	_	0.1488%	0.0377%	0.0161%	0.0263%	0.0627%
	Percent Available, N		97.8098%	99.7724%	99.8912%		97.9335%

Feature Description	Habitat Status	Area (Ac)	Melanerpes erythrocephalus	Nyctanassa violaceus	Nycticorax nycticorax	Pandion haliaetus	Passerculus sandwichensis
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	0.00
Cemetery	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	0.00
	Available	0.00	NA	NA	NA	NA	0.00
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	20.73	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Commercial/Services	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	2.62	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	72.52	NA	NA	NA	NA	NA
Military Installations	Impacted (AFFF Area)	22.07	NA	NA	NA	NA	NA
	Available	903.70	NA NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA	NA NA
Transportation/Communication/Utilities	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA	NA NA
,	Available	2.43	NA NA	NA NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA NA	NA NA	NA NA	0.00
Upland Right-of-Way Undeveloped	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	0.00
opiana inglico: iray onaciopea	Available	3.42	NA	NA NA	NA	NA	3.42
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA	NA NA
Stormwater Basin	Impacted (AFFF Area)	0.00	NA	NA NA	NA	NA	NA NA
	Available	23.49	NA	NA NA	NA	NA NA	NA NA
	Impacted (AFFF Drainage)	67.20	NA	NA NA	NA NA	NA	NA NA
Other Urban or Built-up Land	Impacted (AFFF Area)	8.22	NA	NA NA	NA	NA	NA NA
outer of bank up land	Available	1,729.81	NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.04	NA NA	NA NA	NA NA	NA	NA NA
Managed Wetland in Maintained Lawn Greenspace	Impacted (AFFF Area)	0.00	NA	NA NA	NA NA	NA NA	NA NA
inanagea ir etiana in mameanea zatin ereenspase	Available	10.38	NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Athletic Fields (Schools)	Impacted (AFFF Area)	0.00	NA	NA NA	NA NA	NA NA	NA NA
Administration (consolo)	Available	0.00	NA	NA NA	NA NA	NA NA	NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	0.00
Recreational Land	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	0.00
Nedicational Zana	Available	82.29	NA NA	NA NA	NA NA	NA NA	82.29
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA
Agricultural Wetlands (Modified)	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Agricultural Wetlanus (Mounteu)	Available	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Cropland and Pastureland	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
ι ο ριατία απά τ αθται σταπά	Available	0.00	NA NA	NA NA	NA NA		
						NA NA	NA NA
Desiduous Forest (10 F0% Crown Closure)	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Deciduous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	0.73	NA 0.04	NA NA	NA NA	NA NA	NA NA
Decidence Forcet & FOOCO C. C.	Impacted (AFFF Drainage)	0.01	0.01	NA NA	NA NA	NA	NA NA
Deciduous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	0.01	NA 	NA 	NA	NA
	Available	146.32	146.32	NA	NA	NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	184.15	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Melanerpes erythrocephalus	Nyctanassa violaceus	Nycticorax nycticorax	Pandion haliaetus	Passerculus sandwichensis
	Impacted (AFFF Drainage)	45.67	45.67	NA	NA	NA	NA
Coniferous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	0.01	NA	NA	NA	NA
	Available	1,381.20	1,381.20	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Plantation	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	104.26	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)		0.00	NA	NA NA	NA	NA	NA
,	Available	10.91	NA	NA NA	NA NA	NA	NA NA
	Impacted (AFFF Drainage)	15.12	15.12	NA NA	NA NA	NA NA	NA NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted (AFFF Area)	0.01	0.01	NA NA	NA NA	NA	NA NA
wince rolest (> 50% connerous with > 50% crown closure)	Available	904.40	904.40	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA	NA NA	NA NA	NA NA	NA NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)		0.00					
lwixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted (AFFF Area)		NA NA	NA NA	NA NA	NA NA	NA
	Available	14.62	NA 132	NA NA	NA NA	NA NA	NA NA
Balived Fewert (NFOO/ Deciderary with NFOO/ Comments	Impacted (AFFF Drainage)	1.32	1.32	NA NA	NA NA	NA NA	NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted (AFFF Area)	0.06	0.06	NA NA	NA NA	NA NA	NA
	Available	375.12	375.12	NA NA	NA NA	NA NA	NA 0.20
	Impacted (AFFF Drainage)	0.30	NA	NA	NA	NA	0.30
Old Field (< 25% Brush Covered)	Impacted (AFFF Area)	0.06	NA	NA	NA	NA	0.06
	Available	124.81	NA	NA	NA	NA	124.81
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Deciduous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	1.15	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	5.44	NA	NA	NA	NA	NA
Coniferous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	200.94	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	3.82	NA	NA	NA	NA	NA
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted (AFFF Area)	0.10	NA	NA	NA	NA	NA
	Available	80.91	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	4.45	NA	4.45	4.45	NA	NA
Streams and Canals	Impacted (AFFF Area)	0.00	NA	0.00	0.00	NA	NA
	Available	67.79	NA	67.79	67.79	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	0.00	0.00	0.00	NA
Natural Lakes	Impacted (AFFF Area)	0.00	NA	0.00	0.00	0.00	NA
	Available	0.95	NA	0.95	0.95	0.95	NA
	Impacted (AFFF Drainage)	0.07	NA NA	0.07	0.07	0.07	NA
Artificial Lakes	Impacted (AFFF Area)	0.00	NA NA	0.00	0.00	0.00	NA
	Available	91.55	NA	91.55	91.55	91.55	NA NA
	Impacted (AFFF Drainage)	2.58	2.58	2.58	2.58	NA NA	NA NA
Deciduous Wooded Wetlands	Impacted (AFFF Area)	0.00	0.00	0.00	0.00	NA	NA NA
beciauous vvooded vveilailus	Available	63.59	63.59	63.59	63.59	NA NA	NA NA
	Impacted (AFFF Drainage)	2.06	2.06	2.06	2.06	NA NA	NA NA
Coniferous Wooded Wetlands	Impacted (AFFF Area)	0.00	0.00	0.00	0.00	NA NA	NA NA
Connectous aacoucu aactiulius	Available	289.14	289.14	289.14	289.14	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	0.00	289.14 NA	0.00	NA NA	NA NA
Atlantic White Cedar Wetlands	Impacted (AFFF Area)	0.00	0.00	NA NA	0.00	NA NA	NA NA
Additio venite cedai vectianus	Available	109.14			109.14		
			109.14	NA 2.24		NA NA	NA NA
Deciduous Corub/Chrub Methor de	Impacted (AFFF Drainage)	3.34	NA NA	3.34	3.34	NA NA	NA NA
Deciduous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	NA NA	0.00	0.00	NA NA	NA
	Available	83.03	NA NA	83.03	83.03	NA NA	NA
	Impacted (AFFF Drainage)	0.00	NA 	0.00	0.00	NA	NA
Coniferous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	NA 	0.00	0.00	NA	NA
	Available	51.98	NA	51.98	51.98	NA	NA
L.,	Impacted (AFFF Drainage)	0.65	NA	0.65	0.65	NA	NA
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	NA	0.00	0.00	NA	NA
	Available	60.14	NA	60.14	60.14	NA	NA

Feature Description	Habitat Status	Area (Ac)	Melanerpes erythrocephalus	Nyctanassa violaceus	Nycticorax nycticorax	Pandion haliaetus	Passerculus sandwichensis
	Impacted (AFFF Drainage)	0.00	NA	0.00	0.00	NA	NA
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	NA	0.00	0.00	NA	NA
	Available	63.10	NA	63.10	63.10	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	0.00	0.00	NA	NA
Herbaceous Wetlands	Impacted (AFFF Area)	0.00	NA	0.00	0.00	NA	NA
	Available	21.06	NA	21.06	21.06	NA N	NA
	Impacted (AFFF Drainage)	1.44	1.44	1.44	1.44	NA	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	0.00	0.00	0.00	NA	NA
	Available	47.03	47.03	47.03	0.00         0.00         NA           63.10         63.10         NA           0.00         0.00         NA           0.00         0.00         NA           21.06         21.06         NA           1.44         1.44         NA           0.00         0.00         NA           47.03         47.03         NA           0.00         0.00         NA           0.00         0.00         NA           78.97         NA         NA           0.28         0.28         NA           32.13         NA         NA           NA         NA         NA	NA	
	Impacted (AFFF Drainage)	0.00	0.00	0.00	0.00	NA	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	0.00	0.00	0.00	NA	NA
	Available	78.97	78.97	78.97	78.97	NA	NA
	Impacted (AFFF Drainage)	0.51	NA	0.51	0.51	NA	NA
Disturbed Wetlands (Modified)	Impacted (AFFF Area)	0.28	NA	0.28	0.28	NA	NA
	Available	32.13	NA	32.13	32.13	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Transitional Areas	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	14.34	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	0.00
Undifferentiated Barren Lands	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	0.00
	Available	34.78	NA	NA	NA	NA	34.78
	Impacted (AFFF Drainage)	0.38	NA	0.38	0.38	NA	NA
Phragmites Dominant Interior Wetlands	Impacted (AFFF Area)	0.00	NA	0.00	0.00	NA	NA
	Available	7.23	NA	7.23	7.23	NA	NA
	Species Habitat Impacted, AFFF	Drainage (ac)	68.21	15.47	15.47	0.07	0.30
	Species Habitat Impacted, A	AFFF Area (ac)	0.09	0.28	0.28	0.00	0.06
	Species Habitat Availab	ole Overall (ac)	3,394.91	957.69	1,066.83	92.50	245.30
	Percent Impacted Over Available, A	AFFF Drainage	2.0091%	1.6150%	1.4497%	0.0744%	0.1235%
	Percent Impacted Over Availal	ble, AFFF Area	0.0027%	0.0292%	0.0263%	0.0000%	0.0245%
	Percent Available,	Not Impacted	97.9882%	98.3558%	98.5240%	99.9256%	99.8519%

Feature Description	Habitat Status	Area (Ac)	Podilymbus podiceps	Pooecetes gramineus	Strix varia	Atrytone arogos arogos	Bolaria selene myrina
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Cemetery	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	20.73	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
, , , ,	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Commercial/Services	Impacted (AFFF Area)	0.00	NA	NA	NA NA	NA NA	NA NA
•	Available	2.62	NA	NA NA	NA	NA NA	NA NA
	Impacted (AFFF Drainage)	72.52	NA NA	NA NA	NA NA	NA NA	NA NA
Military Installations	Impacted (AFFF Area)	22.07	NA NA	NA NA	NA NA	NA NA	NA NA
,	Available	903.70	NA NA	NA NA	NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Transportation/Communication/Utilities	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA	NA NA	NA NA
	Available	2.43	NA NA	NA NA	NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	0.00	NA NA
Upland Right-of-Way Undeveloped	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	0.00	NA NA
opidina highe of tray officercloped	Available	3.42	NA NA	NA NA	NA NA	3.42	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA	NA NA
Stormwater Basin	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA	NA NA	NA NA
otorimater basin	Available	23.49	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	67.20	NA NA	NA NA	NA NA	NA NA	NA NA
Other Urban or Built-up Land	Impacted (AFFF Area)	8.22	NA NA	NA NA	NA	NA NA	NA NA
other orban or bune-up taile	Available	1,729.81	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.04	NA NA	NA NA	NA NA	NA NA	NA NA
Managed Wetland in Maintained Lawn Greenspace	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
ividilaged wetland in Maintained Lawn Greenspace	Available	10.38	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Athletic Fields (Schools)	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Atmetic Fields (Schools)	Available	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Recreational Land	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
necieational Land	Available						
	Impacted (AFFF Drainage)	82.29 0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Agricultural Wetlands (Modified)				NA NA			
ngricultulai vvetialius (ivibuilleu)	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	0.00	NA NA	NA 0.00	NA NA	NA NA	NA NA
Cropland and Dartureland	Impacted (AFFF Drainage)	0.00	NA NA	0.00	NA NA		NA NA
Cropland and Pastureland	Impacted (AFFF Area)	0.00	NA NA	0.00	NA NA	NA NA	NA NA
	Available	0.00	NA NA	0.00	NA NA	NA NA	NA NA
Desidueus Ferrest (40 F00/ Commercial	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA
Deciduous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA NA	NA	NA NA	NA NA	NA
	Available	0.73	NA 	NA NA	NA	NA NA	NA
	Impacted (AFFF Drainage)	0.01	NA	NA	0.01	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	NA	NA	0.01	NA	NA
	Available	146.32	NA	NA	146.32	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Coniferous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	184.15	NA	NA	NA	NA	NA

Feature Description	Habitat Status	Area (Ac)	Podilymbus podiceps	Pooecetes gramineus	Strix varia	Atrytone arogos arogos	Bolaria selene myrina
	Impacted (AFFF Drainage)	45.67	NA	NA	45.67	NA	NA
Coniferous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	NA	NA	0.01	NA	NA
	Available	1,381.20	NA	NA	1,381.20	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	0.00	NA	NA	NA
Plantation	Impacted (AFFF Area)	0.00	NA	0.00	NA	NA	NA
	Available	104.26	NA	104.26	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	10.91	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	15.12	NA	NA	15.12	NA	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted (AFFF Area)	0.01	NA	NA	0.01	NA	NA
, ,	Available	904.40	NA	NA	904.40	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
, ,	Available	14.62	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	1.32	NA	NA	1.32	NA	NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted (AFFF Area)	0.06	NA	NA	0.06	NA NA	NA
,	Available	375.12	NA	NA	375.12	NA	NA
	Impacted (AFFF Drainage)	0.30	NA	0.30	NA	0.30	NA NA
Old Field (< 25% Brush Covered)	Impacted (AFFF Area)	0.06	NA	0.06	NA	0.06	NA NA
•	Available	124.81	NA	124.81	NA NA	124.81	NA NA
	Impacted (AFFF Drainage)	0.00	NA	NA NA	NA NA	NA NA	NA NA
Deciduous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA	NA	NA	NA NA	NA
	Available	1.15	NA	NA	NA	NA NA	NA
	Impacted (AFFF Drainage)	5.44	NA	NA NA	NA NA	NA	NA
Coniferous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA	NA NA	NA	NA	NA
	Available	200.94	NA	NA NA	NA	NA	NA NA
	Impacted (AFFF Drainage)	3.82	NA	NA NA	NA	NA NA	NA NA
ixed Deciduous/Coniferous Brush/Shrubland	Impacted (AFFF Area)	0.10	NA	NA	NA	NA NA	NA NA
	Available	80.91	NA	NA	NA	NA NA	NA
	Impacted (AFFF Drainage)	4.45	NA	NA	NA NA	NA NA	NA NA
Streams and Canals	Impacted (AFFF Area)	0.00	NA	NA	NA	NA NA	NA
	Available	67.79	NA	NA	NA NA	NA NA	NA
	Impacted (AFFF Drainage)	0.00	0.00	NA NA	NA NA	NA NA	NA NA
Natural Lakes	Impacted (AFFF Area)	0.00	0.00	NA	NA	NA	NA
	Available	0.95	0.95	NA NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.07	0.07	NA NA	NA NA	NA NA	NA
Artificial Lakes	Impacted (AFFF Area)	0.00	0.00	NA NA	NA NA	NA	NA NA
	Available	91.55	91.55	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	2.58	2.58	NA NA	NA NA	NA NA	NA NA
Deciduous Wooded Wetlands	Impacted (AFFF Area)	0.00	0.00	NA NA	NA NA	NA NA	NA NA
	Available	63.59	63.59	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	2.06	2.06	NA NA	NA NA	NA NA	NA NA
Coniferous Wooded Wetlands	Impacted (AFFF Area)	0.00	0.00	NA NA	NA NA	NA NA	NA NA
	Available	289.14	289.14	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	0.00	NA NA	NA NA	NA NA	NA NA
Atlantic White Cedar Wetlands	Impacted (AFFF Area)	0.00	0.00	NA NA	NA NA	NA NA	NA NA
	Available	109.14	109.14	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	3.34	3.34	NA NA	NA NA	NA NA	3.34
Deciduous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	0.00	NA NA	NA NA	NA NA	0.00
	Available	83.03	83.03	NA NA	NA NA	NA NA	83.03
	Impacted (AFFF Drainage)	0.00	53.03	NA NA	NA NA	NA NA	0.00
Coniferous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00		NA NA	NA NA	NA NA	0.00
	Available	51.98		NA NA	NA NA	NA NA	51.98
	Impacted (AFFF Drainage)	0.65	0.65	NA NA	NA NA	NA NA	0.65
	impacted (Altri Diamage)	0.05					
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	0.00	NA	NA	NA	0.00

Feature Description	Habitat Status	Area (Ac)	Podilymbus podiceps	Pooecetes gramineus	Strix varia	Atrytone arogos arogos	Bolaria selene myrina
	Impacted (AFFF Drainage)	0.00	0.00	NA	NA	NA	0.00
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	0.00	NA	NA	NA	0.00
	Available	63.10	63.10	NA	NA	NA	63.10
	Impacted (AFFF Drainage)	0.00	0.00	NA	NA	NA	0.00
Herbaceous Wetlands	Impacted (AFFF Area)	0.00	0.00	NA	NA	NA	0.00
	Available	21.06	21.06	NA	NA	NA N	21.06
	Impacted (AFFF Drainage)	1.44	NA	NA	1.44	NA	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	NA
	Available	47.03	NA	NA	47.03	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	0.00	NA	NA
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	NA
	Available	78.97	NA	NA	78.97	NA	NA
	Impacted (AFFF Drainage)	0.51	NA	NA	NA	NA	NA
Disturbed Wetlands (Modified)	Impacted (AFFF Area)	0.28	NA	NA	NA	NA	NA
	Available	32.13	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Transitional Areas	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	14.34	NA	NA	NA	NA N	NA
	Impacted (AFFF Drainage)	0.00	NA	0.00	NA	NA	NA
Undifferentiated Barren Lands	Impacted (AFFF Area)	0.00	NA	0.00	NA	NA	NA
	Available	34.78	NA	34.78	NA	NA	NA
	Impacted (AFFF Drainage)	0.38	0.38	NA	NA	NA	NA
Phragmites Dominant Interior Wetlands	Impacted (AFFF Area)	0.00	0.00	NA	NA	NA	NA
	Available	7.23	7.23	NA	NA	NA	NA
	Species Habitat Impacted, AF	FF Drainage (ac)	9.07	0.30	63.57	0.30	3.99
	Species Habitat Impacted	d, AFFF Area (ac)	0.00	0.06	0.09	0.06	0.00
	Species Habitat Avail	able Overall (ac)	788.93	263.85	2,933.03	128.23	279.31
	Percent Impacted Over Available	e, AFFF Drainage	1.1501%	0.1148%	2.1674%	0.2363%	1.4276%
	Percent Impacted Over Avai	lable, AFFF Area	0.0000%	0.0228%	0.0031%	0.0469%	0.0000%
	Percent Availabl	le, Not Impacted	98.8499%	99.8624%	97.8294%	99.7168%	98.5724%

Feature Description	Habitat Status	Area (Ac)	Callophrys irus	Lynx rufus	Clemmys insculpta	Clemmys muhlenbergii	Crotalus horridus
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Cemetery	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, High Density or Multiple Dwelling	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	20.73	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Rural, Single Unit	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Low Density	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Residential, Single Unit, Medium Density	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
, , , ,	Available	0.00	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Commercial/Services	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	2.62	NA	NA	NA NA	NA	NA
	Impacted (AFFF Drainage)	72.52	NA NA	NA NA	NA NA	NA NA	NA NA
Military Installations	Impacted (AFFF Area)	22.07	NA	NA NA	NA	NA	NA
,	Available	903.70	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA NA	NA NA	NA NA	NA NA
Transportation/Communication/Utilities	Impacted (AFFF Area)	0.00	NA	NA NA	NA	NA NA	NA NA
Transportation, communication, others	Available	2.43	NA	NA NA	NA	NA	NA NA
	Impacted (AFFF Drainage)	0.00	0.00	NA NA	NA NA	NA NA	NA NA
Upland Right-of-Way Undeveloped	Impacted (AFFF Area)	0.00	0.00	NA NA	NA NA	NA NA	NA NA
opiana Right-or-way ondeveloped	Available	3.42	3.42	NA NA	NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA	NA NA	NA NA	NA NA	NA NA
Stormwater Basin	Impacted (AFFF Area)	0.00	NA	NA NA	NA	NA NA	NA NA
otoriiiwater busiii	Available	23.49	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	67.20	NA NA	NA NA	NA NA	NA NA	NA NA
Other Urban or Built-up Land	Impacted (AFFF Area)	8.22	NA NA	NA NA	NA NA	NA NA	NA NA
other orban or bunt-up tand	Available	1,729.81	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.04	NA NA	NA NA	NA NA	NA NA	NA NA
Managed Wetland in Maintained Lawn Greenspace	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
I wanaged wedand in Manitained Lawn Greenspace	Available	10.38	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)		NA NA	NA NA	NA NA	NA NA	NA NA
Athletic Fields (Schools)		0.00 0.00					
Athletic Fields (Schools)	Impacted (AFFF Area)		NA NA	NA NA	NA NA	NA NA	NA NA
	Available	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Recreational Land	Impacted (AFFF Drainage)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Recreational Land	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	82.29	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA NA	NA NA	NA
Agricultural Wetlands (Modified)	Impacted (AFFF Area)	0.00	NA	NA	NA NA	NA 	NA
	Available	0.00	NA NA	NA 0.00	NA NA	NA NA	NA
	Impacted (AFFF Drainage)	0.00	NA	0.00	NA	NA	NA
Cropland and Pastureland	Impacted (AFFF Area)	0.00	NA	0.00	NA 	NA 	NA 
	Available	0.00	NA	0.00	NA	NA 	NA 
	Impacted (AFFF Drainage)	0.00	NA	NA	NA 	NA 	NA 
Deciduous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	0.73	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.01	NA	NA	NA	NA	NA
Deciduous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	NA	NA	NA	NA	NA
	Available	146.32	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	0.00	NA	NA	0.00
Coniferous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	0.00	NA	NA	0.00
	Available	184.15	NA	184.15	NA	NA	184.15

Feature Description	Habitat Status	Area (Ac)	Callophrys irus	Lynx rufus	Clemmys insculpta	Clemmys muhlenbergii	Crotalus horridus
	Impacted (AFFF Drainage)	45.67	NA	45.67	NA	NA	45.67
Coniferous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	NA	0.01	NA	NA	0.01
	Available	1,381.20	NA	1,381.20	NA	NA	1,381.20
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Plantation	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	104.26	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	0.00	NA	NA	0.00
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)		0.00	NA	0.00	NA	NA	0.00
	Available	10.91	NA	10.91	NA	NA	10.91
	Impacted (AFFF Drainage)	15.12	NA	15.12	NA	NA	15.12
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted (AFFF Area)	0.01	NA	0.01	NA	NA	0.01
	Available	904.40	NA	904.40	NA	NA	904.40
	Impacted (AFFF Drainage)	0.00	NA	0.00	NA	NA	0.00
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	0.00	NA	NA NA	0.00
,	Available	14.62	NA	14.62	NA	NA	14.62
	Impacted (AFFF Drainage)	1.32	NA NA	1.32	NA NA	NA NA	1.32
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted (AFFF Area)	0.06	NA NA	0.06	NA NA	NA NA	0.06
	Available	375.12	NA NA	375.12	NA NA	NA NA	375.12
	Impacted (AFFF Drainage)	0.30	0.30	NA	NA NA	NA NA	NA
Old Field (< 25% Brush Covered)	Impacted (AFFF Area)	0.06	0.06	NA NA	NA	NA NA	NA NA
out tion ( 125/0 brash covered)	Available	124.81	124.81	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA	NA NA	NA NA	NA NA	NA NA
Deciduous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
Decladous Brushi, Shirabiana	Available	1.15	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	5.44	NA NA	NA NA	NA NA	NA NA	NA NA
Coniferous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	200.94	NA NA	NA NA	NA NA	NA NA	NA NA
	Impacted (AFFF Drainage)	3.82	NA NA	NA NA	NA NA	NA NA	NA NA
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted (AFFF Area)	0.10	NA NA	NA NA	NA NA	NA NA	NA NA
ivilked Deciduous/Colliferous Brush/Shrubland	Available	80.91					
	Impacted (AFFF Drainage)	4.45	NA NA	NA NA	NA 4.45	NA NA	NA NA
Streams and Canals		0.00		NA NA	0.00		
Streams and Canais	Impacted (AFFF Area)		NA			NA NA	NA NA
	Available	67.79	NA NA	NA NA	67.79	NA NA	NA NA
Noticed Lakes	Impacted (AFFF Drainage)	0.00	NA	NA NA	0.00	NA NA	NA NA
Natural Lakes	Impacted (AFFF Area)	0.00	NA	NA NA	0.00	NA NA	NA NA
	Available	0.95	NA NA	NA NA	0.95	NA NA	NA NA
Austrial Labora	Impacted (AFFF Drainage)	0.07	NA	NA	NA NA	NA NA	NA NA
Artificial Lakes	Impacted (AFFF Area)	0.00	NA NA	NA NA	NA NA	NA NA	NA NA
	Available	91.55	NA NA	NA NA	NA 2.50	NA NA	NA NA
Daviduana Waadad Wallanda	Impacted (AFFF Drainage)	2.58	NA	NA NA	2.58	NA NA	NA NA
Deciduous Wooded Wetlands	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA NA	NA NA
	Available	63.59	NA	NA	63.59	NA NA	NA 2.26
out our work have to	Impacted (AFFF Drainage)	2.06	NA	NA	2.06	NA NA	2.06
Coniferous Wooded Wetlands	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA 	0.00
	Available	289.14	NA	NA	289.14	NA 	289.14
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA 	0.00
Atlantic White Cedar Wetlands	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	0.00
	Available	109.14	NA	NA	NA	NA	109.14
	Impacted (AFFF Drainage)	3.34	NA	NA	3.34	3.34	NA
Deciduous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	NA	NA	0.00	0.00	NA
	Available	83.03	NA	NA	83.03	83.03	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	0.00	0.00	NA
Coniferous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	NA	NA	0.00	0.00	NA
	Available	51.98	NA	NA	51.98	51.98	NA
	Impacted (AFFF Drainage)	0.65	NA	NA	0.65	0.65	NA
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	0.00	0.00	NA
	Available	60.14	NA	NA	60.14	60.14	NA

Feature Description	Habitat Status	Area (Ac)	Callophrys irus	Lynx rufus	Clemmys insculpta	Clemmys muhlenbergii	Crotalus horridus
	Impacted (AFFF Drainage)	0.00	NA	NA	0.00	0.00	NA
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	0.00	0.00	NA
	Available	63.10	NA	NA	63.10	63.10	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	0.00	0.00	NA
Herbaceous Wetlands	Impacted (AFFF Area)	0.00	NA	NA	0.00	0.00	NA
	Available	21.06	NA	NA	21.06	21.06	NA
	Impacted (AFFF Drainage)	1.44	NA	NA	1.44	NA	NA
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	NA
	Available	47.03	NA	NA	47.03	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	0.00	NA	0.00
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	NA	NA	0.00	NA	0.00
	Available	78.97	NA	NA	78.97	NA	78.97
	Impacted (AFFF Drainage)	0.51	NA	NA	NA	NA	NA
Disturbed Wetlands (Modified)	Impacted (AFFF Area)	0.28	NA	NA	NA	NA	NA
	Available	32.13	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Transitional Areas	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	14.34	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA	NA	NA	NA
Undifferentiated Barren Lands	Impacted (AFFF Area)	0.00	NA	NA	NA	NA	NA
	Available	34.78	NA	NA	NA	NA	NA
	Impacted (AFFF Drainage)	0.38	NA	NA	0.38	0.38	NA
Phragmites Dominant Interior Wetlands	Impacted (AFFF Area)	0.00	NA	NA	0.00	0.00	NA
	Available	7.23	NA	NA	7.23	7.23	NA
	Species Habitat Impacted, AFFF Drainage (ac)		0.30	62.12	14.89	4.37	64.18
	Species Habitat Impacted, AFFF Area (ac) Species Habitat Available Overall (ac)			0.08	0.00	0.00	0.08
				2,870.40	834.01	286.54	3,347.65
	Percent Impacted Over Available,	AFFF Drainage	0.2363%	2.1641%	1.7856%	1.5246%	1.9171%
	Percent Impacted Over Availal	ble, AFFF Area	0.0469%	0.0027%	0.0000%	0.0000%	0.0023%
	Percent Available,	Not Impacted	99.7168%	97.8332%	98.2144%	98.4754%	98.0806%

Feature Description	Habitat Status	Area (Ac)	Elaphe guttata	Pituophis m melanoleucus
	Impacted (AFFF Drainage)	0.00	NA	NA
Cemetery	Impacted (AFFF Area)	0.00	NA	NA
	Available	0.00	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA
Residential, High Density or Multiple Dwelling	Impacted (AFFF Area)	0.00	NA	NA
	Available	20.73	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA
Residential, Rural, Single Unit	Impacted (AFFF Area)	0.00	NA	NA
	Available	0.00	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA
Residential, Single Unit, Low Density	Impacted (AFFF Area)	0.00	NA	NA
	Available	0.00	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA
Residential, Single Unit, Medium Density	Impacted (AFFF Area)	0.00	NA	NA
	Available	0.00	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA
Commercial/Services	Impacted (AFFF Area)	0.00	NA	NA
	Available	2.62	NA	NA
	Impacted (AFFF Drainage)	72.52	NA	NA
Military Installations	Impacted (AFFF Area)	22.07	NA	NA
•	Available	903.70	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA
Transportation/Communication/Utilities	Impacted (AFFF Area)	0.00	NA	NA
•	Available	2.43	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA
Upland Right-of-Way Undeveloped	Impacted (AFFF Area)	0.00	NA	NA
	Available	3.42	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA
Stormwater Basin	Impacted (AFFF Area)	0.00	NA	NA
	Available	23.49	NA	NA
	Impacted (AFFF Drainage)	67.20	NA	NA
Other Urban or Built-up Land	Impacted (AFFF Area)	8.22	NA	NA
·	Available	1,729.81	NA	NA
	Impacted (AFFF Drainage)	0.04	NA	NA
Managed Wetland in Maintained Lawn Greenspace	Impacted (AFFF Area)	0.00	NA	NA
	Available	10.38	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	NA
Athletic Fields (Schools)	Impacted (AFFF Area)	0.00	NA NA	NA NA
,	Available	0.00	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA
Recreational Land	Impacted (AFFF Area)	0.00	NA NA	NA NA
	Available	82.29	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA
Agricultural Wetlands (Modified)	Impacted (AFFF Area)	0.00	NA NA	NA NA
(	Available	0.00	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA
Cropland and Pastureland	Impacted (AFFF Area)	0.00	NA NA	NA NA
שוטקימווא מווע ו עסנעונועווע	Available	0.00	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	0.00
Deciduous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA NA	0.00
Decidadas i orest (10-30/0 Crown Closure)	Available	0.73	NA NA	0.00
			NA NA	0.73 NA
Deciduous Forest />E00/ Crown Clasura)	Impacted (AFFF Drainage)	0.01		
Deciduous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	NA NA	NA NA
Г	Available	146.32	NA NA	NA 0.00
Coniference Forest /10 FOX/ Consum Classes	Impacted (AFFF Drainage)	0.00	NA NA	0.00
Coniferous Forest (10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA NA	0.00
	Available	184.15	NA	184.15

Feature Description	Habitat Status	Area (Ac)	Elaphe guttata	Pituophis m melanoleucus
	Impacted (AFFF Drainage)	45.67	45.67	NA
Coniferous Forest (>50% Crown Closure)	Impacted (AFFF Area)	0.01	0.01	NA
	Available	1,381.20	1,381.20	NA
	Impacted (AFFF Drainage)	0.00	NA	NA
Plantation	Impacted (AFFF Area)	0.00	NA	NA
	Available	104.26	NA	NA
	Impacted (AFFF Drainage)	0.00	NA	0.00
Mixed Forest (> 50% Coniferous with 10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	0.00
	Available	10.91	NA	10.91
	Impacted (AFFF Drainage)	15.12	15.12	NA
Mixed Forest (> 50% Coniferous with >50% Crown Closure)	Impacted (AFFF Area)	0.01	0.01	NA
	Available	904.40	904.40	NA
	Impacted (AFFF Drainage)	0.00	NA	0.00
Mixed Forest (>50% Deciduous with 10-50% Crown Closure)	Impacted (AFFF Area)	0.00	NA	0.00
	Available	14.62	NA	14.62
	Impacted (AFFF Drainage)	1.32	NA	NA
Mixed Forest (>50% Deciduous with >50% Crown Closure)	Impacted (AFFF Area)	0.06	NA	NA
	Available	375.12	NA	NA
	Impacted (AFFF Drainage)	0.30	NA NA	0.30
Old Field (< 25% Brush Covered)	Impacted (AFFF Area)	0.06	NA	0.06
	Available	124.81	NA	124.81
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA
Deciduous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA	NA
	Available	1.15	NA NA	NA
	Impacted (AFFF Drainage)	5.44	NA NA	NA NA
Coniferous Brush/Shrubland	Impacted (AFFF Area)	0.00	NA NA	NA NA
Connerous brushy shrubland	Available	200.94	NA NA	NA NA
	Impacted (AFFF Drainage)	3.82	NA NA	NA NA
Mixed Deciduous/Coniferous Brush/Shrubland	Impacted (AFFF Area)	0.10	NA NA	NA NA
wince Decideday conferous brasily sin abiana	Available	80.91	NA NA	NA NA
	Impacted (AFFF Drainage)	4.45	NA NA	NA NA
Streams and Canals	Impacted (AFFF Area)	0.00	NA NA	NA NA
Streams and Canais	Available	67.79	NA NA	NA NA
	Impacted (AFFF Drainage)	0.00	NA NA	NA NA
Natural Lakes	, , ,			
Natural Lakes	Impacted (AFFF Area) Available	0.00	NA NA	NA NA
		0.95	NA NA	NA NA
Autiliain Lalena	Impacted (AFFF Drainage)	0.07	NA NA	NA NA
Artificial Lakes	Impacted (AFFF Area)	0.00	NA NA	NA NA
	Available	91.55	NA NA	NA NA
Desiderary Weeded Wetlende	Impacted (AFFF Drainage)	2.58	NA NA	NA NA
Deciduous Wooded Wetlands	Impacted (AFFF Area)	0.00	NA NA	NA NA
	Available	63.59	NA 2.00	NA NA
Coniference Monday d Matterials	Impacted (AFFF Drainage)	2.06	2.06	NA NA
Coniferous Wooded Wetlands	Impacted (AFFF Area)	0.00	0.00	NA NA
	Available	289.14	289.14	NA NA
	Impacted (AFFF Drainage)	0.00	0.00	NA NA
Atlantic White Cedar Wetlands	Impacted (AFFF Area)	0.00	0.00	NA
	Available	109.14	109.14	NA
	Impacted (AFFF Drainage)	3.34	NA	NA
Deciduous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	NA	NA
	Available	83.03	NA	NA
	Impacted (AFFF Drainage)	0.00	0.00	NA
Coniferous Scrub/Shrub Wetlands	Impacted (AFFF Area)	0.00	0.00	NA
	Available	51.98	51.98	NA
	Impacted (AFFF Drainage)	0.65	NA	NA
Mixed Scrub/Shrub Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	NA	NA
	Available	60.14	NA	NA

Feature Description	Habitat Status	Area (Ac)	Elaphe guttata	Pituophis m melanoleucus		
	Impacted (AFFF Drainage)	0.00	0.00	NA		
Mixed Scrub/Shrub Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	0.00	NA		
	Available	63.10	63.10	NA		
	Impacted (AFFF Drainage)	0.00	NA	NA		
Herbaceous Wetlands	Impacted (AFFF Area)	0.00	NA	NA		
	Available	21.06	NA	NA		
	Impacted (AFFF Drainage)	1.44	NA	NA		
Mixed Wooded Wetlands (Deciduous Dom.)	Impacted (AFFF Area)	0.00	NA	NA		
	Available	47.03	NA	NA		
	Impacted (AFFF Drainage)	0.00	0.00	NA		
Mixed Wooded Wetlands (Coniferous Dom.)	Impacted (AFFF Area)	0.00	0.00	NA		
	Available	78.97	78.97	NA		
	Impacted (AFFF Drainage)	0.51	NA	NA		
Disturbed Wetlands (Modified)	Impacted (AFFF Area)	0.28	NA	NA		
	Available	32.13	NA	NA		
	Impacted (AFFF Drainage)	0.00	NA	NA		
Transitional Areas	Impacted (AFFF Area)	0.00	NA	NA		
	Available	14.34	NA	NA		
	Impacted (AFFF Drainage)	0.00	0.00	0.00		
Undifferentiated Barren Lands	Impacted (AFFF Area)	0.00	0.00	0.00		
	Available	34.78	34.78	34.78		
	Impacted (AFFF Drainage)	0.38	NA	NA		
Phragmites Dominant Interior Wetlands	Impacted (AFFF Area)	0.00	NA	NA		
	Available	7.23	NA	NA		
	Species Habitat Impacted, A	FFF Drainage (ac)	62.86	0.30		
	Species Habitat Impacted, AFFF Area (ac)					
	2,912.71	370.00				
	Percent Impacted Over Availab	le, AFFF Drainage	2.1581%	0.0819%		
	Percent Impacted Over Ava	· ·		0.0163%		
	Percent Availab	ole, Not Impacted	97.8414%	99.9018%		

ac = acres

Table B-4. Spatial Exposure Potential of Protected Species at Naval Air Station Patuxent River

Feature Description	Habitat Status	Area (ac)	Gastrophryne carolinensis	Accipiter gentilis	Ammodramus henslowii	Asio flammeus	Bartramia longicauda	Charadrius melodus	Cistothorus platensis
Desidential	Impacted	0.00	NA	NA	NA	NA	NA	NA	NA
Residential	Available	19.80	NA	NA	NA	NA	NA	NA	NA
Institutional	Impacted	21.68	NA	NA	NA	NA	NA	NA	NA
institutional	Available	723.50	NA	NA	NA	NA	NA	NA	NA
Transportation	Impacted	10.13	NA	NA	NA	NA	NA	NA	NA
Transportation	Available	503.97	NA	NA	NA	NA	NA	NA	NA
Open Urban Land	Impacted	37.57	NA	NA	37.57	37.57	37.57	NA	NA
Open Orban Land	Available	1,917.60	NA	NA	1,917.60	1,917.60	1,917.60	NA	NA
A anioultuno	Impacted	0.00	0.00	0.00	NA	0.00	0.00	NA	NA
Agriculture	Available	149.09	149.09	149.09	NA	149.09	149.09	NA	NA
Desidueus Ferest	Impacted	0.02	0.02	0.02	NA	NA	NA	NA	NA
Deciduous Forest	Available	633.81	633.81	633.81	NA	NA	NA	NA	NA
Cycumus on Councit	Impacted	8.90	8.90	8.90	NA	NA	NA	NA	NA
Evergreen Forest	Available	99.75	99.75	99.75	NA	NA	NA	NA	NA
Mixed Forest	Impacted	24.78	24.78	24.78	NA	NA	NA	NA	NA
	Available	1,323.13	1,323.13	1,323.13	NA	NA	NA	NA	NA
Old Field	Impacted	5.26	5.26	5.26	5.26	5.26	5.26	NA	NA
Old Field	Available	221.84	221.84	221.84	221.84	221.84	221.84	NA	NA
Brush	Impacted	5.26	NA	NA	5.26	5.26	5.26	NA	5.26
Drusii	Available	127.32	NA	NA	127.32	127.32	127.32	NA	127.32
Nontidal Water	Impacted	0.98	0.98	NA	NA	NA	NA	0.98	0.98
Nontidal Water	Available	42.79	42.79	NA	NA	NA	NA	42.79	42.79
Tidal Water	Impacted	0.00	NA	NA	NA	NA	NA	0.00	NA
iluai water	Available	47.41	NA	NA	NA	NA	NA	47.41	NA
Emergent Wetland	Impacted	0.00	0.00	0.00	0.00	0.00	NA	0.00	0.00
Emergent Wetland	Available	109.79	109.79	109.79	109.79	109.79	NA	109.79	109.79
Forested Wetland	Impacted	5.53	5.53	5.53	NA	NA	NA	NA	NA
rofesteu wetialiu	Available	270.12	270.12	270.12	NA	NA	NA	NA	NA
Sandy Shoreline	Impacted	0.00	NA	NA	NA	0.00	NA	0.00	NA
Sandy Shoreline	Available	18.28	NA	NA	NA	18.28	NA	18.28	NA
	Species Habitat	Impacted (ac)	45.46	44.48	48.09	48.09	48.09	0.98	6.24
	Species Habitat Availab	le Overall (ac)	2,850.32	2,807.53	2,376.55	2,543.92	2,415.85	218.26	279.89
	Percent Impacted C	Over Available	1.59%	1.58%	2.02%	1.89%	1.99%	0.45%	2.23%
	Percent Available,	Not Impacted	98.41%	98.42%	97.98%	98.11%	98.01%	99.55%	97.77%

ac = acres

Feature Description	Habitat Status	Area (ac)	Contopus cooperi	Gelochelidon nilotica	Geothlypis philadelphia	Lanius ludovicianus	Rynchops niger	Setophaga fusca	Sternula antillarum
Residential	Impacted	0.00	NA	NA	NA	NA	NA	NA	NA
Residential	Available	19.80	NA	NA	NA	NA	NA	NA	NA
Institutional	Impacted	21.68	NA	NA	NA	NA	NA	NA	NA
ilistitutioliai	Available	723.50	NA	NA	NA	NA	NA	NA	NA
Transportation	Impacted	10.13	NA	NA	NA	NA	NA	NA	NA
Transportation	Available	503.97	NA	NA	NA	NA	NA	NA	NA
Open Urban Land	Impacted	37.57	NA	NA	NA	37.57	NA	NA	NA
Open Orban Land	Available	1,917.60	NA	NA	NA	1,917.60	NA	NA	NA
Agriculture	Impacted	0.00	NA	0.00	NA	NA	NA	NA	NA
Agriculture	Available	149.09	NA	149.09	NA	NA	NA	NA	NA
Deciduous Forest	Impacted	0.02	NA	NA	0.02	0.02	NA	0.02	NA
Deciduous Forest	Available	633.81	NA	NA	633.81	633.81	NA	633.81	NA
Currence Conset	Impacted	8.90	8.90	NA	8.90	8.90	NA	8.90	NA
Evergreen Forest	Available	99.75	99.75	NA	99.75	99.75	NA	99.75	NA
Mixed Forest	Impacted	24.78	24.78	NA	24.78	NA	NA	24.78	NA
	Available	1,323.13	1,323.13	NA	1,323.13	NA	NA	1,323.13	NA
Old Field	Impacted	5.26	NA	NA	5.26	5.26	NA	NA	NA
Old Fleid	Available	221.84	NA	NA	221.84	221.84	NA	NA	NA
Brush	Impacted	5.26	5.26	NA	5.26	5.26	NA	NA	NA
Drusii	Available	127.32	127.32	NA	127.32	127.32	NA	NA	NA
Nontidal Water	Impacted	0.98	NA	NA	NA	NA	NA	NA	0.98
Nontidal Water	Available	42.79	NA	NA	NA	NA	NA	NA	42.79
Tidal Water	Impacted	0.00	NA	0.00	NA	NA	0.00	NA	0.00
ildai water	Available	47.41	NA	47.41	NA	NA	47.41	NA	47.41
Emergent Wetland	Impacted	0.00	NA	NA	NA	NA	NA	NA	NA
Emergent Wetland	Available	109.79	NA	NA	NA	NA	NA	NA	NA
Forested Wetland	Impacted	5.53	5.53	NA	NA	NA	NA	5.53	NA
rorestea wetiana	Available	270.12	270.12	NA	NA	NA	NA	270.12	NA
Sandy Shoreline	Impacted	0.00	NA	0.00	NA	NA	0.00	NA	0.00
Sandy Shoreline	Available	18.28	NA	18.28	NA	NA	18.28	NA	18.28
	Species Habitat	t Impacted (ac)	44.46	0.00	44.21	57.01	0.00	39.22	0.98
	Species Habitat Availal	ble Overall (ac)	1,820.32	214.78	2,405.85	3,000.32	65.69	2,326.81	108.47
	Percent Impacted	Over Available	2.44%	0.00%	1.84%	1.90%	0.00%	1.69%	0.90%
	Percent Available	, Not Impacted	97.56%	100.00%	98.16%	98.10%	100.00%	98.31%	99.10%

ac = acres

Feature Description	Habitat Status	Area (ac)	Thalasseus maximus	Acipenser oxyrhynchus	Centrarchus macropterus	Notropis chalybaeus	Callophrys irus	Cicindela d dorsalis	Eubalaeana glacialis
Residential	Impacted	0.00	NA	NA	NA	NA	NA	NA	NA
Residential	Available	19.80	NA	NA	NA	NA	NA	NA	NA
Institutional	Impacted	21.68	NA	NA	NA	NA	NA	NA	NA
mstitutional	Available	723.50	NA	NA	NA	NA	NA	NA	NA
Transportation	Impacted	10.13	NA	NA	NA	NA	NA	NA	NA
Transportation	Available	503.97	NA	NA	NA	NA	NA	NA	NA
Open Urban Land	Impacted	37.57	NA	NA	NA	NA	37.57	NA	NA
Open Orban Land	Available	1,917.60	NA	NA	NA	NA	1,917.60	NA	NA
Agriculture	Impacted	0.00	NA	NA	NA	NA	NA	NA	NA
Agriculture	Available	149.09	NA	NA	NA	NA	NA	NA	NA
Deciduous Forest	Impacted	0.02	NA	NA	NA	NA	NA	NA	NAA
Deciduous Forest	Available	633.81	NA	NA	NA	NA	NA	NA	NA
Cyangnasa Canast	Impacted	8.90	NA	NA	NA	NA	NA	NA	NA
Evergreen Forest	Available	99.75	NA	NA	NA	NA	NA	NA	NA
Mixed Forest	Impacted	24.78	NA	NA	NA	NA	NA	NA	NA
	Available	1,323.13	NA	NA	NA	NA	NA	NA	NA
Old Field	Impacted	5.26	NA	NA	NA	NA	5.26	5.26	NA
Old Field	Available	221.84	NA	NA	NA	NA	221.84	221.84	NA
Brush	Impacted	5.26	NA	NA	NA	NA	5.26	NA	NA
DI USII	Available	127.32	NA	NA	NA	NA	127.32	NA	NA
Nontidal Water	Impacted	0.98	NA	NA	0.98	0.98	NA	NA	NA
Nontidal Water	Available	42.79	NA	NA	42.79	42.79	NA	NA	NA
Tidal Water	Impacted	0.00	0.00	0.00	NA	NA	NA	NA	0.00
iluai water	Available	47.41	47.41	47.41	NA	NA	NA	NA	47.41
Emergent Wetland	Impacted	0.00	NA	NA	NA	NA	NA	NA	NA
Lineigent Wetland	Available	109.79	NA	NA	NA	NA	NA	NA	NA
Forested Wetland	Impacted	5.53	NA	NA	NA	NA	NA	NA	NA
rolesteu Wetlaliu	Available	270.12	NA	NA	NA	NA	NA	NA	NA
Sandy Shoreline	Impacted	0.00	0.00	NA	NA	NA	NA	0.00	NA
Sandy Shoreline	Available	18.28	18.28	NA	NA	NA	NA	18.28	NA
	Species Habita	t Impacted (ac)	0.00	0.00	0.98	0.98	48.09	5.26	0.00
	Species Habitat Availal	ble Overall (ac)	65.69	47.41	42.79	42.79	2,266.76	240.12	47.41
	Percent Impacted	Over Available	0.00%	0.00%	2.29%	2.29%	2.12%	2.19%	0.00%
	Percent Available	, Not Impacted	100.00%	100.00%	97.71%	97.71%	97.88%	97.81%	100.00%

ac = acres

Feature Description	Habitat Status	Area (ac)	Megaptera novaeangliae	Dermochelys coriacea	Caretta caretta	Lepidochelys kempii	Haliaeetus leucocephalus
Residential	Impacted	0.00	NA	NA	NA	NA	NA
Residential	Available	19.80	NA	NA	NA	NA	NA
Institutional	Impacted	21.68	NA	NA	NA	NA	NA
ilistitutioliai	Available	723.50	NA	NA	NA	NA	NA
Transportation	Impacted	10.13	NA	NA	NA	NA	NA
Transportation	Available	503.97	NA	NA	NA	NA	NA
Open Urban Land	Impacted	37.57	NA	NA	NA	NA	NA
Open Orban Land	Available	1,917.60	NA	NA	NA	NA	NA
Agriculture	Impacted	0.00	NA	NA	NA	NA	NA
Agriculture	Available	149.09	NA	NA	NA	NA	NA
Deciduous Forest	Impacted	0.02	NA	NA	NA	NA	0.02
Deciduous Forest	Available	633.81	NA	NA	NA	NA	633.81
Evergreen Forest	Impacted	8.90	NA	NA	NA	NA	8.90
Evergreen Forest	Available	99.75	NA	NA	NA	NA	99.75
Mixed Forest	Impacted	24.78	NA	NA	NA	NA	24.78
	Available	1,323.13	NA	NA	NA	NA	1,323.13
Old Field	Impacted	5.26	NA	NA	NA	NA	5.26
Old Field	Available	221.84	NA	NA	NA	NA	221.84
Brush	Impacted	5.26	NA	NA	NA	NA	NA
Diusii	Available	127.32	NA	NA	NA	NA	NA
Nontidal Water	Impacted	0.98	NA	NA	NA	NA	0.98
Nontidal Water	Available	42.79	NA	NA	NA	NA	42.79
Tidal Water	Impacted	0.00	0.00	0.00	0.00	0.00	0.00
iluai watei	Available	47.41	47.41	47.41	47.41	47.41	47.41
Emergent Wetland	Impacted	0.00	NA	NA	NA	NA	0.00
Lineigent Wetland	Available	109.79	NA	NA	NA	NA	109.79
Forested Wetland	Impacted	5.53	NA	NA	NA	NA	NA
rorested wetland	Available	270.12	NA	NA	NA	NA	NA
Sandy Shoreline	Impacted	0.00	NA	0.00	0.00	0.00	NA
Sandy Shoreline	Available	18.28	NA	18.28	18.28	18.28	NA
	Species Habitat	Impacted (ac)	0.00	0.00	0.00	0.00	39.93
	Species Habitat Availab	le Overall (ac)	47.41	65.69	65.69	65.69	2,478.52
	Percent Impacted (	Over Available	0.00%	0.00%	0.00%	0.00%	1.61%
	Percent Available,	Not Impacted	100.00%	100.00%	100.00%	100.00%	98.39%

ac = acres

